



# **The Comparative Clinical Effectiveness and Value of Novel Combination Therapies for the Treatment of Patients with Genotype 1 Chronic Hepatitis C Infection**

**A Technology Assessment**

**Draft Report**

**November 17, 2014**

**Completed by:**

**Institute for Clinical and Economic Review**



**AUTHORS:**

Jeffrey A. Tice, MD

Associate Professor of Medicine, Division of General Internal Medicine, Department of Medicine, University of California San Francisco

Daniel A. Ollendorf, PhD

Chief Review Officer, Institute for Clinical and Economic Review

Harinder S. Chahal, PharmD, MSc

WHO Collaborating Centre for Pharmaceutical Research and Science Policy, Department of Clinical Pharmacy, University of California San Francisco

James G. Kahn, MD, MPH

Professor, School of Medicine, Institute for Health Policy Studies, University of California San Francisco

Elliot Marseille, DrPH, MPP

Principal, Health Strategies International

Jed Weissberg, MD, FACP

Senior Fellow, Institute for Clinical and Economic Review

Steven D. Pearson, MD, MSc

President, Institute for Clinical and Economic Review

**DATE OF****PUBLICATION:**

November 17, 2014

We wish to thank Rena Fox, MD, Professor of Clinical Medicine, Division of General Internal Medicine, UCSF, for her contributions as a clinical expert to the modeling work and Sachin Garg, MD, Primary Care Research Fellow, UCSF School of Medicine, for his contributions to the data abstractions for this report.

## About ICER

The Institute for Clinical and Economic Review (ICER) is an independent non-profit research organization that evaluates medical evidence and convenes public deliberative bodies to help stakeholders interpret and apply evidence to improve patient outcomes and control costs. ICER receives funding from government grants, non-profit foundations, health plans, provider groups, and health industry manufacturers. Through all its work, ICER seeks to help create a future in which collaborative efforts to move evidence into action provide the foundation for a more effective, efficient, and just health care system. More information about ICER is available at [www.icer-review.org](http://www.icer-review.org)

## About CTAF

The California Technology Assessment Forum (CTAF) – a core program of ICER – reviews evidence reports and provides a public venue in which the evidence on the effectiveness and value of health care services can be discussed with the input of all stakeholders. CTAF seeks to help patients, clinicians, insurers, and policymakers interpret and use evidence to improve the quality and value of health care.

The CTAF Panel is an independent committee of medical evidence experts from across California, with a mix of practicing clinicians, methodologists, and leaders in patient engagement and advocacy, all of whom meet strict conflict of interest guidelines, who are convened to evaluate evidence and vote on the comparative clinical effectiveness and value of medical interventions. More information about CTAF is available at [www.ctaf.org](http://www.ctaf.org)

# Table of Contents

Abbreviations.....	V
Executive Summary.....	ES1
Introduction .....	1
1. Background .....	3
2. Clinical Guidelines .....	12
3. Coverage Policies .....	14
4. Previous Systematic Reviews and Technology Assessments.....	18
5. Ongoing Studies .....	19
6. Evidence Review (Methods & Results) .....	24
7. Model of Clinical and Economic Outcomes of Treatment Strategies for Hepatitis C.....	43
References .....	74
<b>APPENDICES.....</b>	<b>89</b>

## Abbreviations used in this report

AASLD:	American Association for the Study of Liver Diseases
AEs:	Adverse events
AST/ALT:	aspartate aminotransferase/alanine aminotransferase
ASV:	Asunaprevir
CDC:	Centers for Disease Control and Prevention
CHC:	Chronic hepatitis C
CI:	Confidence interval
CMS:	Centers for Medicare & Medicaid Services
CPI:	Consumer Price Index
CTAF:	California Technology Assessment Forum
DARE:	Database of Abstracts of Reviews of Effects
DAA:	Direct-acting antiviral agent
DCV:	Daclatasvir
DR:	Discontinuation rate
FDA:	US Food and Drug Administration
HCC:	Hepatocellular carcinoma
HCV:	Hepatitis C virus
HR:	Hazard ratio
ICER:	Incremental cost-effectiveness ratio
IFN	Interferon
LDV:	Ledipasvir
NDA:	New drug application
NR:	Not reported
NS:	Not significant
OR:	Odds ratio
P:	Pegylated interferon
PBO:	Placebo
PMPM:	Per-member per-month
PR:	Pegylated interferon plus ribavirin
Q8:	Taken every 8 hours
QALY:	Quality-adjusted life year
R:	Ribavirin
RCT:	Randomized Controlled Trial
SMV:	Simeprevir
SOF:	Sofosbuvir
SVR:	Sustained virologic response
SVR12:	SVR at 12 weeks
US:	United States
WTP:	Willingness-to-pay
3D:	Paritaprevir, ritonavir, ombitasvir, and dasabuvir

## Executive Summary

On December 18, 2014, the California Technology Assessment Forum (CTAF) will discuss the comparative clinical effectiveness and value of new interferon-free combinations of direct-acting antiviral (DAA) drugs for the treatment of chronic hepatitis C, genotype 1, the most common genotype in the United States. Our [prior assessment](#) in March 2014 evaluated single DAA drugs used with pegylated interferon and/or ribavirin. Since that review, the FDA has approved two new therapies that each combine two DAAs and do not require the use of either interferon or ribavirin. On October 10, 2014, the FDA approved the combination of ledipasvir/sofosbuvir and on November 5, 2014, the FDA approved the combination of simeprevir/sofosbuvir. At least two more combinations of DAAs are likely to receive FDA approval in the next few months.

Chronic hepatitis C is a common infection that is a major cause of chronic liver disease, liver failure, and hepatocellular carcinoma (HCC), and it is the leading indication for liver transplantation in the Western world.<sup>1</sup> Prior to 2011, the combination of pegylated interferon and ribavirin (PR) was the standard of therapy for the treatment of chronic hepatitis C. Fewer than half of patients with genotype 1 clear the virus from their bloodstream entirely and maintain a sustained virologic response (SVR) 24 weeks after the end of treatment with PR. PR therapy can be difficult, however, as both interferon and ribavirin can produce severe fatigue and body aches, and in some cases, dangerous levels of anemia, neutropenia, and/or thrombocytopenia.<sup>2</sup> The 2011 introduction of first generation DAA protease inhibitors boceprevir and telaprevir resulted in substantially improved SVR rates in many patients when combined with PR. This improvement came with new challenges including significant additional side effects and drug-drug interactions as well as stringent dosing requirements and high pill burdens for patients.<sup>3</sup> In 2013, the FDA approved the second generation of DAAs, simeprevir and sofosbuvir, which in combination with PR increased SVR rates with shorter duration of therapy and fewer adverse events. Since the March 2014 CTAF review on hepatitis C therapies, investigators have published promising results on several interferon-free therapies that combine two or more DAAs.

As highlighted in the prior CTAF assessment, the new drugs are expensive, with new combination therapies costing approximately \$65,000 to \$190,000 per course of therapy depending on treatment duration.<sup>4,5</sup> Because chronic infection with HCV is relatively common, this translates into an enormous potential budget impact for federal, state, and private health insurers. Because of the tension between the potential cost-effectiveness of these new agents (i.e., their “care value”) and their budgetary impact (i.e., health-system value), ICER commissioned a detailed cost-effectiveness model to provide a more robust analysis of the benefits and costs of the new agents for the current assessment.

This assessment addresses the following questions: 1) among patients with genotype 1 hepatitis C infections, what is the comparative clinical effectiveness of combinations of two or more DAAs compared to each other, as well as to single DAA therapy used in combination with interferon and ribavirin in the achievement of SVR as a surrogate for the prevention of longer-term sequelae of chronic liver disease; and 2) what is the comparative value of the new therapies and alternative population treatment strategies (i.e., treat all vs. treat only patients with advanced liver disease). The purpose of this assessment is to help patients, providers, and payers address these important questions and to support dialogue needed for successful action to improve the quality and value of health care for patients with hepatitis C.

This evidence review of treatments for genotype 1 differs from the March 2014 CTAF review in analyzing four clinically relevant subgroups shown in Table ES1 below and derives summary estimates for SVR and discontinuation rates (DR) in each group by treatment regimen listed in Table ES2 below.

**Table ES1: Clinical Subgroups**

Treatment-naïve / non-cirrhotic	Treatment-naïve / cirrhotic
Treatment-experienced / non-cirrhotic	Treatment-experienced / cirrhotic

**Table ES2: Therapies Considered in this Assessment**

Brand Name	Generic Name	Abbreviation	Pharmaceutical Company
<i>FDA-approved comparators from prior review</i>			
Olysio + PR	Simeprevir + PR	SMV + PR	Janssen and Medivir AB
Sovaldi + PR	Sofosbuvir + PR	SOF + PR	Gilead Sciences
Sovaldi + R	Sofosbuvir + R	SOF + R	Gilead Sciences
<i>FDA-approved combinations since prior review</i>			
Olysio + Sovaldi	Simeprevir/sofosbuvir	SMV/SOF	Janssen + Gilead Sciences
Harvoni	Ledipasvir/sofosbuvir	LDV/SOF	Gilead Sciences
<i>Combinations pending FDA approval</i>			
Daklinza + Sovaldi	Daclatasvir + sofosbuvir	DCV/SOF	Bristol-Myers Squibb + Gilead Sciences
3D	Paritaprevir + ritonavir + ombitasvir + dasabuvir	3D	AbbVie

We included all prospective randomized trials and cohorts that reported SVR12 or SVR24 in HCV genotype 1 infected populations. The SVR results for each of the regimens by clinical subgroup are shown in Figure ES1 on page ES4. The evidence on the comparative clinical effectiveness of the newest treatment regimens appears consistent in all four major treatment subgroups. Among treatment-naïve patients without cirrhosis, the SVR12 for simeprevir or sofosbuvir combined with interferon and/or ribavirin is between 75% and 90%. The comparable SVR12 for DAA combination therapy (i.e., SMV/SOF, LDV/SOF, DCV/SOF, 3D) ranges from 95% to 100%. Among treatment-naïve patients with cirrhosis, the SVR12 for primary DAA therapy ranges from 55% to 80% compared to 67% to 95% for DAA combination therapy. For treatment-experienced patients, the SVR12 for older therapy is about 75% for both cirrhotic and non-cirrhotic patients and is 95% to 100% for DAA combination therapy.


Due to the lack of head-to-head trials and the lack of trials with common comparators, it is difficult to know if one of the DAA combination therapies is clearly superior to another. At this time, only two combinations have FDA approval (simeprevir/sofosbuvir, ledipasvir/sofosbuvir). Two of the combinations (simeprevir/sofosbuvir, daclatasvir/sofosbuvir) have been studied in very small samples, and the confidence intervals around the estimates for their SVRs are wide. For the patient population with cirrhosis, the confidence intervals are wide for all four of the new DAA combinations. Furthermore, selection bias may explain some of the differences between the SVRs.

Adverse effects may also impact the choice of therapy, although there were very few discontinuations from therapy in any of the studies due to adverse events, and the rate of serious adverse events was similarly low. When patient characteristics require longer therapy with ribavirin (sofosbuvir + R for 24 weeks, 3D + R for 24 weeks), the adverse event rates are higher, and the rate of significant anemia is higher. Simeprevir also causes photosensitivity and more rashes.

High-quality observational data from real world settings will be essential for evaluating the comparative effectiveness of the combination DAA therapies and to see if the SVR rates achieved in clinical trials are replicated in usual care settings.



Figure ES1: SVR and 95% Confidence Intervals for the Primary DAA Regimens in Four Clinical Subgroups



\* — N values represent the total number of patients in the represented studies for each clinical subgroup

## Care Value Analysis: Cost-Effectiveness Mode

We commissioned a decision-analytic multistate Markov model<sup>125</sup> to determine the cost-effectiveness of six treatment regimens for HCV genotype 1 currently marketed in the US. The model calculated the net costs, health benefits, and incremental cost-effectiveness ratios (ICERs) of these therapies. It is also designed to determine how these ICERs change if treatment is delayed to a more advanced stage of disease as compared to treating people at all disease stages. We thus aimed to address two key policy or program questions with regard to HCV therapy:

- **Comparing regimens:** Which regimens are most cost-effective? Specifically, what is the incremental cost-effectiveness of more expensive and effective regimens?
- **Comparing population treatment strategies:** What is the cost-effectiveness of treating all individuals, as compared with waiting to treat at more advanced disease stages?

The model produced lifetime discounted quality-adjusted life years (QALYs) and costs to calculate ICERs. Costs, QALYs gained, incremental costs, and incremental QALYs were calculated for each regimen in comparison with the next least costly regimen. The ICER for each regimen's "treat all" strategy also was calculated against "treat at F3, F4" (i.e., treat only when patients have severe fibrosis or cirrhosis) in order to assess the cost-effectiveness of a global treatment approach versus a prioritized one.

In the base-case analysis, we found that LDV/SOF regimens for treatment-naïve and treatment-experienced patients were very cost-effective, producing ICERs  $\leq$ \$20,000 per QALY gained regardless of the comparison (i.e., PR alone vs. next-least costly alternative, treat all vs. treat at F3/F4, weighted estimates for a combined treatment-naïve and treatment-experienced cohort).

Our analysis also found that, while treating patients at all fibrosis stages was more expensive in comparison to waiting to treat until patients reached F3 or F4, it was also more effective. For example, treating all naïve patients with LDV/SOF 8/12 (according to viral load and fibrosis stage) or LDV/SOF 12 (all patients get 12 weeks of therapy) or PR alone produced ICERs  $<$ \$40,000 per QALY gained in comparison to treating only at F3/F4. Among treatment-experienced patients, differences in effectiveness were more pronounced, with more than two years of quality-adjusted life expectancy gained for single DAA sofosbuvir-based regimens relative to PR alone (generating ICERs of \$10,000-\$20,000 per QALY gained). Comparisons of the "treat all" vs. "treat at F3, F4" approaches in the treatment-experienced subgroup generated more costs (in part because single DAA sofosbuvir-based regimens are longer) but still produced estimates of cost-effectiveness of  $\sim$ \$50,000 per QALY gained.

## Health System Value Analysis

We assessed the potential budgetary impact of new hepatitis C therapy over three periods of follow-up: one, five, and 20 years after treatment initiation from a health system perspective. As with the cost-effectiveness analyses, the regimen of interest for genotype 1 was the LDV/SOF strategy (8/12 weeks for treatment-naïve, 12/24 weeks for treatment-experienced), as this represents the cost-effective strategy that is currently available and most likely to receive widespread use in this population. For each of these time points, we used outputs from the care value model to inform expected numbers (per 1,000 treated) of patients experiencing HCV-related complications (cirrhosis, decompensated cirrhosis, hepatocellular carcinoma, liver transplant) and dying of HCV-related causes. Findings for the performance of LDV/SOF vs. PR are presented in Table ES3 on the next page. LDV/SOF produces incremental clinical benefits very soon after treatment initiation; for example, compared with PR alone, LDV/SOF prevents approximately six cases of cirrhosis and two HCV-related deaths per 1,000 patients treated in the first year alone. Benefits are more fully realized at later time points; at five years, LDV/SOF would avert 44 cases of cirrhosis (15 of which would be decompensated), five cases of HCC, and 17 HCV-related deaths per 1,000 treated. Cost offsets would total approximately 7% of incremental treatment costs. At 20 years, there would be a nearly six-fold reduction in the incidence of cirrhosis, HCC incidence would be reduced by more than half, and 140 HCV-related deaths would be averted per 1,000 treated. More than 25% of treatment costs would be offset by these reductions.

We then combined these results with findings from the March 2014 CTAF review for genotypes 2 and 3<sup>180</sup> to assess the one-, five-, and 20-year budgetary impact to two California state agencies (Medi-Cal and the Department of Corrections) of adopting LDV/SOF for genotype 1 and the most effective therapies available for genotypes 2 and 3 (SOF + R for 12 weeks for genotype 2 and 24 weeks for genotype 3). The number of individuals with chronic hepatitis C who rely on these two state entities was recently estimated to total 93,000.<sup>177</sup>


The availability of new HCV treatments would increase costs by approximately \$1.6 billion, \$545 million, and \$901 million for genotypes 1, 2, and 3 respectively (see Figure ES2 on page ES8), resulting in a total increase of \$3 billion, or \$33 PMPM. This represents a 5% increase over the base per-member per-month (PMPM) Medi-Cal costs of \$611. Cost offsets after five years would total \$254 million, reducing net expenditures modestly to \$2.8 billion. More substantial offsets after 20 years (\$1.2 billion) would reduce net expenditures further to \$1.8 billion.

**Table ES3: Clinical Outcomes (per 1,000 Patients Treated) and Costs for LDV/SOF and PR Therapy over One, Five, and 20 Years of Follow-up**

Timeframe/Regimen	Cirrhosis	Liver-Related Complications			HCV	Costs (per patient, \$)		
		Decompensation	HCC	Transplant	Death	Treatment	Other	Total
<u>1 Year</u>								
PR	6.8	3.5	1.8	0.0	5.4	\$34,966	\$1,636	\$36,602
LDV/SOF	0.8	0.6	1.2	0.0	3.4	\$84,341	\$696	\$85,037
Difference (LS-PR)	(5.9)	(3.0)	(0.6)	0.0	(2.0)	\$49,375	(\$940)	\$48,435
<u>5 Years</u>								
PR	34.8	18.7	11.9	0.4	35.3	\$34,966	\$6,681	\$41,647
LDV/SOF	6.1	3.4	6.7	0.3	18.7	\$84,341	\$3,260	\$87,601
Difference (LS-PR)	(28.8)	(15.3)	(5.1)	(0.1)	(16.5)	\$49,375	(\$3,421)	\$45,954
<u>20 Years</u>								
PR	120.9	66.8	45.3	4.9	248.8	\$34,966	\$23,442	\$58,409
LDV/SOF	21.5	11.8	23.0	1.5	109.1	\$84,341	\$10,214	\$94,555
Difference (LS-PR)	(99.4)	(55.0)	(22.3)	(3.3)	(139.7)	\$49,375	(\$13,229)	\$36,146

LS-PR: Difference between LDV/SOF and PR therapy

**Figure ES2: Budgetary Impact of New Hepatitis C Treatments in the Medi-Cal/Department of Corrections Hepatitis C Population in California, with and without Cost Offsets from Reduced Liver-related Complications**



## Drug Pricing to Meet Per-Member Per-Month Benchmarks

PMPM increases of 0.5%-1% in a given year were used in this report as a range of potential budget impact that, when exceeded, are likely to drive specific efforts to manage the costs of a new health care intervention. We examined the incremental drug expenditures at which PMPM increases of 0.5% and 1% would be met for genotype 1, the patient subpopulation of interest in this review. Based on the assumed baseline PMPM in this analysis (\$611) as well as the size of the population to be treated (approximately 33,000 patients in the Medi-Cal/Department of Corrections population in California if 50% of genotype 1 patients present for treatment), a course of treatment with a new agent would need to be priced at \$34,000 - \$42,000 to meet the 0.5% and 1% thresholds respectively.

We also conducted a hypothetical analysis of the number of treatment-naïve Medi-Cal/Department of Corrections patients who could be treated without exceeding these thresholds, based on the current wholesale acquisition costs of LDV/SOF (approximately \$63,000 and \$95,000 for 8 and 12 weeks, respectively). Only two-thirds of these patients (approximately 16,500 of the 26,000 patients presenting for care) could receive treatment at these prices if the one-year PMPM increase were to be held to less than 1% (i.e.,  $\leq \$6.11$ ), leaving nearly 10,000 Medi-Cal/Department of Corrections patients without access to new therapy. When considering a 0.5% threshold for PMPM increase ( $\leq \$3.06$ ), less than half of eligible patients (12,600 of 26,000) could be treated at current prices. In contrast, if the population of treatment-naïve genotype 1 patients is restricted to those with F3 and F4 stage disease ( $n \sim 6,700$ ), LDV/SOF could replace historical PR therapy in all of these patients at current prices and remain under the 1% threshold for PMPM increase. When considering a 0.5% increase in PMPM (\$3.06), LDV/SOF could replace PR in 91% of F3/F4 patients ( $n \sim 6,100$ ) at current prices.

## Summary

Our findings have important implications for patients, physicians, and payers. Specifically, model results suggest that the introduction of LDV/SOF for both treatment-naïve and treatment-experienced individuals would confer substantial clinical benefits in comparison to historical treatment standards and even in relation to other sofosbuvir-based regimens. While the use of this new regimen would increase treatment costs, such use appears to be cost-effective by conventional standards. However, the additional expenditures required to treat all patients with genotype 1 infection (even if only 50% of them are aware of their infection) are substantial; when added to the additional expenditures required for genotypes 2 and 3, this represents a per-member per-month premium increase that is five-fold higher than frequently-discussed manageable thresholds for new interventions. It is clear that patients, physicians, insurers, and health systems will have to grapple with the budget impact of new, highly effective, and expensive treatments for hepatitis C. Whether this will result in prioritization of clinical care, new contracting and financing mechanisms, evolving market dynamics, or policy actions remains to be seen.

# Introduction

---

This assessment for the California Technology Assessment Forum (CTAF) evaluates the evidence on the comparative clinical effectiveness and value of new interferon-free combinations of direct-acting antiviral (DAA) drugs for the treatment of chronic hepatitis C, genotype 1, which is the most common genotype in the United States. Our March 2014 assessment evaluated single DAA drugs used with pegylated interferon and ribavirin. Since that review, the FDA approved two new therapies that each combine two DAAs and do not require the use of either interferon or ribavirin. On October 10, 2014, the FDA approved the combination of ledipasvir/sofosbuvir and on November 5, 2014, the FDA approved the combination of simeprevir/sofosbuvir. At least two more combinations of DAAs are likely to receive FDA approval in the next few months.

Chronic hepatitis C is a common infection that is a major cause of chronic liver disease, liver failure, and hepatocellular carcinoma (HCC), and it is the leading indication for liver transplantation in the Western world.<sup>1</sup> Prior to 2011, the combination of pegylated interferon and ribavirin (PR) was the standard of therapy for the treatment of chronic hepatitis C. Fewer than half of patients with genotype 1 clear the virus from their bloodstream entirely and maintain a sustained virologic response (SVR) 24 weeks after the end of treatment with PR. PR therapy can be difficult, however, as both interferon and ribavirin can produce severe fatigue and body aches, and in some cases, dangerous levels of anemia, neutropenia, and/or thrombocytopenia.<sup>2</sup> The 2011 introduction of first generation DAA protease inhibitors boceprevir (Victrelis®, Merck & Co.) and telaprevir (Incivek®, Vertex Pharmaceuticals, Inc.) resulted in substantially improved SVR rates in many patients when combined with PR. This improvement came with new challenges including significant additional side effects and drug-drug interactions as well as stringent dosing requirements and high pill burdens for patients.<sup>3</sup> In 2013, the FDA approved the second generation of DAAs, simeprevir and sofosbuvir, which in combination with PR increased the SVR, decreased the duration of therapy and decreased adverse events. Since the March 2014 CTAF assessment of hepatitis C therapies, investigators published promising results on several interferon-free therapies that combine two or more DAAs.

As highlighted in the prior CTAF assessment, the new drugs are expensive, with new combination therapies costing approximately \$65,000 to \$190,000 per course of therapy, depending on treatment duration.<sup>4,5</sup> Because chronic infection with HCV is relatively common, this translates into an enormous potential budget impact for federal, state, and private health insurers. ICER commissioned a detailed cost-effectiveness model to provide a more detailed assessment of the benefits and costs of the new drugs for our new assessment.

This assessment will address the following questions: 1) among patients with genotype 1, what is the comparative clinical effectiveness of combinations of two or more DAAs compared to each other as well as to single DAA therapy used in combination with interferon and ribavirin; and 2) what is the comparative value of the new therapies, including analysis of their care value at the patient level and of their potential health system value when budget impact is also taken into

consideration. The purpose of this assessment is to help patients, providers, and payers address these important questions and to support dialogue needed for successful action to improve the quality and value of health care for patients with hepatitis C.



# 1. Background

---

## 1.1 Hepatitis C

The worldwide prevalence of hepatitis C infection is estimated to be between 120 and 170 million.<sup>6</sup> Estimates for the prevalence of hepatitis C in the United States range from 3.0 to 5.2 million people.<sup>7-10</sup> It is the leading cause of liver failure requiring liver transplant.<sup>11</sup>

There are six major genotypes of hepatitis C.<sup>12</sup> The most common genotype in the United States is genotype 1 (70-75%), followed by genotype 2 (13-17%) and genotype 3 (8-12%).<sup>13-18</sup> Genotypes 4 to 6 are uncommon in the United States (1% or less). Knowledge of the viral genotype is important because response to therapy varies by genotype. The new combination therapies considered in this assessment have primarily been studied in genotype 1, and this assessment will focus exclusively on genotype 1.

The majority of patients with chronic hepatitis C infections are asymptomatic and unaware of their infections unless they have been screened. It is estimated that approximately half of patients infected with hepatitis C in the United States are unaware of their infection and that less than 15% have received treatment.<sup>9,19,20</sup> The majority (approximately 76%) of Americans infected with the hepatitis C virus (HCV) were born between the years of 1945 and 1965.<sup>20</sup> Both the Centers for Disease Control and Prevention (CDC) and the US Preventive Services Task Force (USPSTF) now recommend hepatitis C screening for all Americans born during those years.<sup>21,22</sup>

The CDC estimates that among 100 people infected with hepatitis C, only 20 to 30 will develop symptoms acutely (see Table 1 on the next page).<sup>121</sup> The symptoms are primarily fatigue, decreased appetite, nausea, and jaundice. Of 100 people infected with hepatitis C, 75 to 85 will remain chronically infected with hepatitis C.<sup>23-25</sup> Between 60 and 70 of these individuals will develop chronic liver disease, and from 5 to 20 will develop cirrhosis over 20 years.<sup>26,27</sup> If untreated, approximately 1 to 5 individuals out of the original 100 infected will die from cirrhosis or liver cancer. The most common causes of death among patients with chronic hepatitis C are drug overdose, HIV, and liver disease.<sup>28-30</sup> This reflects the epidemiology of hepatitis C infection: many are infected through injection drug use, which puts them at risk for both HIV and drug overdose. Evaluation of death certificates and modeling studies suggest that these statistics may underestimate the morbidity and mortality from HCV infection.<sup>122-124</sup>

**Table 1. Natural History of Hepatitis C Infection over 20 Years**

Condition	Number of individuals
Infection with hepatitis C	100
Develop symptoms	20-30
Remain asymptomatic	70-80
Develop chronic infection	75-85
Develop chronic liver disease	60-70
Develop cirrhosis	5-20
Die from cirrhosis or liver cancer	1-5

As described above, chronic hepatitis C is a slowly progressive disease. Up to 20% of patients develop cirrhosis over 20 to 30 years of infection.<sup>26,27</sup> The risk for cirrhosis may increase with time. One study estimated that the probability of cirrhosis was 16% after 20 years of infection, but increased to 41% after 30 years of infection.<sup>26</sup> Once bridging fibrosis or cirrhosis develops, patients with chronic HCV infection are at risk for the development of hepatocellular carcinoma. Factors associated with an increased risk for progression to cirrhosis include male sex, older age, co-infection with hepatitis B or HIV, obesity, alcohol intake, diabetes, and insulin resistance.<sup>26,27,31-40</sup>

## 1.2 Definitions

- *Cirrhosis*: progressive scarring of liver tissue that may affect the effectiveness of chronic hepatitis C treatment. Cirrhosis is typically biopsy-proven in clinical trials of chronic hepatitis C therapies.
- *Decompensated cirrhosis*: the presence of cirrhosis plus one or more complications including esophageal varices, ascites, hepatic encephalopathy, spontaneous bacterial peritonitis, hepatorenal syndrome, or hepatocellular carcinoma.
- *Genotype*: a classification of hepatitis C based on genetic material in the RNA strands of the virus. There are 6 main genotypes, which are further divided into subtypes in some cases.
- *Interferon-ineligible*: patients in whom interferon therapy is contraindicated due to such conditions as anemia, alcohol abuse, advanced or decompensated cirrhosis, or severe psychiatric disorder.
- *Interferon-intolerant*: patients who discontinue interferon therapy prematurely due to side effects.
- *Sustained virologic response (SVR)*: absence of detectable HCV RNA, measured 12-24 weeks following the completion of treatment.
- *Relapse*: recurrence of detectable viral RNA at some point after achieving an undetectable HCV viral load during treatment.

- *Null response*: no reduction of at least 1 log<sub>10</sub> in HCV RNA during prior treatment.
- *Partial response*: greater than a 1 log<sub>10</sub> reduction in HCV RNA during prior treatment, but never achieving undetectable viral RNA.
- *Treatment-naïve*: not previously treated for chronic hepatitis C infection.
- *Treatment-experienced*: one or more previous attempts at treatment of chronic hepatitis C infection. This group may contain a mix of patients who relapsed, those with a partial response, and those with a null response to prior treatment.

The **METAVIR score** is a standardized measure of fibrosis and inflammation seen on a liver biopsy. The fibrosis score ranges from 0 to 4, and the inflammation activity score is measured from 0 to 3.

Fibrosis score:

F0 = no fibrosis

F1 = portal fibrosis without septa

F2 = portal fibrosis with few septa

F3 = numerous septa without cirrhosis

F4 = cirrhosis

Activity score:

A0 = no activity

A1 = mild activity

A2 = moderate activity

A3 = severe activity

The fibrosis score is particularly useful because patients with higher fibrosis scores are more likely to progress to cirrhosis and HCC and may warrant earlier treatment.

The **Ishak scale** is a second commonly reported histologic grading system for liver fibrosis that ranges from 0 to 6.

Ishak Scale

1 = no fibrosis (normal)

2 = fibrous expansion of some portal areas ± short fibrous septa

3 = fibrous expansion of most portal areas ± short fibrous septa

4 = fibrous expansion of portal areas with marked bridging (portal to portal, portal to central)

5 = marked bridging with occasional nodules (incomplete cirrhosis)

6 = cirrhosis

A rough approximation of how the two scoring systems compare is as follows:

Ishak	METAVIR
0	0
1, 2	1
3	2
4, 5	3
6	4

### 1.3 Treatment of Chronic Hepatitis C Infection

The primary goal of HCV treatment is the prevention of cirrhosis and hepatocellular carcinoma. The combination of pegylated interferon plus ribavirin (commonly referred to as “PR”) has until recently been the backbone of treatment for patients infected with HCV. However, patients infected with genotype 1 tend to have a poor response to PR. As noted earlier, the first generation DAAs – the protease inhibitors boceprevir and telaprevir – were approved for treatment of genotype 1 in 2011. The viral clearance rate with first generation triple therapy (boceprevir or telaprevir + PR) is approximately double the cure rate of the combination of interferon and ribavirin alone. The approvals of simeprevir and sofosbuvir in 2013 were based on data demonstrating improved viral clearance rates in genotype 1 with less toxicity and shorter treatment duration. New DAAs and new combinations that eliminate the need for interferon are poised to enter clinical use with the promise of even higher rates of viral clearance, shorter treatment courses, and fewer side effects.

Because the natural history for the development of cirrhosis and HCC is long, treatment success is usually measured by the maintenance of a sustained virologic response (SVR), defined as undetectable serum HCV RNA for at least 24 weeks (SVR24) after the completion of treatment. In recent trials, the FDA has allowed the SVR 12 weeks after the completion of treatment (SVR12) to be the primary outcome. SVR is a reasonable, but imperfect measure of a clinical “cure”, and it varies somewhat based on when it is measured. For example, the PILLAR trial,<sup>41</sup> a phase 2B trial of simeprevir, reported the number of participants who had undetectable RNA at the end of treatment and at 12, 24, and 72 weeks after treatment. The number of patients with undetectable HCV RNA declined from 336 at the end of treatment to 303 (12 weeks), 300 (24 weeks), and 293 (72 weeks), respectively. Thus SVR12 was a reasonably stable representation of SVR24 (only 3/303 or about 1% relapsed between those two time points). However, relapses did continue over time, with an additional 7/300 (2.3%) relapsing between 24 and 72 weeks of follow-up. One meta-analysis summarized the data on relapse rates among patients treated with PR who achieved SVR12.<sup>42</sup> They found that approximately 6% of patients relapsed between 12 and 24 weeks (SVR12 53% versus SVR24 47%).<sup>42</sup> This may be less of a problem with the newer DAAs, although the data are still limited. A summary of five trials of sofosbuvir-containing regimens found that only 2 of 779 patients achieving SVR12 had detectable viral RNA at 24 weeks (0.3% relapse rate).<sup>43</sup> In a meta-analysis of long-term outcomes with PR, the percent of patients with long-term viral clearance following SVR24

ranged from 98% to 100%.<sup>44</sup> Comparable data are not yet available for the newer DAA-based regimens.

Clinical trial results are typically better than real-world results.<sup>45</sup> Recent data from CVS/Caremark indicate that real world discontinuation rates for sofosbuvir regimens requiring interferon and/or ribavirin may be as high as five times greater than the rates reported in clinical trials.<sup>46</sup> In their data, 10.2% of 738 patients prescribed sofosbuvir + PR discontinued therapy compared to the standard of approximately 2% in the clinical trials. Similarly, 9.0% of 680 patients prescribed sofosbuvir + R discontinued therapy compared to 0-2.0% in the pivotal clinical trials.<sup>46</sup> However, preliminary results from the HCV-TARGET real-world registry, funded by a consortium of pharmaceutical companies manufacturing drugs to treat hepatitis C, reported discontinuation rates that were similar to those observed in the clinical trials.<sup>47</sup> In their data, 2.5% of 366 patients prescribed sofosbuvir + PR discontinued therapy compared to 2% in the clinical trials. Similarly, 3.6% of 645 patients prescribed sofosbuvir + R discontinued therapy compared to 0-2.0% in the clinical trials. It is important to note that in the HCV-TARGET registry, 25.7% of the patients treated with sofosbuvir + PR and 52.0% of the patients treated with sofosbuvir + R had not yet completed treatment, so these reported discontinuation rates are likely to be underestimates of the true values.<sup>47</sup>

## **Treatment of Genotype 1**

### ***Pegylated interferon plus ribavirin***

Pegylated interferon plus ribavirin (PR) was the primary treatment of HCV for more than 10 years. In clinical trials, the SVR24 for patients with genotype 1 treated with PR ranged from 40% to 50%, but it was about 20% lower in real-world studies in part because of the poor tolerability of PR therapy and because of the special nature of patients willing to participate in clinical trials.<sup>48-50</sup> Interferon requires a weekly injection and commonly causes fatigue (50% to 60%), headache (50% to 60%), myalgias (40% to 55%), and fever (40% to 45%).<sup>51</sup> Other common side effects of PR include anemia (hemoglobin < 10 g/dL) in up to 30% of patients, generalized pruritus (25% to 30%), and psychiatric symptoms such as depression (up to 25%), insomnia, and anxiety (15% to 25%).<sup>51</sup> Ribavirin may cause birth defects, so women of child-bearing age must be on birth control during treatment.

For genotype 1, patients are treated for 48 weeks with once-weekly subcutaneous injections of pegylated interferon and twice-daily oral ribavirin taken with food. Routine monitoring is performed with dose reductions recommended for neutropenia, thrombocytopenia, anemia, depression, and worsening renal function.

### ***Boceprevir and Telaprevir***

The first generation protease inhibitors boceprevir and telaprevir were the first two DAAs approved by the FDA. After their approval in 2011, the standard of care for the treatment of genotype 1 became PR in combination with either boceprevir or telaprevir.<sup>52-54</sup> However, the manufacturer of telaprevir discontinued sales in the United States on October 16, 2014 due to declining use after the approval of simeprevir and sofosbuvir. Among treatment-naïve patients in clinical trials, PR plus boceprevir or telaprevir has a SVR24 between 70% and 75%.

Treatment with PR plus either boceprevir or telaprevir is challenging. Patients are required to take either six or 12 pills per day spaced every seven to nine hours with specific dietary restrictions. Both medications increase the risk for severe anemia, which is already common with PR treatment (increased from 30% with PR to 50% with either boceprevir or telaprevir).<sup>51</sup> The combination of PR plus boceprevir or telaprevir is associated with serious adverse event rates between 40% and 50%.<sup>45,51,55</sup> Neither can be used as monotherapy because resistance develops quickly.<sup>56,57</sup> Finally, boceprevir and telaprevir are strong inhibitors of the cytochrome P450 (CYP) 3A4 enzyme, leading to many potential drug interactions with statins, benzodiazepines, colchicine, St. John's wort, anticonvulsants, sulfonyleureas, and some reverse transcriptase inhibitors.

### ***Simeprevir and Sofosbuvir***

Simeprevir is a NS3/4A protease inhibitor that was approved by the FDA for the treatment of HCV genotype 1 in November 2013. It is a second-generation protease inhibitor (boceprevir and telaprevir were first generation protease inhibitors). Simeprevir has several advantages over the earlier protease inhibitors. It may be taken once a day rather than six to 12 pills divided into doses taken every eight hours. It does not appear to increase the risk for anemia, which is a common, often severe, problem with the first generation protease inhibitors. Simeprevir must be used in combination with PR because viral resistance develops rapidly with monotherapy. Simeprevir is taken once daily with PR for 12 weeks followed by an additional 12 weeks of PR for treatment-naïve patients and patients who relapsed or by an additional 36 weeks of PR for prior partial and null responders (see Table 3 on page 11).

Sofosbuvir is the first drug in the class of HCV NS5B nucleotide analog polymerase inhibitors to be approved. Like the other DAAs, sofosbuvir should not be prescribed as monotherapy. It has been studied in combination with PR, with ribavirin alone, with simeprevir, and in combination with other DAAs that have not yet received FDA approval. Like simeprevir, sofosbuvir only needs to be taken once daily. The details of therapy are guided by genotype, prior treatment status, interferon eligibility, and liver histology. The FDA indication for patients with genotype 1 is sofosbuvir 400 mg daily with PR for 12 weeks; patients who are interferon-ineligible may consider sofosbuvir 400 mg plus R alone for 24 weeks (see Table 3 on page 11). For patients who are HIV co-infected, the treatment is the same as for patients who are not HIV co-infected.

### ***Interferon-free therapy combining more than one DAA for Genotype 1***

Boceprevir, telaprevir, simeprevir, and sofosbuvir were the first four DAAs approved by the FDA. More than 30 additional DAAs are in clinical trials. The new drugs attack different targets in the HCV life cycle and include NS3/4A protease inhibitors, nucleoside and nucleotide polymerase inhibitors, non-nucleoside polymerase inhibitors, NS5A inhibitors, and cyclophilin inhibitors. The names and classes of some of the new drugs are summarized in Table 2 on the following page.

At the time of the March 2014 CTAF assessment, preliminary results using several combinations of simeprevir/sofosbuvir with or without ribavirin had been presented at conferences. The study results have now been published, and on November 5, 2014, the FDA approved the combination of simeprevir 150 mg once daily plus sofosbuvir 400 mg once daily without ribavirin for patients with genotype 1 infection (see Table 3 on page 11).<sup>58</sup> Prior to FDA approval, observational studies reported that between 23% and 47% of patients with hepatitis C treated with sofosbuvir-containing combinations were being treated with off-label combinations of simeprevir/sofosbuvir.<sup>46,47</sup>

The FDA approved the combination of ledipasvir/sofosbuvir (LDV/SOF) formulated in a single tablet (Harvoni®) on October 10, 2014. It is taken one pill a day for eight to 24 weeks and is not taken with any additional drugs (see Table 3 on page 11).

Bristol-Meyers Squibb (BMS) has several drug combinations in development. Initial studies show promising results for the combination of daclatasvir/sofosbuvir.<sup>59</sup> BMS has three phase 3 studies of this combination in progress (ALLY 1, 2, and 3), and FDA approval is expected in late 2014.

**Table 2: Therapies for Hepatitis C by Class**

Brand Name	Generic Name	Internal Name	Pharmaceutical Company
<i>Pegylated Interferon Alfa</i>			
PegIntron	peginterferon alfa-2b		Merck
Pegasys	peginterferon alfa-2a		Genentech
<i>Nucleoside analog</i>			
Ribasphere, Virazole	Ribavirin		Genentech
<i>NS3/4A Protease inhibitors</i>			
Incivek	telaprevir		Vertex
Victrelis	boceprevir		Merck
Olysio	simeprevir	TMC435	Janssen and Medivir AB
Sunvepra	asunaprevir	BMS-650032	Bristol-Myers Squibb
n/a	vaniprevir	MK-7009	Merck
n/a	paritaprevir	ABT-450	AbbVie
n/a		MK-5172	Merck
<i>Nucleoside and Nucleotide NS5B Polymerase Inhibitor</i>			
Sovaldi	sofosbuvir	GS-7977	Gilead Sciences
n/a	mericitabine	RG7128	Roche
<i>Non-Nucleotide NS5B Polymerase Inhibitor</i>			
n/a	dasabuvir	ABT-333	AbbVie
n/a		BMS-791325	Bristol-Myers Squibb
n/a		ABT-072	AbbVie
<i>NS5A Inhibitors</i>			
Daklinza	daclatasvir	BMS-790052	Bristol-Myers Squibb
n/a	ledipasvir	GS-5885	Gilead Sciences
n/a	ombitasvir	ABT-267	AbbVie
n/a		GS-5816	Gilead
n/a		MK-8742	Merck
<i>Combination pills</i>			
Harvoni	ledipasvir/sofosbuvir	--	Gilead Sciences

The European Commission approved the use of daclatasvir as part of combination therapy in August 2014, but it has not been approved in the United States. BMS withdrew its application for the combination of asunaprevir/daclatasvir from the FDA in late 2014, but the combination is approved for use in Japan. BMS also has phase 3 studies of the combination of daclatasvir, asunaprevir, and BMS-791325 in progress (UNITY 1, 2, and 3).

In April 2014, AbbVie submitted an interferon-free combination to the FDA of paritaprevir/ritonavir (150/100mg) co-formulated with ombitasvir 25mg, dosed once daily, and dasabuvir 250mg with or without R (weight-based), dosed twice daily. This is known as the “3 DAA” or “3D” regimen. FDA approval is expected in late 2014.

Many physicians have been monitoring patients with chronic HCV infections but not treating them while waiting for new medical therapies (sometimes referred to as “warehousing”). Treatment rates have increased since the approval of simeprevir and sofosbuvir, but many patients have been waiting for additional interferon- and ribavirin-free treatments.



**Table 3. FDA Indications for New DAAs to Treat Genotype 1**

<b>Drug</b>	<b>Genotype</b>	<b>Treatment</b>
Simeprevir	1	<ul style="list-style-type: none"><li>• 150 mg daily with PR x <b>12 weeks</b> plus PR for an additional <b>12 to 36 weeks</b></li></ul>
Sofosbuvir	1	<ul style="list-style-type: none"><li>• 400 mg daily with PR x <b>12 weeks</b></li><li>• Alternate if interferon (IFN)-ineligible: 400 mg daily with R x <b>24 weeks</b></li></ul>
Simeprevir/ sofosbuvir	1	<ul style="list-style-type: none"><li>• 150 mg simeprevir with 400 mg sofosbuvir once daily x <b>12 weeks</b> for treatment-naïve and treatment-experienced without cirrhosis</li><li>• 150 mg simeprevir with 400 mg sofosbuvir once daily x <b>24 weeks</b> for treatment-naïve and treatment-experienced with cirrhosis</li></ul>
Ledipasvir/ sofosbuvir	1	<ul style="list-style-type: none"><li>• 90 mg / 400 mg once daily x <b>12 weeks</b> for treatment-naïve with or without cirrhosis and treatment-experienced without cirrhosis</li><li>• 90 mg / 400 mg once daily x <b>24 weeks</b> for treatment-experienced with cirrhosis</li><li>• Alternate therapy for treatment-naïve patients without cirrhosis and HCV RNA &lt; 6 million IU/ml: 90 mg / 400 mg once daily x <b>8 weeks</b></li></ul>

## 2. Clinical Guidelines

---

Each of the guidelines referenced below may address multiple hepatitis C genotypes. For the purposes of this review, only information specific to genotype 1 will be included. Websites were accessed on October 27, 2014.

**The American Association for the Study of Liver Diseases (AASLD) / Infectious Diseases Society of America (IDSA) / International Antiviral Society – USA (IAS USA) (2014)**

<http://www.hcvguidelines.org>

On January 29, 2014, the AASLD, IDSA, and IAS-USA launched an online guideline for the treatment of chronic hepatitis. The guidelines do not yet include consideration of ledipasvir/sofosbuvir. For genotype 1, current recommendations are 12 weeks of sofosbuvir + PR for interferon-eligible patients, and simeprevir/sofosbuvir ± R for interferon-ineligible patients. Alternative therapies for patients with genotype 1 with genotype 1b or genotype 1a without the Q80K polymorphism are 12 weeks of simeprevir + 24 weeks of PR for interferon-eligible patients and 12 weeks of sofosbuvir + 24 weeks of R for interferon-ineligible patients.

On August 11, 2014, the guidelines were updated to include recommendations on when and in whom to initiate therapy. Recommendations are that highest priority for treatment be given to patients at highest risk for severe complications, including those with advanced liver disease (METAVIR F3 or F4), liver transplant recipients, and patients with severe extrahepatic manifestations of hepatitis C.

**The Department of Veterans Affairs (VA)**

<http://www.hepatitis.va.gov/provider/guidelines/index.asp#S2X>

The VA guidelines have not yet addressed the use of ledipasvir/sofosbuvir. For treatment-naïve genotype 1 patients, the current VA recommendations are for 12 weeks of sofosbuvir + PR, with 12 weeks of simeprevir + 24 weeks of PR as an alternative for patients without the Q80K polymorphism. For treatment-naïve patients who are interferon-ineligible, the recommendation for non-cirrhotic patients is 24 weeks of sofosbuvir + R; an alternative treatment for this group and the recommended treatment for interferon-ineligible cirrhotics is 12 weeks of simeprevir/sofosbuvir + R (not FDA-approved at the time the guidelines were published). Treatment-experienced patients who are interferon-eligible are recommended to receive 12 weeks of sofosbuvir + PR. The recommendation for treatment-experienced, interferon-ineligible patients is 12 weeks of simeprevir/sofosbuvir + R (not FDA-approved at the time the guidelines were published). Alternative recommendations for treatment-experienced patients are 12 weeks of simeprevir + 24-48 weeks of PR for patients without Q80K polymorphism, and 12 weeks of

simeprevir/sofosbuvir + R for patients with cirrhosis (not FDA-approved at the time the guidelines were published).

The VA guidelines currently state that it is reasonable to defer treatment in non-cirrhotic patients without significant extrahepatic disease due to the FDA's expected approval of several highly-effective, low side-effect, interferon-free treatments within the next one to two years.

### **European Association for the Study of the Liver (EASL)**

[http://www.easl.eu/\\_clinical-practice-guideline](http://www.easl.eu/_clinical-practice-guideline)

EASL has also not yet addressed the use of ledipasvir/sofosbuvir in its guidance. The most recent guideline update in April 2014 includes the same regimens for genotype 1 infections as the AASLD and VA guidelines but also includes 12-24 weeks of daclatasvir + PR as an alternative for patients with genotype 1b infections. Interferon-ineligible patients are recommended to receive 24 weeks of sofosbuvir + R, 12 weeks of simeprevir/sofosbuvir ± R, or 12-24 weeks of sofosbuvir/daclatasvir ± R.

EASL recommends that all patients with compensated liver disease due to HCV be considered for treatment and that treatment be prioritized for patients with significant fibrosis (METAVIR F3 or F4) or significant extrahepatic manifestations. EASL states that treatment of patients with METAVIR score F2 is justified. They suggest that treatment for patients with METAVIR scores of F0-F1 may be deferred and that regular assessments be made to assess for disease progression or other reasons to initiate treatment.

### **National Institute for Health and Care Excellence (NICE)**

<http://www.nice.org.uk/guidance/conditions-and-diseases/liver-conditions/hepatitis>

<http://cks.nice.org.uk/hepatitis-c>

NICE has nearly completed its technology appraisals of simeprevir and sofosbuvir and is currently developing technology assessments of daclatasvir, faldaprevir, and two combination therapies: ledipasvir/sofosbuvir and paritaprevir + ritonavir + ombitasvir + dasabuvir (3D).

Broader hepatitis C treatment guidelines have not been updated, however, since 2012 and continue to recommend treatment with telaprevir or boceprevir + PR for patients with genotype 1 infection. The NICE website does not indicate when its guideline will be updated.

## 3. Coverage Policies

---

Coverage policies of a variety of public and private payers for sofosbuvir, simeprevir, and ledipasvir/sofosbuvir were reviewed on November 3, 2014. Each of the policies may address multiple hepatitis C genotypes, but for the purposes of this review, only policies for genotype 1 will be included. Tables summarizing details of coverage policies are provided in Appendix A and include website links for each payer/drug regimen.

### 3.1 Ledipasvir/sofosbuvir (Harvoni)

#### ***Medicare & Medicaid***

No publicly-available coverage policies, prior authorization protocols, or formulary designations for ledipasvir/sofosbuvir were available from CMS or Medi-Cal, California's Medicaid agency.

#### ***Regional Private Payers***

**Health Net** (revised October 28, 2014)

[https://www.healthnet.com/static/general/unprotected/html/national/pa\\_guidelines/harvoni\\_natl.html](https://www.healthnet.com/static/general/unprotected/html/national/pa_guidelines/harvoni_natl.html)

Health Net's interim guidelines for ledipasvir/sofosbuvir provide coverage for patients with genotype 1 chronic hepatitis C infections who have not failed previous treatment that included sofosbuvir and who have fibrosis demonstrated by liver biopsy or noninvasive test corresponding to METAVIR score  $\geq 2$  or biopsy corresponding to Ishak score  $\geq 3$ . Coverage is not available for those with decompensated liver disease.

#### ***National Private Payers/Pharmacy Benefit Managers***

**Aetna** (revised October 31, 2014)

[http://www.aetna.com/products/rxnonmedicare/data/2014/GI/hepatitis\\_c.html](http://www.aetna.com/products/rxnonmedicare/data/2014/GI/hepatitis_c.html)

Aetna covers ledipasvir/sofosbuvir for patients with genotype 1 chronic hepatitis C infections and compensated liver disease who are treatment-naïve or have failed previous treatment with PR  $\pm$  any protease inhibitor. Aetna's policy bulletin states that for patients meeting the criteria for ledipasvir/sofosbuvir, its use will be required over other simeprevir or sofosbuvir regimens unless the patient has a contraindication or intolerance to any of its ingredients. Ledipasvir/sofosbuvir is noted as being less costly and/or more effective in achieving SVR than any other simeprevir or sofosbuvir regimens for previously treated, non-cirrhotic patients. For reauthorization at six weeks

of treatment, hepatitis C RNA levels must have declined more than  $2\log_{10}$  IU/ml at treatment week four.

**Anthem/WellPoint/Express Scripts** (revised October 15, 2014)

[http://www.anthem.com/provider/noapplication/f0/s0/t0/pw\\_e225443.pdf?na=pharminfo&](http://www.anthem.com/provider/noapplication/f0/s0/t0/pw_e225443.pdf?na=pharminfo&)

Anthem covers ledipasvir/sofosbuvir for adults with genotype 1 chronic hepatitis C infections and compensated liver disease who are post-liver transplant, have serious extrahepatic manifestations, or have advanced liver disease demonstrated by imaging or biopsy corresponding to METAVIR, IASL, Batts-Ludwig scores  $\geq 3$  or Ishak score  $\geq 4$ . Ledipasvir/sofosbuvir is not covered for patients with severe renal impairment, patients who have failed prior treatment with sofosbuvir- or ledipasvir-based regimens, and in combination with other NS5A or NS5B inhibitors. Patients must not be actively abusing illicit drugs and/or alcohol, or must be in concurrent substance abuse treatment.

**UnitedHealthcare** (effective October 15, 2014)

[https://www.unitedhealthcareonline.com/ccmcontent/ProviderII/UHC/en-US/Assets/ProviderStaticFiles/ProviderStaticFilesPdf/Tools%20and%20Resources/Policies%20and%20Protocols/Medical%20Policies/Ox\\_MPUB\\_Future\\_Pharmacy/PA\\_Med\\_Nec\\_Harvoni\\_101414.pdf](https://www.unitedhealthcareonline.com/ccmcontent/ProviderII/UHC/en-US/Assets/ProviderStaticFiles/ProviderStaticFilesPdf/Tools%20and%20Resources/Policies%20and%20Protocols/Medical%20Policies/Ox_MPUB_Future_Pharmacy/PA_Med_Nec_Harvoni_101414.pdf)

UnitedHealthcare limits the use of ledipasvir/sofosbuvir to patients with genotype 1 chronic hepatitis C infections who have advanced liver disease (biopsy or imaging corresponding to METAVIR score  $\geq F3$  or its equivalent on the Batts-Ludwig, Knodell, or Ishak scales) or have serious extrahepatic manifestations. Patients meeting these criteria may be either treatment-naïve or have previously failed regimens with PR  $\pm$  any protease inhibitor or sofosbuvir. Patients re-infected with genotype 1 hepatitis C post liver transplant are eligible for treatment with ledipasvir/sofosbuvir. Cirrhotic patients must have stage 4 hepatic fibrosis (METAVIR score of F4 or equivalent). All patients prescribed ledipasvir/sofosbuvir must either have no history of substance abuse or have abstained from illicit drug/alcohol abuse for the past 6 months.

### 3.2 Sofosbuvir

Sofosbuvir in combination with PR has been covered by most payers included in our review, with three payers requiring a fibrosis score of  $\geq F3$  and one requiring a fibrosis score of  $\geq F2$ ; Medi-Cal also allowed for treatment of patients with a lower fibrosis score if they have severe extrahepatic manifestations. Coverage for sofosbuvir + R and simeprevir/sofosbuvir  $\pm$  R was generally limited to patients who were interferon-ineligible. Several payers had limits on sofosbuvir coverage for treatment-experienced patients, often requiring that they not have a previous treatment failure with a regimen inclusive of sofosbuvir. Of the four payers that have released policies on ledipasvir/sofosbuvir since its approval by the FDA in October 2014, Aetna and Health Net have restricted coverage for SOF + PR, SOF + R, or SMV/SOF  $\pm$  R to patients with an intolerance or contraindication to either ledipasvir or sofosbuvir.

### 3.3 Simeprevir

Simeprevir in combination with PR has been covered by most payers included in our review, with two payers requiring a fibrosis score of  $\geq F3$  and one requiring a fibrosis score of  $\geq F2$ ; Medi-Cal also allowed for treatment of patients with a lower fibrosis score if they have severe extrahepatic manifestations. All but one of the payers excluded coverage for genotype 1a patients with the Q80k polymorphism; UnitedHealthcare (UHC) noted that SMV + PR is not the recommended treatment for these patients and an alternative is encouraged. Several payers had limits on simeprevir coverage for treatment-experienced patients, often requiring that they not have a previous treatment failure with a protease inhibitor. As with sofosbuvir, Aetna and Health Net have restricted coverage for SMV + PR to patients with an intolerance or contraindication to either ledipasvir or sofosbuvir.

### 3.4 Coverage Policies across Payers

Aetna and Humana's coverage policies did not specify a level of liver fibrosis needed for coverage of these treatments, and CVS/Caremark required a METAVIR score  $\geq F3$  only for SMV/SOF  $\pm$  R. Medi-Cal, Anthem, and UHC covered treatment with a fibrosis score of  $\geq F3$ ; Medi-Cal also allowed for treatment of patients with a lower fibrosis score if they have severe extrahepatic manifestations. Health Net covered these treatments with a fibrosis score of  $\geq F2$  (except for SMV/SOF  $\pm$  R, for which Health Net has no publicly available policy). As noted above, two of the four payers with ledipasvir/sofosbuvir policies (Aetna and Health Net) have restricted coverage for simeprevir- or sofosbuvir-based regimens to patients with an intolerance or contraindication to either ledipasvir or sofosbuvir.

Coverage for several patient characteristics is summarized below:

- *Treatment-experienced* – for most payers, patients were generally eligible for treatment with a protease or polymerase inhibitor if they had not failed previous treatment with the same type of inhibitor. UHC covered ledipasvir/sofosbuvir for patients who had any previous treatment failure, including sofosbuvir-based regimens. Anthem did not cover simeprevir- or sofosbuvir-based regimens for patients who had failed therapy with any protease or polymerase inhibitor in combination with PR and did not cover LDV/SOF for patients who had failed either LDV or SOF.
- *Decompensated cirrhosis* – most payers covered SOF + R or SMV/SOF  $\pm$  R if decompensation was the reason for a patient's interferon-ineligibility
- *Hepatocellular carcinoma* – most payers covered SOF + R if for patients who were awaiting liver transplants and required that treatment be discontinued if a liver transplant occurs
- *Post-liver transplant* – most payers covered SOF + PR, SOF + R, or SMV/SOF  $\pm$  R for patients who had a liver transplant, although CVS/Caremark only covered SMV/SOF  $\pm$  R for patients who are treatment-naïve post-transplant. Anthem and UHC covered ledipasvir/sofosbuvir

for all post-liver transplant patients, Aetna did not cover this treatment, and Health Net did not specify in this category.

- *Severe renal impairment* – generally not covered or not specified for sofosbuvir-based regimens, and generally not specified for simeprevir-based regimens

Several other coverage requirements are summarized below:

- *Treatment discontinuation if HCV RNA levels not reduced* – five of the seven payers required or recommended this for one or more of the DAA drug regimens
- *Specialist to prescribe or consult on these treatments* – three of the seven payers recommended or required this
- *Abuse of illicit drugs and/or alcohol* – Medi-Cal, Anthem, and UHC had requirements related to this, including concurrent substance abuse treatment, toxicology tests, and/or six months of abstinence prior to treatment

## 4. Previous Systematic Reviews and Technology Assessments

---

We were unable to identify any systematic reviews or formal technology assessments that address the interferon-free combinations of two or more DAAs considered in this assessment.

### 4.1 Formal Health Technology Assessments

No formal health technology assessments were identified for the new multiple DAA combinations. However, the Canadian Agency for Drugs and Technologies in Health (CADTH, <http://www.cadth.ca>) is currently reviewing new DAA agents (among patients with genotype 1 chronic hepatitis C only). Similarly, the National Institute for Health and Care Excellence (NICE, <http://www.nice.org.uk>) in England is reviewing the new DAAs and has draft guidance on sofosbuvir.

### 4.2 Systematic Reviews

No published systematic reviews of the newest DAAs were identified.



## 5. Ongoing Studies

The table below summarizes the ongoing and recently completed Phase III and IV trials with at least one arm including the following combinations of two or more DAAs:

- 1) Simeprevir/sofosbuvir
- 2) Daclatasvir/sofosbuvir
- 3) Ledipasvir/sofosbuvir
- 4) 3D ± ribavirin

We did not include studies focusing exclusively on the treatment of HCV genotypes 2, 3, 4, 5, or 6, or on combinations of drugs that were not considered in this assessment.

Title/Trial Sponsor	Study Design	Comparators	Patient Population	Primary Outcomes	Estimated Completion Date
<b>Simeprevir/Sofosbuvir</b>					
Simeprevir / Sofosbuvir With or Without Ribavirin (RBV) for Interferon-intolerant or Ineligible (IFN-II) Patients With Chronic Hepatitis C (Phase IV)  NCT02214420	Interventional  N = not provided	SMV/SOF vs. SMV/SOF + R	<ul style="list-style-type: none"> <li>• HCV</li> <li>• Interferon intolerant or ineligible</li> </ul>	SVR12	August 2015
The SIM-SOF Trial: A Randomized Trial Comparing Simeprevir-Sofosbuvir Versus Peginterferon/Ribavirin/Sofosbuvir for the Treatment of Chronic Hepatitis C Genotype-1a-infected Patients With Cirrhosis (Phase IV)  NCT02168361	RCT  Open label  N = 82	SMV/SOF vs. SOF + PR	<ul style="list-style-type: none"> <li>• GT 1a</li> <li>• Cirrhosis, compensated</li> </ul>	SVR12	Nov 2014
Efficacy and Safety of a 12-Week Regimen of Simeprevir in Combination With Sofosbuvir in Treatment-Naïve or -Experienced Subjects With Chronic Genotype 1 Hepatitis C Virus Infection and Cirrhosis (Phase III)  NCT02114151	Cohort, single arm  Open-label  N = 103	None	<ul style="list-style-type: none"> <li>• GT 1</li> <li>• Treatment-naïve and experienced</li> <li>• Cirrhosis</li> </ul>	SVR12	April 2015
Efficacy and Safety of a 12- or 8-Week Treatment Regimen of Simeprevir in Combination With Sofosbuvir in Treatment-Naïve and -Experienced Subjects With Chronic Genotype 1 Hepatitis C Virus Infection Without Cirrhosis (Phase III)  NCT02114177	RCT  Open-label  N = 310	SMV/SOF for 8 weeks Vs. SMV/SOF for 12 weeks	<ul style="list-style-type: none"> <li>• GT 1</li> <li>• Non-cirrhotic</li> <li>• Treatment-naïve and experienced</li> </ul>	SVR12	April 2015
Safety and Efficacy of 12 Weeks of Simeprevir and Sofosbuvir, for HIV-infected, HCV Genotype 1 Patients With Advanced Fibrosis (Phase IV)  NCT02206932	Cohort, single arm  Open-label  N = 20	None	<ul style="list-style-type: none"> <li>• GT 1 HCV</li> <li>• Cirrhosis (METAVIR score F3 or F4)</li> <li>• Co-infection with HIV-1</li> </ul>	SVR12	TBD (not yet recruiting)

<b>Daclatasvir/Sofosbuvir</b>					
ALLY-1: Evaluation of Daclatasvir, Sofosbuvir, and Ribavirin in Genotype 1-6 Chronic Hepatitis C Infection Subjects With Cirrhosis Who May Require Future Liver Transplant and Subjects Post-Liver Transplant (Phase III)  NCT02032875	Cohort, multiple arm  Open label  N = 110	None	<ul style="list-style-type: none"> <li>• GT 1, 2, 3, 4, 5, or 6</li> <li>• Chronic HCV before or after liver transplantation</li> </ul>	SVR12	March 2015
ALLY-2: Evaluation of Daclatasvir Plus Sofosbuvir in Treatment-naïve and Treatment-experienced Chronic Hepatitis C (Genotype 1- 6) Subjects Coinfected With HIV (Phase III)  NCT02032888	RCT  Open label  N = 200	None	<ul style="list-style-type: none"> <li>• GT 1, 2, 3, 4, 5, or 6</li> <li>• Treatment-naïve or experienced</li> <li>• HIV-1 co-infection</li> </ul>	SVR12	Jan 2015
<b>Ledipasvir/Sofosbuvir</b>					
Efficacy and Safety of Sofosbuvir/Ledipasvir Fixed-Dose Combination ± Ribavirin in Treatment-Naïve and Treatment-Experienced Japanese Subjects With Chronic Genotype 1 HCV Infection (Phase IIIb)  NCT01975675	RCT, multiple arm  Open Label  N = 341	LDV/SOF vs. LDV/SOF + R	<ul style="list-style-type: none"> <li>• GT1</li> <li>• Treatment-naïve or experienced</li> <li>• Japanese patients</li> </ul>	SVR 12  Major adverse events	Aug 2014 (completed recently)
Efficacy and Safety of Sofosbuvir/Ledipasvir Fixed-Dose Combination for 12 Weeks in Subjects With Chronic Genotype 1 or 4 Hepatitis C Virus (HCV) and Human Immunodeficiency Virus (HIV)-1 Co-infection (Phase III)  NCT02073656	Cohort, single arm  Open label  N = 300	None	<ul style="list-style-type: none"> <li>• GT1 and GT4</li> <li>• HIV-1 co-infection</li> <li>• Treatment-naïve and experienced</li> </ul>	SVR12  Major adverse events	June 2016
Efficacy and Safety of Sofosbuvir/Ledipasvir Fixed-Dose Combination in Treatment-Naïve and Treatment-Experienced Subjects With Chronic Genotype 1 HCV Infection (Phase III)  NCT02021656	Cohort, single arm  Open label  N = 360	None	<ul style="list-style-type: none"> <li>• GT1</li> <li>• Treatment-naïve and experienced</li> <li>• Korean/Taiwanese patients</li> </ul>	SVR12  Major adverse events	June 2017

3D ± R					
<p>MALACHITE-1: Efficacy and Safety of ABT-450/Ritonavir/ABT-267 and ABT-333 Co-administered With and Without Ribavirin Compared to Telaprevir Co-administered With Pegylated Interferon α-2a and Ribavirin in Treatment-Naïve Adults With Chronic Hepatitis C Genotype 1 Virus Infection (Phase III)</p> <p>NCT01854697</p>	<p>RCT</p> <p>Open label</p> <p>N = 314</p>	<p>3D + R vs. 3D vs. Telaprevir + PR</p>	<ul style="list-style-type: none"> <li>• GT1</li> <li>• Treatment-naïve</li> <li>• Non-cirrhotic</li> </ul>	<p>SVR12</p>	<p>July 2015</p>
<p>MALACHITE-2: Efficacy and Safety of ABT-450/Ritonavir/ABT-267 and ABT-333 Co-administered With Ribavirin Compared to Telaprevir Co-administered With Pegylated Interferon α-2a and Ribavirin in Treatment-Experienced Adults With Chronic Hepatitis C Genotype 1 Virus Infection (Phase III)</p> <p>NCT01854528</p>	<p>RCT</p> <p>Open label</p> <p>N = 150</p>	<p>3D + R vs. 3D vs. Telaprevir + PR</p>	<ul style="list-style-type: none"> <li>• GT1</li> <li>• Treatment-experienced</li> </ul>	<p>SVR12</p>	<p>July 2015</p>
<p>TURQUOISE-CPB: Safety and Efficacy of ABT-450/Ritonavir/ABT-267 and ABT-333 With Ribavirin in Adults With Genotype 1 Chronic Hepatitis C Virus Infection and Decompensated Cirrhosis (Phase III)</p> <p>NCT02219477</p>	<p>RCT</p> <p>Open label</p> <p>N = 50</p>	<p>Treatment for 12 vs. 24 weeks</p>	<ul style="list-style-type: none"> <li>• GT1</li> <li>• Cirrhosis, decompensated (Child Pugh score 7-9)</li> </ul>	<p>SVR12</p>	<p>March 2016</p>
<p>TURQUOISE-I: Safety and Efficacy of ABT-450/Ritonavir/ABT-267 (ABT-450/r/ABT-267) and ABT-333 Coadministered With Ribavirin (RBV) in Adults With Genotype 1 Chronic Hepatitis C Virus (HCV) Infection and Human Immunodeficiency Virus, Type 1 (HIV-1) Coinfection</p> <p>NCT01939197</p>	<p>RCT</p> <p>Open Label</p> <p>N = 300</p>	<p>Treatment for 12 vs. 24 weeks</p>	<ul style="list-style-type: none"> <li>• GT1</li> <li>• HIV-1 Co-infection</li> </ul>	<p>SVR12</p>	<p>May 2016</p>
<p>TURQUOISE-III: Safety and Efficacy of Ombitasvir/ABT-450/Ritonavir and Dasabuvir in Adults With Genotype 1b Chronic Hepatitis C Virus (HCV) Infection and Cirrhosis (Phase III)</p> <p>NCT02219503</p>	<p>Cohort, single arm</p> <p>Open label</p> <p>N = 50</p>	<p>None</p>	<ul style="list-style-type: none"> <li>• GT1b</li> <li>• Cirrhosis (Child-Pugh score 5 or 6)</li> </ul>	<p>SVR12</p>	<p>Nov 2015</p>

<p>TURQUOISE-IV: Safety and Efficacy of ABT-450/Ritonavir/ABT-267 (ABT-450/r/ABT-267) and ABT-333 Co-administered With Ribavirin (RBV) in Adults With Genotype 1b Chronic Hepatitis C Virus (HCV) Infection and Cirrhosis (Phase III)</p> <p>NCT02216422</p>	<p>Cohort, single arm</p> <p>Open label</p> <p>N = 36</p>	<p>None</p>	<ul style="list-style-type: none"> <li>• GT1b</li> <li>• Cirrhosis</li> </ul>	<p>SVR12</p>	<p>Sep 2015</p>
<p>TOPAZ-I: Long-Term Outcomes With ABT-450/Ritonavir/ ABT-267 (ABT-450/r/ABT-267) and ABT-333 With or Without Ribavirin (RBV) in Adults With Genotype 1 Chronic Hepatitis C Virus (HCV) Infection (Phase III)</p> <p>NCT02219490</p>	<p>Cohort, single arm</p> <p>Open label</p> <p>N = 1650</p>	<p>3D ± R, for 12 vs. 24 weeks</p>	<ul style="list-style-type: none"> <li>• GT1</li> </ul>	<p>TBD</p>	<p>Dec 2020</p>
<p>TOPAZ-II: Long-term Outcomes With ABT-450/Ritonavir/ABT-267 (ABT-450/r/ABT-267) and ABT-333 With or Without Ribavirin (RBV) in Adults With Genotype 1 Chronic Hepatitis C Virus (HCV) Infection (Phase III)</p> <p>NCT02167945</p>	<p>Cohort, multiple arm</p> <p>Open label</p> <p>N = 600</p>	<p>3D ± R, for 12 vs. 24 weeks</p>	<ul style="list-style-type: none"> <li>• GT1</li> </ul>	<p>TBD</p>	<p>March 2020</p>
<p>RUBY-I: Safety and Efficacy of Ombitasvir/ABT-450/Ritonavir and Dasabuvir With or Without Ribavirin (RBV) in Treatment-Naïve Adults With Genotype 1 Chronic Hepatitis C Virus (HCV) Infection, With Severe Renal Impairment or End-Stage Renal Disease (Phase III)</p> <p>NCT02207088</p>	<p>Cohort, single arm</p> <p>Open label</p> <p>N = 40</p>	<p>3D vs. 3D + R</p>	<ul style="list-style-type: none"> <li>• GT1</li> <li>• Treatment-naïve</li> <li>• Severe or end-stage renal impairment</li> </ul>	<p>SVR12</p>	<p>March 2016</p>

## 6. Evidence Review (Methods & Results)

The goal of this technology assessment is to evaluate the comparative effectiveness and value of new combinations of two or more DAAs in the treatment of chronic HCV genotype 1 infection. We compared the four combination therapies that are expected to be approved by the end of 2014 based on new drug applications (NDAs) to the FDA with the three FDA-approved uses of single DAA therapy with simeprevir or sofosbuvir that were evaluated in our March 2014 assessment (see Table 4 below). There are no randomized or other studies that directly compare the new therapies. The majority of the studies compare different dosing regimens of the same drug combinations to each other but not to older therapies like PR or PR plus one of the first generation protease inhibitors. For our prior review, there were sufficient randomized trials comparing boceprevir, telaprevir, simeprevir, and sofosbuvir to the combination of pegylated interferon and ribavirin (PR) to perform a network meta-analysis. Because there are no randomized trials or other studies directly comparing the interferon-free combinations considered in this review to PR or to each other, it is not possible to perform a network meta-analysis in this review. Instead, we summarize the proportion of patients achieving SVR12 with each new combination and combine them using a meta-analysis of proportions. To allow comparisons with the drug combinations for genotype 1 considered in the prior review, we also calculate new summary estimates for the proportion of patients who achieve SVR using the same methodology. These estimates differ somewhat from those reported in the prior review because of the different method used to produce the summary estimate and because we are now estimating the results in four patient subgroups (naïve, non-cirrhotic; naïve, cirrhotic; experienced, non-cirrhotic; experienced, cirrhotic) rather than two subgroups (naïve, experienced).

**Table 4: Therapies Considered in this Assessment**

Brand Name	Generic Name	Abbreviation	Pharmaceutical Company
<i>FDA-approved comparators from prior review</i>			
Olysio + PR	Simeprevir + PR	SMV + PR	Janssen and Medivir AB
Sovaldi + PR	Sofosbuvir + PR	SOF + PR	Gilead Sciences
Sovaldi + R	Sofosbuvir + R	SOF + R	Gilead Sciences
<i>FDA-approved combinations since prior review</i>			
Olysio + Sovaldi	Simeprevir/sofosbuvir	SMV/SOF	Janssen + Gilead Sciences
Harvoni	Ledipasvir/sofosbuvir	LDV/SOF	Gilead Sciences
<i>Combinations pending FDA approval</i>			
Daklinza + Sovaldi	Daclatasvir/sofosbuvir	DCV/SOF	Bristol-Myers Squibb + Gilead Sciences
3D	Paritaprevir + ritonavir + ombitasvir + dasabuvir	3D	AbbVie

We included all prospective randomized trials and cohorts that reported SVR12 or SVR24 in HCV genotype 1 infected populations. We used fixed effects meta-analysis to summarize the SVR12 and

discontinuation rates within each treatment regimen, but any comparison of these summary SVR12 rates between treatments should be made cautiously because differences in the study samples may explain some of the differences in response rates. To calculate the SVR and discontinuation rates in each individual study, we used the number of patients randomized, even if study subjects were later found to be ineligible, never received treatment, or withdrew consent for the trial. The discontinuation rate includes patients who were lost to follow-up, withdrew consent, or stopped treatment due to adverse events. For our primary analyses, we focused on the four subgroups noted above: treatment-naïve patients with and without cirrhosis and treatment-experienced patients with and without cirrhosis. These represent the primary criteria guiding the choice of therapy for HCV genotype 1.

The Medline database, Embase, Cochrane clinical trials database, Cochrane reviews database, the Database of Abstracts of Reviews of Effects (DARE), the Web of Science, and BIOSIS previews were searched using the key words “simeprevir” OR “sofosbuvir” OR “daclatasvir” OR “ombitasvir” OR “abt-450.” The search was performed for the period from 1945 through September 10, 2014. Full details of the search are in Appendix B. The bibliographies of systematic reviews and key articles were manually searched for additional references. The abstracts of citations were reviewed for relevance and all potentially relevant articles were reviewed in full. Because of the paucity of published data, we included meeting abstracts, FDA documents, and press releases as sources of information. For the results of a study to be included in the meta-analysis of SVR, at least one study group must have received a treatment regimen with dosing similar to the likely final FDA dose for the particular indication. We did not treat the data from study abstracts or FDA documents differently from that abstracted from published studies. If both were available, we preferentially used data from the published study.

The search identified 608 potentially relevant references (see Figure 1 on page 27). After elimination of duplicate and non-relevant references, the search identified 54 publications and abstracts describing clinical trials of new DAAs for the treatment of HCV genotype 1. The primary reasons for study exclusion were (a) early dose finding studies, (b) no data on genotype 1, (c) lack of SVR or other clinical outcomes, or (d) reviews and commentaries. Some of the publications reported the results from more than one study. For genotype 1, there were five studies of simeprevir + PR using the dose recommended by the FDA<sup>41,60-63</sup> and an additional four publications describing five studies of a lower dose alternative in Japan.<sup>64-67</sup> There were three studies of sofosbuvir + PR<sup>68-70</sup> and three studies of sofosbuvir + R.<sup>71-73</sup> For combination therapy with sofosbuvir, there was one published study of simeprevir/sofosbuvir,<sup>58</sup> six publications<sup>74-79</sup> and two abstracts<sup>80,81</sup> of ledipasvir/sofosbuvir, and one published study of daclatasvir/sofosbuvir.<sup>59</sup> Evidence on additional combination therapies included six publications on daclatasvir/asunaprevir<sup>82-87</sup> and six publications on paritaprevir (ABT-450) with ritonavir plus ombitasvir plus dasabuvir, with or without ribavirin (3D ± R).<sup>88-93</sup> In addition, there were 11 publications on other combinations,<sup>94-103</sup> three using the new combinations in HIV co-infected patients,<sup>104-106</sup> and three in patients around the time of liver transplant.<sup>107-109</sup>

We adopted the approach of the ICER Evidence Rating Matrix to evaluate the evidence for each therapy ([ICER Evidence Rating Matrix](#)).<sup>110</sup> The quality of individual studies was assessed by considering the domains listed below, which are adapted from the methods guide of the Agency for Healthcare Research & Quality (AHRQ):


- Similarity of baseline characteristics and prognostic factors between comparison groups
- Well-described methods for randomization and concealment of treatment assignment
- Use of valid, well-described primary outcomes
- Blinding of subjects, providers, and outcome assessors
- Intent-to-treat analysis (all randomized subjects included)
- Limited and non-differential loss to follow-up
- Disclosure of any conflicts of interest

The evidence rating reflects a joint judgment of two critical components:

- a) The **magnitude** of the difference between a therapeutic agent and its comparator in “net health benefit” – the balance between clinical benefits and risks and/or adverse effects AND
- b) The level of **certainty** in the best point estimate of net health benefit.



**Figure 1. Selection of Studies for Inclusion in Review**



### **Key Patient Outcomes**

The four most important outcomes in chronic HCV infection are the development of decompensated liver cirrhosis, hepatocellular carcinoma, liver transplantation, or death from liver-related causes. Because HCV has such a long natural history (often 20-40 years before the development of cirrhosis and HCC), large randomized trials with long-term follow-up are needed to demonstrate improvement in these outcomes. None of the studies identified in the search evaluated these four outcomes. For new drug evaluation, the primary outcome has been the sustained absence of HCV viral RNA for at least 24 weeks after the end of therapy (SVR24). The FDA

has accepted recent studies with a primary outcome of SVR 12 weeks after the end of therapy, and SVR12 was the primary outcome for the majority of the recent phase 3 studies of DAAs.

The vast majority of patients with SVR24 remain HCV free during long-term follow-up. In several studies with five or more years of follow-up, 91% to 100% of patients remained virus free.<sup>111-114</sup> Additionally, patients with SVR24 have marked improvements or normalization of their liver function enzymes as well as improvements in liver histology.<sup>111-116</sup> More importantly, SVR24 has been associated with improvements in quality of life and a reduction in fatigue within months of treatment.<sup>117,118</sup> Recent studies have demonstrated that SVR24 is associated with decreases in decompensated liver disease, hepatocellular carcinoma, liver transplant, and all-cause mortality.<sup>111,119-123</sup> For example, in the HALT-C trial, the investigators prospectively followed 526 patients with advanced fibrosis who received treatment with PR (140 patients with SVR; 386 patients with either non-response, breakthrough, or relapse to therapy) for a median of approximately seven years.<sup>120</sup> The primary outcomes were death, liver transplant, death from liver-related causes, and decompensated liver failure. There was more than an 80% reduction in all clinically important outcomes including death or liver transplantation (HR=0.17, 95% CI: 0.06–0.46), decompensated liver disease or death from liver-related causes (HR=0.15, 95% CI: 0.06–0.38), and incident HCC (HR=0.19, 95% CI: 0.04–0.80).

In a much larger observational study of VA patients using data from their electronic medical records, the benefits of achieving SVR were somewhat lower. Over six years of follow-up, there was a 27% reduction in liver-related complications (HR 0.73, 95% CI 0.66 to 0.82) and a 45% reduction in all-cause mortality (HR 0.55, 95% CI 0.47 to 0.64). The VA study compared patients with an undetectable viral load at one point in time following therapy to those with no documentation of an undetectable viral load.<sup>123</sup> Confounding by indication (sicker patients may be more likely to receive treatment) in the VA study may explain some of the difference between it and studies like HALT-C, which compared responders to non-responders in a population of treated patients.

All of the studies linking SVR to clinical outcomes are observational and thus may be subject to residual confounding. In addition, it is important to note that among patients with SVR, those with cirrhosis prior to treatment were still at risk for HCC during follow-up.<sup>111,112,114,119,120,124</sup> Thus, achieving an SVR24 will not prevent the complications of chronic HCV infection for all patients.

## 6.1 Overview of the Key Studies by Treatment Regimen

This review begins with a summary of the three single DAA treatments reviewed in the March 2014 CTAF assessment, simeprevir and sofosbuvir, because these represent the current standard used to assess the new drug therapies. Then we will review the two new FDA-approved combinations of two DAAs, simeprevir/sofosbuvir and ledipasvir/sofosbuvir. Finally, we will consider the two additional DAA combinations likely to be approved by the end of 2014, daclatasvir/sofosbuvir and

3D. Tables summarizing the results for the individual studies are in Appendix C. In addition, tables summarizing the results of the combination of daclatasvir/asunaprevir, which was withdrawn from the FDA, can also be found in Appendix C. Following this overview, the summary estimates for each of the seven primary treatment regimens will be compared.

### ***Simeprevir + PR***

As described in our prior assessment, there are data available from 10 trials of simeprevir in patients with HCV genotype 1 infections (see Appendix Tables C1 and C2 for details). There are two phase 2 trials (PILLAR, ASPIRE), three phase 3 trials (QUEST-1, QUEST-2, PROMISE), and five Japanese trials (DRAGON, CONCERTO 1-4). The evidence base is remarkable for the large number of randomized trials with an appropriate comparator as a control (8/10 trials). However, the Japanese trials use a lower dose of simeprevir (100 mg rather than 150 mg), so results from those trials do not directly apply to patients in the United States. Without the Japanese trials, 847 patients were randomized to the FDA-approved dose and duration of simeprevir + PR. The quality of the data for simeprevir + PR is higher than that for any of the other therapies, because of the large number of patients randomized and the number of randomized trials with an appropriate comparator. The primary weaknesses of the evidence base for simeprevir + PR is the use of the intermediate outcome, SVR. As noted in the prior review, patients with the Q80k polymorphism have a lower response rate to combination therapy with simeprevir, which decreases the population of patients eligible for simeprevir + PR.

### ***Sofosbuvir + PR***

The clinical trial data for sofosbuvir are more complex (see Appendix Tables C3 and C4). There are data available from only three trials of sofosbuvir that included patients infected with genotype 1 (PROTON, ATOMIC, NEUTRINO), and none of the trials included a control group without sofosbuvir. None of the trials compared sofosbuvir to PR plus another active agent, and a total of 391 patients were randomized to sofosbuvir + PR for 12 weeks. The quality of the trials was lower than that for simeprevir because there were no randomized trials comparing sofosbuvir + PR to a prior standard therapy. As with simeprevir, the outcome was SVR, an intermediate outcome. In addition, there are no data on the effectiveness of sofosbuvir + PR in treatment-experienced patients.

### ***Sofosbuvir + R***

The evidence base for sofosbuvir + R for 24 weeks is even sparser (see Appendix Tables C5 and C6). Only 54 patients with genotype 1 have been studied in clinical trials. There are no treatment-experienced patients treated for 24 weeks in the studies and only six patients with cirrhosis treated for 24 weeks. There are no controlled studies, and the outcomes were all intermediate (SVR).

### ***Simeprevir/sofosbuvir***

The COSMOS trial is the only published study of the combination of simeprevir/sofosbuvir (see Appendix Tables C7 and C8). The study enrolled 81 treatment-experienced patients with genotype 1 fibrosis stages F0 to F2 (Cohort 1) and treated them with four different combinations: simeprevir/sofosbuvir for 12 weeks; simeprevir/sofosbuvir for 24 weeks; simeprevir/sofosbuvir + ribavirin for 12 weeks; or simeprevir/sofosbuvir + ribavirin for 24 weeks. Only 14 patients in Cohort 1 received the FDA-indicated dose of simeprevir/sofosbuvir for 12 weeks.

The study also enrolled 87 patients with genotype 1 fibrosis stages F3 or F4 (Cohort 2) and treated them with the same four combinations. About half of the patients in Cohort 2 (40/87) were treatment-naïve. Only 10 patients in Cohort 2 had cirrhosis and were treated with the FDA-indicated dose: 24 weeks of simeprevir/sofosbuvir.

Eleven patients did not complete the study (6.5%) and the overall SVR12 was 92% (154/168). The number of patients treated according to the FDA indication was small (n=31, see Appendix Table C8), but their overall SVR12 was high (97%). As with the prior studies, the quality of data is limited by the lack of any appropriate control group, the use of an intermediate outcome, and the level of uncertainty due to the small number of patients studied in each of the key patient subgroups.

### ***Ledipasvir/sofosbuvir***

The evidence base is larger for the combination of ledipasvir/sofosbuvir (see Appendix Tables C9 and C10). There are five phase 2 studies and three phase 3 studies. These studies include 841 patients with HCV genotype 1 who received the FDA indicated dose of ledipasvir/sofosbuvir. The SVR12 rates are almost uniformly high (94% to 100%) with the exception of the small ELECTRON 2 trial. The primary methodological concern is the lack of a control group in any of the trials. However, the magnitude of benefit (SVR rate 94% to 100% compared to historical controls of approximately 60%, fewer adverse events) somewhat mitigates this concern.

### ***Daclatasvir/sofosbuvir***

There is only a single published trial of daclatasvir/sofosbuvir (see Appendix Tables C11 and C12). The study assigned 167 patients with HCV genotype 1 to one of seven treatment groups, all of which contained daclatasvir/sofosbuvir. They varied by the length of treatment, inclusion of ribavirin, and whether or not the patients had received prior treatment for HCV. There was no control group. Overall, 98% of patients achieved SVR12. There are three ongoing phase 3 trials of the combination of daclatasvir/sofosbuvir.

## ***Daclatasvir/asunaprevir***

BMS withdrew the NDA for daclatasvir/asunaprevir from the FDA in late 2014, so this combination will not be considered further in our assessment. Details of the six trials of this two-DAA combination are summarized in Appendix Tables C13 and C14.

## **3D**

The last therapy combines three DAAs (paritaprevir (ABT-450), ombitasvir, and dasabuvir) with ritonavir. The combination has been studied most often with ribavirin. There are data from one phase 2 trial (AVIATOR, 14 groups studied) and six phase 3 studies (PEARL II, PEARL III, PEARL IV, SAPPHIRE I, SAPPHIRE II, and TURQUOISE II). The study results are summarized in Appendix Tables C15 and C16. A total of 1,677 patients were treated with either 12 or 24 weeks of 3D + R and the SVR12 rates ranged from 90% to 100%. Two of the trials had placebo groups (SAPPHIRE I, SAPPHIRE II), but none of the trials had active control groups with PR or a single DAA therapy.

## **Important Subgroups**

### ***HIV co-infection***

The data for HIV co-infected patients are sparse but encouraging. Two therapies containing one DAA (simeprevir + PR, sofosbuvir + R) and one dual DAA therapy (ledipasvir/sofosbuvir) have been studied in HIV co-infected patients (see Appendix Tables C17 and C18). For all three of these drug regimens, the SVR12 was approximately the same for HIV co-infected patients as it was for HCV genotype 1 mono-infected patients. There do not appear to be any unexpected interactions with anti-retroviral medications. The numbers in each trial are small, particularly when examining the subgroups defined by prior treatment and cirrhosis. Some of the results are preliminary. For example, the ERADICATE trial treated 50 co-infected patients, but SVR12 results were only available for 10. Large observational studies will be helpful to more firmly establish the efficacy of each of these drug combinations. It is worth noting that the combinations without interferon appear to have lower discontinuation rates than those with interferon.

### ***Pre- or post-transplant***


Similarly, data on the outcomes of treatments for patients on the liver transplant waiting list or post-transplant are rapidly emerging. There are three published trials: one in patients awaiting transplant and two in patients with recurrent infections after liver transplant (see Appendix Tables C19 and C20). The initial results are encouraging, but the discontinuation rates are high, reflecting the illness burden of the near- and post-transplant population. Interactions with immunosuppressive drugs did not interfere with therapy. Data from the pre-transplant population suggest that the earlier SVR is achieved prior to therapy, the more likely for a durable cure after transplant.

## 6.2 SVR Outcomes of Treatment of HCV Genotype 1 in Treatment-naïve, Non-cirrhotic Patients

Figure 2 below presents the results of our fixed-effects meta-analysis of the proportion of treatment-naïve, non-cirrhotic patients achieving SVR in the available prospective cohorts for the seven primary treatment combinations reviewed in this report. The height of each blue bar represents the best estimate of patients achieving SVR, and the vertical black line running through each bar represents the 95 percent confidence interval (95% CI) for the results of each treatment. As noted earlier, there were insufficient placebo-controlled and comparative trials to allow for a network meta-analysis. The first three bars represent treatment with a single DAA plus PR or R alone. The following four bars represent combinations of two or more DAAs without interferon.

The SVR estimates for simeprevir + PR, sofosbuvir + PR, and sofosbuvir + R differ from those in our March 2014 CTAF assessment because of the change in methods used for the meta-analyses and because we did not separate out patients with cirrhosis from those without cirrhosis in the prior assessment. For example, in the prior analysis, our summary estimate from the network meta-analysis for the SVR12 of sofosbuvir + PR in treatment-naïve patients with genotype 1 was 83%. In our updated analysis, our summary estimate for the SVR12 of sofosbuvir + PR in treatment-naïve patients with genotype 1 is 92% in patients without cirrhosis and 81% in those with cirrhosis.

**Figure 2: SVR and 95% Confidence Intervals for the Primary DAA Regimens in Treatment-naïve, Non-cirrhotic Patients**



It is worth noting that some of the estimates have wide confidence intervals. For example, in Figure 2, the combination of simeprevir/sofosbuvir for 12 weeks was only studied in four patients, and the 95% CI for the SVR ranges from 39.8% to 100%.

Additional information, including the number of patients studied for each drug combination as well as the treatment duration and discontinuation rates are summarized in Table 5 below. As noted above, the discontinuation rate includes patients who withdrew consent or were lost to follow-up in addition to those who stopped treatment due to adverse events. Table 5 also includes data for combination therapy used for shorter or longer durations than the FDA indication or for multiple durations when there is not yet an indication for a particular drug combination. We also included the data for 3D without ribavirin, although we did not include it in Figure 2 because it has been less studied and appears to have a lower SVR than the combination of 3D + R. For Figure 2, we chose to represent the most commonly recommended length of treatment for this population of patients (genotype 1, treatment-naïve, non-cirrhotic).

**Table 5: Summary Estimates of SVR and Discontinuation Rates for Treatment-naïve Patients without Cirrhosis**

Therapy	N	Tx Duration	SVR (95% CI)	DR (95% CI)
SMV + PR	473	SMV 12 weeks PR 24-48	.825 (.789-.858)	.062 (.042-.086)
SOF + PR	348	12 weeks	.920 (.888-.948)	.103 (.072-.139)
SOF + R	157	24 weeks	.750 (.675-.819)	.078 (.036-.131)
SMV/SOF	4	12 weeks	1.00 (.398-1.00)	.000 (.000-.602)
SMV/SOF	2	24 weeks	1.00 (.158-1.00)	.000 (.000-.842)
DCV/SOF	35	12 weeks	1.00 (.900-1.00)	.000 (.000-.100)
DCV/SOF	25	24 weeks	1.00 (.863-1.00)	.040 (.001-.204)
LDV/SOF	235	8 weeks	.948 (.913-.976)	.002 (.000-.018)
LDV/SOF	482	12 weeks	.985 (.968-.997)	.013 (.002-.029)
LDV/SOF	184	24 weeks	.984 (.953-.997)	.038 (.015-.077)
3D	493	12 weeks	.949 (.927-.967)	.029 (.015-.046)
3D + R	823	12 weeks	.976 (.963-.986)	.010 (.003-.019)
3D + R	40	24 weeks	.900 (.763-.972)	.075 (.016-.024)

**Tx** Treatment

**SVR** Sustained virologic response

**DR** Discontinuation rate

**PR** Pegylated interferon + ribavirin

**R** Ribavirin

**SMV** Simeprevir

**SOF** Sofosbuvir

**LDV** Ledipasvir

**DCV** Daclatasvir

**3D** AbbVie combination therapy

None of the treatment combinations has been directly compared to any of the others in clinical trials. Thus, the differences in the heights of each bar may in part reflect differences in the populations studied and not true differences in the effectiveness of the respective treatment combinations. Several trends do appear. First, the DAA combinations appear to have higher SVRs than the single DAAs + PR or R with the exception of sofosbuvir + R. Second, the SVRs for these same four combinations do not appear to differ from one another, although there is considerable uncertainty in the estimates for both simeprevir/sofosbuvir and daclatasvir/sofosbuvir. Third, the discontinuation rates during therapy are lower in the new combination therapies with the exception of the 24 week 3D therapy that includes ribavirin.

### 6.3 SVR Outcomes of Treatment of HCV Genotype 1 in Treatment-naïve, Cirrhotic Patients

A similar picture emerges for treatment-naïve patients with cirrhosis, although there is much greater uncertainty for each of the individual treatments (see Figure 3 below). The new, multiple DAA combinations have higher SVRs than the earlier single DAA treatments. It is worth noting in Table 6 on the next page that the SVR12 for 12 weeks of simeprevir/sofosbuvir was only 67%. However, as described in section 6.5 below, the same combination of simeprevir/sofosbuvir for 12 weeks has a 100% SVR when studied in a larger sample of treatment-experienced cirrhotic patients who should be more difficult to treat. It is likely that the SVR of simeprevir/sofosbuvir in a larger sample of treatment-naïve, cirrhotic patients will be higher than the 67% reported in the COSMOS trial. This example highlights the imprecision in the estimates derived from the small number of patients studied for each combination in important patient subgroups.

**Figure 3: SVR and 95% Confidence Intervals for the Primary DAA Regimens in Treatment-naïve, Cirrhotic Patients**




Table 6 gives more detail on each combination therapy as well as additional treatment combinations, primarily varying by length of treatment. The discontinuation rates are generally lower for the new combination therapies, but the confidence intervals are very wide, reflecting the small number of patients with cirrhosis enrolled in these trials.



**Table 6: Summary Estimates of SVR and Discontinuation Rates for Treatment-naïve Patients with Cirrhosis**

Therapy	N	Tx Duration	SVR (95% CI)	DR (95% CI)
<b>SMV + PR</b>	48	SMV 12 weeks PR 24-48	.605 (.459-.742)	.061 (.005-.155)
<b>SOF + PR</b>	43	12 weeks	.814 (.666-.916)	.116 (.039-.251)
<b>SOF + R</b>	11	24 weeks	.545 (.227-.848)	.000 (.000-.013)
<b>SMV/SOF</b>	3	12 weeks	.667 (.094-.992)	.333 (.008-.906)
<b>SMV/SOF</b>	6	24 weeks	1.00 (.541-1.00)	.167 (.004-.641)
<b>DCV/SOF</b>	6	12 weeks	1.00 (.541-1.00)	.000 (.000-.459)
<b>DCV/SOF</b>	4	24 weeks	1.00 (.398-1.00)	.000 (.000-.602)
<b>LDV/SOF</b>	-	8 weeks	-	-
<b>LDV/SOF</b>	57	12 weeks	.892 (.778-.974)	.000 (.000-.043)
<b>LDV/SOF</b>	33	24 weeks	.939 (.798-.993)	.061 (.007-.202)
<b>3D</b>	-	-	-	-
<b>3D + R</b>	86	12 weeks	.942 (.870-.981)	.023 (.003-.081)
<b>3D + R</b>	74	24 weeks	.946 (.867-.985)	.054 (.015-.133)

**Tx** Treatment

- No data

**SVR** Sustained virologic response

**DR** Discontinuation rate

**PR** Pegylated interferon + ribavirin

**LDV** Ledipasvir

**R** Ribavirin

**DCV** Daclatasvir

**SMV** Simeprevir


**3D** AbbVie combination therapy

**SOF** Sofosbuvir

## 6.4 SVR Outcomes of Treatment of HCV Genotype 1 in Treatment-experienced, Non-cirrhotic Patients

There were no studies of sofosbuvir + PR or sofosbuvir + R in treatment-experienced patients with genotype 1 infection (see Figure 4 on the following page). The multiple DAA combinations have similar SVR rates that are consistently higher than simeprevir + PR, although there is greater uncertainty in the estimates for simeprevir/sofosbuvir and daclatasvir/sofosbuvir.

**Figure 4: SVR and 95% Confidence Intervals for the Primary DAA Regimens in Treatment-experienced, Non-cirrhotic Patients**



The discontinuation rates were remarkably low for these treatment-experienced patients (see Table 7 below), perhaps reflecting the tenacity of patients who elect for retreatment.

**Table 7: Summary Estimates of SVR and Discontinuation Rates for Treatment-experienced Patients without Cirrhosis**

Therapy	N	Tx Duration	SVR (95% CI)	DR (95% CI)
SMV + PR	274	SMV 12 weeks PR 24-48	.777 (.725-.825)	.015 (.002-.035)
SOF + PR	-	12 weeks	-	-
SOF + R	-	24 weeks	-	-
SMV/SOF	17	12 weeks	.970 (.781-1.00)	.000 (.000-.083)
SMV/SOF	19	24 weeks	.922 (.724-1.00)	.078 (.000-.276)
DCV/SOF	-	12 weeks	-	-
DCV/SOF	18	24 weeks	1.00 (.815-1.00)	.000 (.000-.185)
LDV/SOF	-	8 weeks	-	-
LDV/SOF	95	12 weeks	.977 (.924-1.00)	.000 (.000-.004)
LDV/SOF	87	24 weeks	.989 (.938-1.00)	.023 (.003-.081)
3D	91	12 weeks	.934 (.862-.975)	.066 (.025-.138)
3D + R	414	12 weeks	.967 (.945-.984)	.015 (.004-.031)
3D + R	20	24 weeks	1.00 (.832-1.00)	.000 (.000-.168)

**Tx** Treatment

**SVR** Sustained virologic response

- No data

**DR** Discontinuation rate

**PR** Pegylated interferon + ribavirin

**R** Ribavirin

**SMV** Simeprevir

**SOF** Sofosbuvir

**LDV** Ledipasvir


**DCV** Daclatasvir

**3D** AbbVie combination therapy

## 6.5 SVR Outcomes of Treatment of HCV Genotype 1 in Treatment-experienced, Cirrhotic Patients

The final patient population considered is patients infected with genotype 1 who are both treatment-experienced and cirrhotic (see Figure 5 below). The study sizes are generally small: 52 patients treated with SMV + PR, 76 patients treated with the three dual DAA regimens combined, and 220 patients treated with 3D + R (see Table 8 on the following page). The point estimate is for nearly 100% SVR rates for the interferon-free therapies compared to 73% for SMV + PR. Furthermore, none of the patients treated with the interferon-free combinations discontinued therapy. If these results are reproduced in larger studies, then we will have confidence that even the most difficult-to-treat patients have an excellent chance to achieve lasting SVR. A study published too recently to be included in the meta-analysis offers additional evidence that this may be the future. Osinusi and colleagues studied 14 patients with HCV genotype 1 who had relapsed after 24 weeks of treatment with sofosbuvir + R in the NIH SPARE trial.<sup>73,79</sup> Half of the patients had advanced liver disease by the Knodell Histology Activity Index. All 14 patients achieved SVR12 (100%) following 12 weeks of therapy with ledipasvir/sofosbuvir.<sup>79</sup>

**Figure 5: SVR and 95% Confidence Intervals for the Primary DAA Regimens in Treatment-experienced, Cirrhotic Patients**



**Table 8: Summary Estimates of SVR and Discontinuation Rates for Treatment-experienced Patients with Cirrhosis**

Therapy	N studied	Tx Duration	SVR (95% CI)	DR (95% CI)
SMV + PR	52	SMV 12 weeks PR 24-48	.734 (.601-.850)	.166 (.071-.286)
SOF + PR	-	12 weeks	-	-
SOF + R	-	24 weeks	-	-
SMV/SOF	4	12 weeks	1.00 (.398-1.00)	.000 (.000-.602)
SMV/SOF	4	24 weeks	1.00 (.398-1.00)	.000 (.000-.602)
DCV/SOF	-	12 weeks	-	-
DCV/SOF	3	24 weeks	1.00 (.292-1.00)	.000 (.000-.708)
LDV/SOF	-	8 weeks	-	-
LDV/SOF	43	12 weeks	.846 (.712-.948)	.000 (.000-.044)
LDV/SOF	22	24 weeks	1.00 (.846-1.00)	.000 (.000-.154)
3D	-	12 weeks	-	-
3D + R	122	12 weeks	.902 (.834-.948)	.016 (.002-.058)
3D + R	98	24 weeks	1.00 (.832-1.00)	.000 (.000-.168)

**Tx** Treatment

- No data

**SVR** Sustained virologic response

**DR** Discontinuation rate

**PR** Pegylated interferon + ribavirin

**LDV** Ledipasvir

**R** Ribavirin

**DCV** Daclatasvir

**SMV** Simeprevir

**3D** AbbVie combination therapy

**SOF** Sofosbuvir

## 6.6 Harms of Treatment


The adverse events reported in the clinical trials are summarized in Table 9 on the next page. The combinations that include ribavirin have an increased incidence of anemia, particularly when taken for 24 weeks or when combined with interferon. The combinations that include simeprevir are associated with a greater incidence of rashes. However, it is evident in Table 9 that the elimination of interferon from the treatment regimen markedly decreases the risk for several adverse events including fatigue, headache, flu-like illness, anemia, pruritus, nausea, and rashes. There were also significantly fewer grade 3 or 4 adverse events, when those were reported.

**Table 9: Adverse Events in the Clinical Trials of New Drug Combinations for Hepatitis C**

	SMV12 + PR24/48	SOF12 + PR12	SOF24 + R24	SMV/SOF12	SMV/SOF24	LDV/SOF8	LDV/SOF12	LDV/SOF24	DCV/SOF12	DCV/SOF24	DCV/ASV	3D + R12	3D + R24
	N = 781	N = 327	N = 566	N = 28	N = 31	N = 215	N = 539	N = 326	N = 41	N = 80	N = 645	N = 1379	N = 172
<b>Any Adverse Event</b>	95%	95%	88%	71%	94%	76%	69%	81%	93%	84%	85%	85%	91%
<b>Significant Adverse Events</b>	2%	1%	4%	0%	3%	2%	2%	6%	2%	8%	6%	3%	5%
<b>Grade 3 or 4 AE</b>	23%	15%	7%	7%	13%	NR	NR	NR	2%	2%	NR	NR	NR
<b>Therapy stopped due to AE</b>	3%	2%	1%	0%	7%	0%	1%	0%	0%	1%	2%	1%	2%
<b><u>Common AEs</u></b>													
<b>Fatigue</b>	36%	59%	40%	25%	25%	21%	22%	24%	37%	36%	22%	33%	46%
<b>Headache</b>	33%	36%	23%	21%	21%	14%	21%	24%	22%	25%	24%	30%	31%
<b>Flu-like illness</b>	26%	16%	3%	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
<b>Insomnia</b>	17%	25%	16%	14%	14%	5%	8%	9%	10%	5%	2%	14%	18%
<b>Anemia (hemoglobin &lt; 10 g/dL)</b>	12%	23%	9%	0%	3%	0%	0%	0%	0%	0%	NR	3%	10%
<b>Pruritus</b>	22%	17%	9%	11%	11%	1%	4%	3%	2%	4%	7%	15%	19%
<b>Nausea</b>	22%	34%	20%	21%	21%	7%	11%	11%	20%	28%	12%	20%	20%
<b>Rash</b>	28%	18%	8%	11%	16%	1%	4%	6%	5%	4%	NR	11%	14%
<b>Photosensitivity</b>	3%	NR	NR	7%	7%	NR	NR	NR	NR	NR	NR	NR	NR
<b>Diarrhea</b>	NR	NR	NR	NR	16%	7%	7%	10%	5%	10%	NR	12%	17%

## 6.7 ICER Staff Evidence Rating

The ICER clinical effectiveness rating arises from a joint judgment of the level of certainty provided by the body of evidence and the magnitude of the net health benefit -- the overall balance between benefits and harms. This method for rating the clinical effectiveness is modeled on the “Evidence-Based Medicine (EBM) matrix” developed by a multi-stakeholder group convened by America’s Health Insurance Plans. This matrix is depicted below:



**A = “Superior”** - High certainty of a substantial (moderate-large) net health benefit

**B = “Incremental”** - High certainty of a small net health benefit

**C = “Comparable”** - High certainty of a comparable net health benefit

**D=“Negative”** - High certainty of an inferior net health benefit

**B+=“Incremental or Better”** – Moderate certainty of a small net health benefit, with high certainty of at least incremental net health benefit

**C+=“Comparable or Better”** - Moderate certainty of a comparable net health benefit, with high certainty of at least comparable net health benefit

**P/I = “Promising but Inconclusive”** - Moderate certainty of a small or substantial net health benefit, small (but nonzero) likelihood of a negative net health benefit

**I = “Insufficient”** – Either moderate certainty that the best point estimate of comparative net health benefit is comparable or inferior; or any situation in which the level of certainty in the evidence is low

When the four multiple DAA therapies are compared to the three older SMV or SOF + PR or R regimens, there is moderate certainty of substantial net benefit with high certainty of at least a small benefit. **Rating: B+.**

**Rationale:** The net benefit reflects the clinically important increase in SVR12 with the multiple DAA-containing therapies and fewer side effects, shorter duration of therapy, and less burdensome treatment (fewer pills, no injections, no interferon); the limitations are the small study sizes with no relevant comparators and SVR12 being only a moderately validated intermediate outcome.

When the four multiple DAA therapies are compared to each other, there is low certainty about the superiority of any one therapy. **Rating: I**

**Rationale:** There are no studies directly comparing two or more of the therapies. In addition, the number of patients in the existing studies is often small, so the estimates of benefits and harms have wide confidence intervals. In addition, the four therapies had roughly comparable net benefits in each of the four subgroups studied.

## 6.8 Summary

Treatment for chronic hepatitis C infection has come a long way from 2010, when interferon combined with ribavirin was the sole therapy. This early drug combination, while providing the first effective treatment for chronic hepatitis C, caused fever and flu-like symptoms in almost half of patients, required a year of injections, and led to viral clearance in fewer than half of patients with the most common form of infection, genotype 1. The combination of PR with first-generation DAAs telaprevir or boceprevir increased the rate of viral clearance above 50% but caused severe anemia in up to half of patients, along with significant nausea, and many drug interactions in addition to the side effects of interferon and ribavirin. The clinical trial data on simeprevir and sofosbuvir demonstrated further increases in the rate of viral clearance, shortened length of therapy, and decreased side effects but still required interferon for patients with genotype 1. Treatments that combine two or more DAAs are simpler, shorter, and cause very few side effects while producing extremely high rates of viral clearance in clinical trials.

The evidence on the comparative clinical effectiveness of the newest treatment regimens appears consistent in all four major treatment subgroups. Among treatment-naïve patients without cirrhosis, the SVR12 for simeprevir or sofosbuvir combined with interferon and/or ribavirin is between 75% and 92%. The comparable SVR12 for DAA combination therapy (simeprevir/sofosbuvir, ledipasvir/sofosbuvir, daclatasvir/sofosbuvir, 3D) ranges from 95% to 100%. Among treatment-naïve patients with cirrhosis, the SVR12 for primary DAA therapy ranges from 55% to 81% compared to 67% to 95% for DAA combination therapy. For treatment-experienced patients, the SVR12 for older therapy is about 75% for both cirrhotic and non-cirrhotic patients compared with 95% to 100% for DAA combination therapy.

Due to the lack of head-to-head trials and the lack of trials with common comparators, it is difficult to know if one of the DAA combination therapies is clearly superior to another. At this time only two combinations have FDA approval (simeprevir/sofosbuvir, ledipasvir/sofosbuvir). Two of the

combinations (simeprevir/sofosbuvir, daclatasvir/sofosbuvir) have been studied in very small samples, and the confidence intervals around the estimates for their SVRs are wide. For the patient population with cirrhosis, the confidence intervals are wide for all four of the new DAA combinations. Furthermore, selection bias may explain some of the differences between the SVRs.

Adverse effects may also impact the choice of therapy, although there were very few discontinuations from therapy in any of the studies due to adverse events, and the rate of serious adverse events was similarly low. When patient characteristics require longer therapy with ribavirin-based therapy (sofosbuvir + R for 24 weeks, 3D + R for 24 weeks), the adverse event rates are higher, and the rate of significant anemia is higher. Simeprevir also causes photosensitivity and more rashes.

High-quality observational data from real world settings will be essential for evaluating the comparative effectiveness of the combination DAA therapies. It is unlikely that there will be head-to-head randomized trials of the current therapies, and many more new drug combinations are being tested in clinical trials today. The SVR12 rates of the studied combination therapies will undoubtedly be lower in observational studies than those reported in the clinical trials, as has been seen with earlier DAAs. Patients who qualify to be in clinical trials are generally more motivated, adherent, and have fewer comorbidities than the larger population of patients with chronic HCV infection who need to be treated. Studies including larger numbers of patients treated with each of the drug combinations will help to identify rare adverse events that have not yet been anticipated and should help to clarify specific patient populations that benefit more from one combination therapy than another. It is incumbent upon researchers working closely with the clinical community to continue to collect high quality observational data to help answer the many remaining questions.



## 7. Model of Clinical and Economic Outcomes of Treatment Strategies for Hepatitis C

As noted in this review, new medications for hepatitis C have the potential to change clinical expectations for achieving sustained virologic response in many more patients than previously thought possible. However, these medications also have the potential to substantially increase health-system costs. We developed simulation models of these new regimens for the express purpose of assessing their potential value along two important constructs:

- **Care Value:**
  1. Comparative clinical effectiveness of each regimen vs. alternatives (considering both clinical benefits and harm)
  2. Any additional “non-clinical” benefits (e.g., reduced caregiver burden)
  3. Contextual considerations (no other acceptable treatment, vulnerable populations)
  4. Cost-effectiveness (incremental cost to achieve important patient outcomes vs. alternatives)
- **Health System Value:**
  1. Care value of the regimen of interest (as above): and
  2. Potential effects of short-term budgetary impact from each regimen on other patients in the health care system

Discussion of the methods and results of our modeling efforts can be found starting in Section 7.2. For comparison purposes, we also identified published studies of the cost-effectiveness of both existing and proposed treatment options for hepatitis C, which are summarized in Section 7.1 below. We limited our summary to those studies focusing on the agents of interest in this review and also included studies that focused on hypothetical all-oral regimens.

### **7.1 Prior Published Evidence on Costs and Cost-effectiveness**

We identified a total of seven studies evaluating the cost-effectiveness of sofosbuvir-based regimens, including two that also assessed the use of simeprevir. We found no published studies that have as of yet assessed the cost-effectiveness of ledipasvir/sofosbuvir (LDV/SOF). However, we did identify three studies that focused on the potential cost-effectiveness of hypothetical all-oral regimens for hepatitis C. Populations analyzed, regimens evaluated, and primary findings are summarized in the sections that follow; not surprisingly, most of these analyses found that results were highly sensitive to the assumed costs of treatment and SVR rates.

## **Sofosbuvir vs. Simeprevir**

Hagan and colleagues developed a Markov state-transition model to assess the lifetime cost-effectiveness of SMV/SOF (12 weeks) vs. SOF + R (24 weeks) in a 50-year old cohort of genotype 1 patients ineligible or intolerant to interferon.<sup>167</sup> A SMV/SOF strategy was found to produce three months of additional quality-adjusted life expectancy relative to SOF + R and was cost-saving, reducing overall costs by nearly \$80,000 per patient on a lifetime basis.

Another recent Markov model evaluated the lifetime economic impact of PR therapy alone as well as in combination with sofosbuvir, simeprevir, telaprevir, or boceprevir in a cohort of genotype 1 patients aged 52 years.<sup>168</sup> Outcomes and costs were evaluated separately for treatment-naïve, treatment-experienced, and HIV-coinfected patients. SOF + PR was less costly and more effective than any other triple therapy in all three cohorts of interest and yielded cost-effectiveness estimates of <\$10,000 per QALY gained vs. no treatment as well as <\$30,000 per QALY gained vs. PR alone.

## **Sofosbuvir vs. Older Regimens**

Two studies compared the cost-effectiveness of sofosbuvir to older regimens among patients with genotype 1 infection.<sup>169, 170</sup> One was a lifetime simulation model conducted from the perspective of the Italian National Health Service, and it involved separate comparisons of triple therapy with sofosbuvir vs. boceprevir and telaprevir in genotype 1 patients who were naïve to treatment and age 50 years.<sup>169</sup> Strategies with an incremental cost per life-year gained less than €25,000 (\$35,000) were considered to be cost-effective. Sofosbuvir triple therapy was estimated to increase life expectancy by approximately eight months relative to boceprevir and three months vs. telaprevir. Sofosbuvir was considered to be cost-effective in comparison to either of the competing strategies but not universally so across all subgroups. For example, sofosbuvir was considered to be cost-effective among cirrhotic patients and those with the IL28b CC allele but not in patients with lower levels of fibrosis or in patients with the genotype 1b subtype.

The other study assessed the lifetime cost-effectiveness of sofosbuvir to older regimens among incarcerated individuals in the US serving either short (<1.5 years) or long (≥1.5 years) prison terms.<sup>170</sup> Among those serving short sentences (with no treatment as the only alternative), SOF + PR produced three- to four-fold reductions in the incidence of severe liver-related complications, generated over two additional years of quality-adjusted life expectancy, and resulted in a cost-effectiveness estimate of ~\$26,000 per QALY gained. Findings were similar for those incarcerated long-term, and sofosbuvir triple therapy had more favorable cost-effectiveness ratios than boceprevir triple therapy or PR alone. This study also addressed the affordability question, estimating that sofosbuvir would increase treatment costs for 500,000 prisoners by \$27-\$30 billion, and cost offsets from reductions in liver-related complications (\$2-\$5 billion) would likely be realized outside the prison system.

An additional two analyses assessed the cost-effectiveness of sofosbuvir-based regimens across genotypes 1, 2, and 3 vs. the previous standard of care from the perspectives of the French and Spanish national health systems respectively.<sup>171, 172</sup> For genotype 1, the comparison was to triple therapy with telaprevir or boceprevir as well as to PR alone. Both studies considered a benchmark of €40,000 (\$50,000) per QALY gained to represent a cost-effective use of resources. The French evaluation found that, across all genotypes, sofosbuvir-based regimens increased quality-adjusted life expectancy by an average of two years and resulted in an incremental cost-effectiveness ratio of approximately €16,000 (\$20,000) per QALY gained.<sup>171</sup> Cost-effectiveness improved with increasing fibrosis stage, but treatment met the cost-effectiveness threshold at all stages. In contrast, the Spanish evaluation found that sofosbuvir-based regimens were below the cost-effectiveness benchmark only for genotypes 1 and 3; genotype 2 regimens exceeded this threshold, as did SOF + R for 24 weeks when used in any of the three genotypes.<sup>172</sup>

### **Cost-Effectiveness of All-Oral Hepatitis C Regimens**

As mentioned previously, we found no published assessments of the economic impact of LDV/SOF. However, three simulation models have assessed the potential cost-effectiveness of hypothetical combinations of all-oral drugs.<sup>173, 174, 175</sup> In an NIH-funded analysis, Hagan and colleagues assessed the cost-effectiveness of a hypothetical 2-drug regimen over a lifetime vs. standard care (i.e., triple therapy with older DAAs or PR) across all genotypes in a 50 year-old treatment-naïve cohort using a societal perspective.<sup>173</sup> Based on SVR and drug cost estimates of 90% and \$70,000 respectively, all-oral therapy resulted in an overall gain of five months of quality-adjusted life expectancy while generating approximately \$20,000 more in costs. The resulting cost-effectiveness ratio was \$45,000 per quality-adjusted life year (QALY) gained. However, all-oral therapy was no longer considered cost-effective in this model (at a \$50,000 per QALY threshold) at prices exceeding \$75,000. An industry-funded analysis involving the same comparators produced a lower cost-effectiveness ratio (\$15,709 per QALY gained), which appears to be closely tied to the assumptions that (a) all-oral drug costs would be equivalent to those of existing triple therapy with telaprevir; and (b) SVR rates with all-oral therapy would be 99%, with no discontinuation.<sup>174</sup>

The third evaluation involved a comparison of hypothetical all-oral treatment to both older triple therapy with telaprevir and boceprevir as well as to SOF + PR in treatment-naïve genotype 1 patients.<sup>175</sup> SVR rates were assumed to be 89% for SOF + PR and 85-95% for all-oral treatment, depending on fibrosis stage. Costs of SOF+PR were estimated to be approximately €5,100 (\$6,375) per week based on the French early access price; costs of all-oral therapy were assumed to be double this amount. Treatment with SOF + PR was cost effective relative to older triple therapy (~\$47,000 per QALY gained), but only for patients treated at F2 and above. All-oral regimens were not cost-effective at assumed prices (ICERs of \$170,000-\$400,000 per QALY gained, depending on fibrosis stage) but would be considered cost-effective at weekly prices similar to those of SOF + PR.

## 7.2 Model of Care Value: Overview and Methods

### Overview

We constructed a decision-analytic multistate Markov model<sup>125</sup> to determine the cost-effectiveness of six treatment regimens for HCV genotype 1 currently marketed in the US, as shown in Table 10 below. Note that there are two rows for LDV/SOF; we alternatively assumed that 1) a percentage of treatment-naïve non-cirrhotic patients would be candidates for eight weeks of therapy (LDV/SOF 8/12) and 2) all treatment-naïve patients would receive 12 weeks of therapy (LDV/SOF 12). The percentage of patients eligible for eight weeks of therapy in the LDV/SOF 8/12 strategy was assumed to be 67% based on the proportion of clinical trial subjects with viral loads <6 million IU/ml; this percentage was varied from 30% to 90% in sensitivity analyses.

**Table 10. Modeled Therapies: Interferon-based and Interferon-free Treatments**

		Duration of therapy (weeks)	
		Treatment-naïve	Treatment-experienced
Interferon-based therapies			
1	Peg-Interferon + ribavirin (PR)	48	48
2	Sofosbuvir + PR (SOF + PR)	12	12
Interferon-free therapies			
3	Sofosbuvir + R (SOF + R)	24	--
4	Simeprevir/sofosbuvir (SMV/SOF)	12	12
5	Ledipasvir/sofosbuvir (LDV/SOF 8/12)	8/12*	--
6	Ledipasvir/sofosbuvir (LDV/SOF 12)	12	12/24†

\* – F0-F3 – treatment duration for 67% of patients is 8 weeks, duration for 33% is 12 weeks; F4 – treatment duration is 12 weeks

† – F0-F3 – treatment duration is 12 weeks, F4 – treatment duration is 24 weeks.

The FDA-recommended dosing used in this model is daily 400mg of sofosbuvir, daily 1200mg of ribavirin, and weekly 180mcg subcutaneous injection of peg-interferon alfa-2a.<sup>136</sup>

We limited our inclusion of simeprevir to its recently-approved use with sofosbuvir, as utilization data indicate that simeprevir + PR, while FDA-approved for genotype 1, is rarely used.<sup>46</sup> We also did not consider the first-generation DAAs (boceprevir and telaprevir), as their use has either formally or essentially been discontinued in the US. Finally, we excluded daclatasvir and the 3D regimen from these analyses, as these agents are not yet FDA-approved and no estimates are currently available on their projected cost. As another referent category, we also calculated outcomes and costs among patients receiving no antiviral therapy (i.e., “no treatment”).

The model is designed to calculate the net costs, health benefits, and incremental cost-effectiveness ratios (ICERs) of these therapies. It is also designed to determine how these ICERs change if treatment is delayed to a more advanced stage of disease as compared with treating

people at all disease stages. We thus aimed to address two key policy or program questions with regard to HCV therapy:

- Comparing *regimens*. Which regimens are most cost-effective? Specifically, what is the incremental cost-effectiveness of more expensive and effective regimens?
- Comparing population treatment *strategies*. What is the cost-effectiveness of treating all individuals, as compared with waiting to treat at more advanced disease stages?

To address these issues, the model portrays HCV natural history: the lifetime progression of a prevalent cohort based on the fibrosis stage (i.e., METAVIR F0-F4) of individuals who are aware of their HCV status. The model also portrays *regression* of liver damage after successful treatment.<sup>125</sup> Costs include those of treatment, other medical care outside of and after treatment, and costs of treating serious HCV-related complications such as decompensated cirrhosis and hepatocellular carcinoma. Effectiveness is measured primarily in terms of quality-adjusted life years; however, the incidence of serious HCV-related complications also is assessed.

All results are portrayed for the individual's lifetime and discounted to the present. Separate analyses were conducted for treatment-naïve and treatment-experienced patients. While regimens also differ in terms of whether patients have cirrhosis, this was incorporated into our calculations based on disease progression and regression; for example, LDV/SOF patients treated at METAVIR stage F4 (cirrhosis) received a longer duration of treatment and had different rates of viral clearance. For each of these two groups, we also present results for the two treatment strategies, "treat all" and "wait until more advanced disease." Finally, we present results for a mixed cohort of treatment-naïve and treatment-experienced patients.

Health benefits, including rates of sustained virologic response (SVR), were adjusted for rates of discontinuation as reported in clinical trials (see Appendix Tables D3 and D4). For each treatment regimen both the costs of managing treatment-associated adverse events and the accompanying "disutility" (reduction in well-being) were estimated and incorporated. Consistent with standard methods for health-economic evaluations, future benefits and costs were discounted by 3%,<sup>126</sup> and all cost inputs were adjusted to 2014 dollars by the medical component of the US Consumer Price Index (CPI) (<http://www.bls.gov/cpi/cpid1408.pdf>).

The model was constructed in TreeAge® Pro 2014, with additional analyses in Microsoft Excel®.

## Perspective

In keeping with CTAF standards, analyses were conducted from the health care payer perspective such as a state Medicaid agency or a managed care organization. Cost estimates were thus limited to direct medical costs only (i.e., costs of drug treatment, HCV management, and treatment of HCV

complications). Direct costs to patients (e.g., transportation) and time costs (i.e., productivity losses associated with getting treated) were not included. Potential increases in future lifetime productivity resulting from successful treatment were also not quantified.

There are no universally accepted criteria for what constitutes an acceptable cost-effectiveness threshold for medical care interventions in the United States. Historically, an ICER under \$50,000 per QALY has been used as one threshold, whereas more recent investigators and policy makers have suggested that ICERs under \$150,000 per QALY may be a reasonable threshold for an intervention to be deemed “cost-effective.”<sup>129, 130</sup> Recently, the World Health Organization has promulgated suggested cost-effectiveness thresholds linked to national Gross Domestic Product (GDP).<sup>131, 132</sup> According to the WHO, an intervention with a cost per QALY less than 1 x GDP can be considered “highly” cost-effective, whereas a cost per QALY higher than 3 x GDP is considered not cost-effective. Current GDP for the US is approximately \$50,000 per capita, and therefore thresholds of \$50,000 per QALY and \$150,000 per QALY are considered in this report as important benchmarks.

## **Patient Population**

Patients for this model were assumed to weigh 75kg and be 60-years of age, selected on the basis of a 2010 analysis of National Health and Nutrition Examination Survey (NHANES) data, indicating that the highest HCV prevalence, (3.5%), is found among individuals born between 1945 and 1965 (i.e., ages 45-65).<sup>133</sup> Since 2010, the age distribution has likely shifted, suggesting that an average age of 60 for a prevalent population is appropriate for estimating the impact of HCV therapy. The distribution of patients across fibrosis stages F0-F4 in our modeled cohort is 0.17, 0.35, 0.22, 0.14, and 0.12, respectively (see Appendix Table D1 for details).<sup>137</sup> This distribution is based on empirical assessments of individuals with known HCV infection.<sup>134</sup> The model does not distinguish patients by viral concentrations, sex, or race, although these factors may affect treatment outcomes and disease progression.<sup>135</sup>

## **Natural History of Progression and Treatment Effects**

The natural history of HCV progression and the related disease-state transition probabilities are based on a review of published literature (see Table 11 on the following page). The SVR rates for all treatments except PR were derived from the meta-analyses described in Section 6 of this report. More details on the design of the natural history model including graphical depictions are available in Appendix E.

**Table 11: Key model Inputs: Chronic Hepatitis C Annual Transition Probabilities, Background Mortality, Weekly Cost of Drugs, Cost of Treatment-related Medical Care, and Annual Cost of CHC-related Health Care.** Note: All costs are in 2014 dollars.

Natural History					
Source State	Target State	Base case	Lower limit	Upper limit	Referenc
F0	No progression (proportion)*	0.24	0.10	0.40	138
	F1	0.077	0.067	0.088	137
	Spontaneous Resolution	0.002	0	0.005	139
F1	F2	0.074	0.064	0.086	137
F2	F3	0.089	0.077	0.103	137
F3	F4 (Compensated Cirrhosis)	0.088	0.075	0.104	137
	Decompensated Cirrhosis	0.012	0.01	0.014	140
	Hepatocellular Carcinoma*	0.00725	0	0.02669	141
F4	Decompensated Cirrhosis	0.039	0.03	0.048	141
	Hepatocellular Carcinoma	0.019	0.017	0.055	141
Decompensate d Cirrhosis	Hepatocellular Carcinoma	0.014	0.011	0.017	140
	Liver Transplant	0.017	0.0169	0.045	142
	Death	0.129	0.1032	0.1548	141
Hepatocellular Carcinoma	Liver Transplant	0.017	0.0169	0.045	142
	Death	0.4270	0.3416	0.5124	141
Liver Transplant	Death (Year 1)	0.107	0.09	0.13	142
	Death (Year 2+)	0.0485	0.0385	0.0585	142
Background Mortality					
Source State	Target State	Base case	Lower limit	Upper limit	Referenc
CHC all-cause mortality ratio	Compared to no CHC (General population)	2.37*	1.28	4.38	143
All-cause mortality ratio after SVR	Compared to no CHC (General population)	1.4*	1.0	2.5	144
Background mortality	Death	Age-specific mortality from US 2009 Life Tables			145
Weekly cost of drugs†					
Drug		Base	Min‡	Max‡	Referenc
P 180mcg subcutaneous injection weekly		825	413	1238	146
R 1200mg daily		48	24	72	146
Simeprevir 150mg daily		5,530	2765	8295	146
Sofosbuvir 400mg daily		7,000	3500	10500	146
Ledipasvir 90mg + Sofosbuvir 400mg (daily, fixed-dose combination)		7,875	3938	11813	146
Treatment-related medical care costs (excluding drugs) §					
Service type		Base	Min	Max	Reference
Anti-HCV (antibody) test		26	13	39	147

HCV RNA quantification	79	39	118	147
Genotype assay	475	237	712	147
CBC w/Differential	14	7	22	147
Hepatic function panel	15	8	23	147
Office visit (outpatient)	97	49	146	148
Fibrosis assessment	262	131	393	149
Annual cost of CHC-related health care by disease state				
<b>Health State</b>	<b>Base</b>	<b>Min</b>	<b>Max</b>	<b>Reference</b>
F0 – No fibrosis#	810	405	3,240	150, 151
F1 – Portal Fibrosis without septa#	810	405	3,240	150, 151
F2 – Portal fibrosis with rare septa#	810	405	3,240	150, 151
F3 – Numerous septa without cirrhosis#	2,150	1,075	8,600	150, 151
F4 – Compensated cirrhosis	2,516	1,258	10,064	150, 151
Decompensated cirrhosis	29,795	27,962	31,627	142, 152
Hepatocellular carcinoma	47,525	46,653	52,392	142
Liver transplant, year 1	188,671	173,986	203,351	142
Liver transplant, year 2+	41,090	33,576	48,606	142
Post-SVR costs for F0-F3	50% of no SVR			150,151
Post-SVR costs for compensated cirrhosis	50% of no SVR			150, 151

\* – Increased by a factor of 2.37 or 1.4 for patients in F3, F4 fibrosis stages with CHC and after SVR, respectively (patients in F0-F2 stages experience the same baseline mortality as no-CHC population based on 2009 US life tables)

† – Wholesale Acquisition Cost, WAC – from Red Book Online.

‡ – The lower and upper bounds for sensitivity analyses are set at 50%-150% of base case.

§ – Cost per unit. For frequency of tests and office visits and the number of each, see Appendix Table D5.

# – F0 to F3 costs based on \$900 weighted average. The cost gradient from F0 to F3 leading into F4 costs was established using fibrosis stage prevalence shown in Appendix Table D1.

In response to treatment, the risk of progressing to worsening stages of disease is reduced.<sup>140,141</sup> It is also possible for the liver damage caused by HCV to be at least partially reversed in some patients following successful therapy (see Appendix Table D2).<sup>140, 153-157</sup> Therefore, the model assumes a proportion of patients regress to an improved fibrotic state as indicated by the proportions listed under the heading “Fibrosis Regression Post-SVR (Proportions)” in Appendix Table D2. In stages F3 and F4, patients are subject to an all-cause mortality rate that is 2.37 times the background population rate for their ages. This is reduced to 1.4 in patients achieving SVR.

## Costs

*Cost of drugs (intervention):* The weekly costs of sofosbuvir, simeprevir, and ledipasvir/sofosbuvir, peg-interferon, and ribavirin were determined using wholesale acquisition price (WAC) from Red Book Online in October 2014 (see Table 11 on the previous page).<sup>146</sup>



*Treatment-related health care costs:* The non-drug treatment-related costs shown in Table 11 are applied only for the duration of the treatment. They include HCV testing, genotyping, fibrosis staging, and therapy monitoring, including clinic visits, blood and hepatic tests, and HCV RNA quantification. See Appendix Table D5 for the frequency of these costs.

*Health care costs:* The annual medical care costs associated with the chronic hepatitis C (CHC) health states were determined from previously published research.<sup>138,140</sup> These costs were determined using Medicare reimbursement schedule and published literature.<sup>147-149</sup> Due to substantial uncertainty, we conducted wide sensitivity analyses.

*Adverse event costs:* There is limited experience with the cost of side-effect management with newer therapies. Costs were estimated by combining published cost estimates for similar events with frequencies of serious and common side-effects from clinical trials (see Table 12 below).

**Table 12: Total Treatment Costs of Associated Adverse Events, 2014 (USD)**

	Base*	Min†	Max†	
PR (48 weeks)	2073	1037	3110	Calculated
Sofosbuvir + PR (12 weeks)	1711	856	2567	Calculated
Sofosbuvir + R (24 weeks)	928	464	1392	Calculated
Ledipasvir/sofosbuvir (8 weeks)	868	434	1302	Calculated
Ledipasvir/sofosbuvir (12 weeks)	775	388	1163	Calculated
Simeprevir/sofosbuvir (12 weeks)	751	376	1127	Calculated

\* — Based on cost of serious adverse events of \$2,706 and cost of common adverse events of \$516. Costs are weighted by frequency of serious and common adverse events and summed to calculate the costs in the table

† — The lower and upper bounds for SA are set at 50%-150% of base case.

*Adjusting costs for early discontinuation:* For patients who discontinue therapy, we assumed discontinuation mid-way through the treatment and thus both the treatment costs and the costs of managing adverse events were decreased by 50%.

## Quality-of-life / Health State Utilities

*Pre- and post-SVR health state utilities:* CHC, independent of its progression to liver disease, can adversely impact patients' lives at all stages. The model uses health state utilities associated with each stage of CHC, including utilities post-SVR, and temporary loss of quality of life during treatment. These utilities represent individuals' preferences for a specific health care state associated with CHC and range from 0 (death) to 1 (normal health).<sup>158</sup> Significant decrements in quality of life accelerate as patients move from F2 to F3. The utility values are determined based on a literature review as shown in Table 13 on the next page.

**Table 13: Health State Utilities in CHC Pre-SVR and Post-SVR**

State	Base case	Lower limit	Upper limit	Reference
Utilities for HCV states				
F0	0.98	0.92	1	138, 159
F1	0.98	0.92	1	138, 159
F2	0.92	0.72	1	159
F3	0.79	0.77	0.81	160
F4 (Compensated Cirrhosis)	0.76	0.70	0.79	160
Decompensated Cirrhosis	0.69	0.44	0.69	160
Hepatocellular Carcinoma	0.67	0.60	0.72	160
Liver Transplant, Year 1	0.5	0.40	0.69	160
Liver Transplant, Year 2+	0.77	0.57	0.77	160
Death	0	0	0	
Utilities after SVR per Markov cycle				
SVR F0	1	0.98	1	138
SVR F1	1	0.98	1	138
SVR F2	0.933	0.92	1	138
SVR F3	0.86	0.82	0.90	140
SVR Compensated Cirrhosis	0.83	0.79	0.87	140

*Utility loss with treatment:* Treatment-related side-effects contribute to transient loss of quality of life. A utility penalty (or loss) due to treatment was therefore also modeled. The utility loss is calculated using utility weights of serious and common AEs weighted by the frequency of AEs reported in clinical trials and adjusted for duration of therapy.<sup>161-164</sup> The base case values of these disutilities range from -0.1782 for PR (48 weeks) to -0.0116 for LDV/SOF (8 weeks) (see Appendix Table D6).

## Calculating Results

The model produced lifetime discounted QALYs and costs to calculate incremental cost-effectiveness ratios (ICERs). Costs, QALYs gained, incremental costs, and incremental QALYs were calculated for each regimen in comparison with the next least costly regimen. ICERs by definition compare the additional costs and clinical outcomes for regimens ordered sequentially from least to most costly. This method is usually the most policy-relevant way to portray the cost-effectiveness of a set of options, provided that all of them are feasible. However, for completeness, we also included cost-effectiveness ratios in which each treatment option is compared alternatively with no treatment, as well as with PR as a universal historical control. We did this because some differences between regimen costs and efficacy are small and subject to uncertainty, making direct comparisons less definitive than comparisons to no intervention or PR. These results are displayed in tables 14 - 20 in this section of the report. The ICER for each regimen's "treat all" strategy also was calculated against "treat at F3, F4" in order to assess the cost-effectiveness of a global treatment approach versus a prioritized one.

## Scenario and Sensitivity Analyses

We portrayed scenarios in which alternative treatment discontinuation rates, the distribution of the patient cohort by fibrosis stage, the cost of care gradient from F0 to F3, and the cohort's age were altered. We documented the effect that these different, plausible values have on results. We also conducted sensitivity analyses on each of the key model inputs one at a time, to determine the model's sensitivity to the level of uncertainty with each input. The range of each variable was based on confidence intervals from published articles when these are available, as they are for example, on the probabilities of disease progression.

The confidence intervals for the SVR rates were provided by the meta-analysis described in Section 6 of this report. When formal confidence intervals were not available, as in the case of drug costs, for example, we varied each input from 50% to 150% of its base case value. To reflect the greater uncertainty in health state utility values, we adopted a wider range of 50% - 300% for those variables. To quantify the uncertainty in all inputs considered simultaneously, we carried out Monte Carlo probabilistic sensitivity analysis, using uniform distributions for all variables and 10,000 iterations. Results of the probabilistic multi-way sensitivity analyses were displayed as cost-effectiveness acceptability curves. In these figures, the X axis shows various costs per QALY gained that might be acceptable to a payer, sometimes called a "willingness-to-pay" (WTP). The Y axis shows the likelihood of any particular WTP being achieved given the range of results observed in the iterations.

## 7.3 Model of Care Value: Results

The cost-effectiveness results are presented in three parts:

1. *Results for the base-case.* "Base-case" refers to results associated with the values of input for the model that we believe are most likely to be accurate and relevant. This is further divided into sub-sections according to whether the modeled cohort was treatment-naïve; treatment-experienced; or a mixed naïve and experienced cohort; and according to whether the population strategy is "treat all" or "treat at F3, F4." The base-case analysis also reports the results of a comparison with PR (48 weeks) only, and a comparison for each treatment regimen considered separately of "treat all" versus "treat at F3, F4".
2. *"Scenario analyses".* This section presents results for different but plausible alternative values for four key inputs, in order to document how robust the base-case results are to different characteristics of the patient cohort.
3. *"Sensitivity analyses".* In this section, the values of all key inputs are altered across a wide range in order to assess the effect of uncertainty on model results.

### 7.3.1 Base-case Results

#### *Treatment-naïve cohort and “treat all” strategy*

In a prevalent, treatment-naïve cohort, PR had an ICER of \$11,385 compared with no treatment. LDV/SOF (8/12 weeks) added 1.41 QALYs compared with PR, yielding an ICER of \$20,132, well under the \$50,000 per QALY threshold to be considered highly cost-effective. All other sofosbuvir-based regimens were found to be “dominated”, meaning that the regimen both costs more and is less effective and is therefore excluded from consideration, with the exception of LDV/SOF for 12 weeks in all patients. This regimen was only slightly more effective than the 8/12 strategy (approximately three additional weeks of quality-adjusted life expectancy), but much more expensive, yielding an ICER of nearly \$300,000 per QALY gained (see Table 14 below).

**Table 14. Cost-effectiveness of Alternative Treatment Regimens for Hepatitis C, for Treatment-naïve Patients and a “Treat All” Strategy**

Strategy	Incremental comparison of regimens					
	Net cost	Incr Net cost	Eff	Incr Eff	ICER	Comment
<b>Tx naïve, treat all</b>						
No Treatment	\$ 45,313	\$ -	11.82	0.00	\$ -	undominated
PR (48 weeks)	\$ 62,540	\$ 17,227	13.34	1.51	\$ 11,385	undominated
LDV/SOF (8/12 weeks)	\$ 90,991	\$ 28,451	14.75	1.41	\$ 20,132	undominated
SOF+PR (12 weeks)	\$ 107,942	\$ 16,951	14.52	-0.23	\$ (73,572)	abs. dominated
LDV/SOF (12 weeks)	\$ 108,619	\$ 17,628	14.81	0.06	\$ 283,927	undominated
SMV/SOF (12 weeks)	\$ 163,336	\$ 54,717	14.74	-0.08	\$ (719,351)	abs. dominated
SOF+R (24 weeks)	\$ 186,513	\$ 77,894	13.99	-0.82	\$ (95,006)	abs. dominated

Incr Net Cost — Incremental Net Cost; Eff — Effectiveness; Incr Eff — Incremental Effectiveness; abs. — absolutely

#### *Treatment-naïve cohort and “treat F3, F4” strategy*

As shown in Table 15 on the next page, with ICERs of \$2,727 and \$15,940 respectively, PR and LDV/SOF 8/12, are somewhat more cost-effective if treatment is delayed until stages F3 or F4. Other regimens either have unfavorable ICERs (e.g., LDV/SOF 12 weeks) or are more costly and less effective (i.e., dominated).

**Table 15. Cost-effectiveness of Alternative Treatment Regimens for Hepatitis C, for Treatment-naïve Patients and a “Treat at F3, F4” Strategy**

Strategy	Incremental comparison of regimens					
	Net cost	Incr Net cost	Eff	Incr Eff	ICER	Comment
<b>Tx Naive, treat at F3, F4</b>						
No Treatment	\$ 45,313	\$ -	11.82	0.00	\$ -	undominated
PR (48 weeks)	\$ 48,435	\$ 3,121	12.97	1.14	\$ 2,727	undominated
LDV/SOF (8/12 weeks)	\$ 65,287	\$ 16,853	14.02	1.06	\$ 15,940	undominated
SOF+PR (12 weeks)	\$ 70,701	\$ 5,414	13.85	-0.17	\$ (31,593)	abs. dominated
LDV/SOF (12 weeks)	\$ 80,653	\$ 15,365	14.07	0.04	\$ 349,851	undominated
SMV/SOF (12 weeks)	\$ 99,733	\$ 19,080	13.98	-0.09	\$ (223,631)	abs. dominated
SOF+R (24 weeks)	\$ 115,070	\$ 34,417	13.42	-0.65	\$ (53,256)	abs. dominated

Incr Net Cost — Incremental Net Cost; Eff — Effectiveness; Incr Eff — Incremental Effectiveness; abs. — absolutely

### ***Treatment-experienced cohort and “treat all” strategy***

Net costs are somewhat higher in treatment-experienced patients compared with treatment-naïve patients in large part due to the longer regimens these patients require, while effectiveness is somewhat lower. LDV/SOF (12/24 weeks) has a very favorable ICER of \$10,200. This regimen costs more than SOF + PR (12 weeks) but added enough QALYs to have a better ICER; hence, extended dominance – while both regimens are more effective than PR alone, LDV/SOF (12/24 weeks) has a better cost-effectiveness ratio than SOF + PR (12 weeks) (see Table 16 below). Note that SOF + R is not considered an option for treatment-experienced patients.

**Table 16. Cost-effectiveness of Alternative Treatment Regimens for Hepatitis C, for Treatment-experienced Patients and a “Treat All” Strategy**

Strategy	Incremental comparison of regimens					
	Net cost	Incr Net cost	Eff	Incr Eff	ICER	Comment
<b>Tx exp, treat all</b>						
No Treatment	\$ 45,313		11.82		\$ -	
PR (48 weeks)	\$ 72,305	\$ 26,992	12.13	0.31	\$ 88,022	undominated
SOF+PR (12 weeks)	\$ 112,226	\$ 39,922	14.11	1.98	\$ 20,130	ext. dominated
LDV/SOF (12/24 weeks)	\$ 119,603	\$ 7,376	14.84	0.72	\$ 10,200	undominated
SMV/SOF (12 weeks)	\$ 165,800	\$ 46,197	14.70	-0.14	\$ (341,582)	dominated

Incr Net Cost — Incremental Net Cost; Eff — Effectiveness; Incr Eff — Incremental Effectiveness; ext. — extended

### ***Treatment-experienced cohort and “Treat F3, F4” strategy***

As shown in Table 17 on the next page, the ICER for PR is \$186,159 relative to no treatment, followed by a far more favorable ICER for LDV/SOF (12/24 weeks) of \$8,585. As in the “treat all” strategy, SOF + PR was cost-effective relative to PR alone (\$9,734 per QALY gained), but the ICER for LDV/SOF 12/24 was better (i.e., extended dominance). SMV/SOF was both more expensive and less effective than LDV/SOF 12/24 (i.e., dominated).

**Table 17. Cost-effectiveness of Alternative Treatment Regimens for Hepatitis C, for Treatment-experienced Patients and a “Treat at F3, F4” Strategy**

Strategy	Incremental comparison of regimens					
	Net cost	Incr Net cost	Eff	Incr Eff	ICER	Comment
<b>Tx exp, treat at F3, F4</b>						
No Treatment	\$ 45,313		11.82		\$ -	
PR (48 weeks)	\$ 59,873	\$ 14,560	11.90	0.08	\$ 186,159	undominated
SOF+PR (12 weeks)	\$ 75,121	\$ 15,248	13.47	1.57	\$ 9,734	ext. dominated
LDV/SOF (12/24 weeks)	\$ 80,382	\$ 5,261	14.08	0.61	\$ 8,585	undominated
SMV/SOF (12 weeks)	\$ 101,840	\$ 21,458	14.00	-0.08	\$ (276,952)	dominated

Incr Net Cost — Incremental Net Cost; Eff — Effectiveness; Incr Eff — Incremental Effectiveness; ext. — extended

### ***Comparisons with PR only***

Table 18 on the following page presents the base case results for both treatment-naïve and experienced patients and for both the “treat all” and “treat at F3, F4” strategies. However, rather than presenting incremental results for each successively more costly intervention, each regimen is compared directly with PR. For a treatment-naïve cohort, the ICERs are under \$50,000 when all patients are treated, with the exception of SMV/SOF (12 weeks), which has an ICER of \$72,038, and SOF + R (24 weeks) with an ICER of \$189,160. In the treatment-naïve and “treat at F3, F4” strategy, all ICERs were under \$50,000 except SOF + R (24 weeks) with an ICER of \$146,472.

For treatment-experienced cohorts, all sofosbuvir-containing regimens had highly favorable ICERs of under \$36,000 in the “treat all” strategy and under \$20,000 in the “treat at F3, F4” strategy when compared to PR alone. For both treatment-naïve and experienced patients, lower (more favorable) ICERs resulted from the “treat at F3, F4” strategy than from the “treat all” strategy.

**Table 18: Cost-effectiveness of Alternative Treatment Regimens for Hepatitis C, Compared to PR Alone**

	Vs. PR		
Strategy	Net cost	Incr Eff	ICER
<b>Tx naïve, treat all</b>			
No Treatment PR (48 weeks)	\$ (17,227.08)	-1.513	\$ 11,385
LDV/SOF (8/12 weeks)	\$ 28,450.78	1.413	\$ 20,132
SOF+PR (12 weeks)	\$ 45,401.89	1.183	\$ 38,386
LDV/SOF (12 weeks)	\$ 46,078.83	1.475	\$ 31,234
SMV/SOF (12 weeks)	\$ 100,795.53	1.399	\$ 72,038
SOF+R (24 weeks)	\$ 123,972.50	0.655	\$ 189,160
	Vs. PR		
Strategy	Net cost	Incr Eff	ICER
<b>Tx Naive, treat at F3, F4</b>			
No Treatment PR (48 weeks)	\$ (3,121.47)	-1.145	\$ 2,727
LDV/SOF (8/12 weeks)	\$ 16,852.92	1.057	\$ 15,940
SOF+PR (12 weeks)	\$ 22,266.46	0.886	\$ 25,134
LDV/SOF (12 weeks)	\$ 32,218.13	1.101	\$ 29,257
SMV/SOF (12 weeks)	\$ 51,298.18	1.016	\$ 50,497
SOF+R (24 weeks)	\$ 66,635.17	0.455	\$ 146,472
	Vs. PR		
Strategy	Net cost	Incr Eff	ICER
<b>Tx exp, treat all</b>			
No Treatment PR (48 weeks)	\$ (26,991.61)	-0.307	\$ 88,022
SOF+PR (12 weeks)	\$ 39,921.83	1.983	\$ 20,130
LDV/SOF (12/24 weeks)	\$ 47,297.98	2.706	\$ 17,477
SMV/SOF (12 weeks)	\$ 93,495.25	2.571	\$ 36,364
	Vs. PR		
Strategy	Net cost	Incr Eff	ICER
<b>Tx exp, treat at F3, F4</b>			
No Treatment PR (48 weeks)	\$ (14,560.10)	-0.078	\$ 186,159
SOF+PR (12 weeks)	\$ 15,247.96	1.566	\$ 9,734
LDV/SOF (12/24 weeks)	\$ 20,508.59	2.179	\$ 9,411
SMV/SOF (12 weeks)	\$ 41,967.02	2.102	\$ 19,968

Incr Eff — Incremental Effectiveness

### ***“Treat all” versus “treat at F3, F4” within regimens***

For both treatment-naïve and treatment-experienced patients, we made within-regimen comparisons of the “treat all” versus “treat at F3, F4” strategies (see Table 19 below). For each regimen, treating at all fibrosis stages was a more costly approach than treating only at F3, F4, but also yielded substantial health benefit (one-half to three-quarters of a year of quality-adjusted life expectancy for sofosbuvir-based regimens). For example, treating all naïve patients with LDV/SOF (8/12 weeks) added ~\$26,000 in lifetime costs versus a “Treat F3, F4” strategy, but also >0.7 QALYs, for an ICER of ~\$35,000 per QALY gained. Incremental costs for LDV/SOF were higher among treatment-experienced patients (where duration is 12 weeks of treatment for non-cirrhotic patients and 24 weeks for cirrhotic patients), but incremental cost-effectiveness is still approximately \$50,000 per QALY gained.

**Table 19: Cost-effectiveness of Alternative Treatment Regimens for Hepatitis C, Comparing a “Treat-All” Strategy with “Treat at F3, F4 Only”**

<b>Treat All vs. Treat at F3, F4</b>			
<b>Strategy</b>	<b>Net cost</b>	<b>Incr Eff</b>	<b>ICER</b>
<b>Tx naïve, treat all</b>			
PR (48 weeks)	\$14,106	0.368	\$ 38,282
LDV/SOF (8/12 weeks)	\$25,703	0.724	\$ 35,484
SOF+PR (12 weeks)	\$37,241	0.665	\$ 55,975
LDV/SOF (12 weeks)	\$27,966	0.743	\$ 37,663
SMV/SOF (12 weeks)	\$63,603	0.752	\$ 84,602
SOF+R (24 weeks)	\$71,443	0.569	\$ 125,577
<b>Treat All vs. Treat at F3, F4</b>			
<b>Strategy</b>	<b>Net cost</b>	<b>Incr Eff</b>	<b>ICER</b>
<b>Tx exp, treat all</b>			
PR (48 weeks)	\$12,432	0.228	\$ 54,421
SOF+PR (12 weeks)	\$37,105	0.645	\$ 57,510
LDV/SOF (12/24 weeks)	\$39,221	0.756	\$ 51,911
SMV/SOF (12 weeks)	\$63,960	0.698	\$ 91,662

Incr Net Cost — Incremental Net Cost; Eff — Effectiveness; Incr Eff — Incremental Effectiveness

### ***Combined treatment-experienced and treatment-naïve cohort***

In this comparison, we present cost-effectiveness results for a cohort containing a mix of treatment-naïve (79%) and treatment-experienced patients (21%). This is the mix reported in a recent study that examined the natural history of HCV in clinical practice, which we adjusted for those who achieved SVR.<sup>165</sup> We present only the results for the comparison of LDV/SOF to PR, given that it is



the regimen with the most favorable cost-effectiveness findings in base-case analyses. Table 20 below shows that the ICERs for LDV/SOF relative to PR (48 weeks) are highly favorable, under \$20,000 per QALY gained for both the “Treat all” and the “treat at F3, F4” strategies (\$19,229 and \$13,611, respectively). “Treat at F3, F4” is somewhat more cost-effective due to the lower total net treatment cost from delaying therapy in most individuals.

**Table 20: Cost-effectiveness of LDV/SOF vs. PR Alone in a Mixed Cohort of Treatment-naïve and Treatment-experienced Patients with Hepatitis C\***

Strategy	Vs. PR		
	Net cost	Incr Eff	ICER
Treat all			
LDV/SOF (8/12 weeks) <sup>†</sup> - LDV/SOF (12/24 weeks) <sup>‡</sup>	\$ 32,446	1.687	\$ 19,229
Treat at F3, F4			
LDV/SOF (8/12 weeks) <sup>†</sup> - LDV/SOF (12/24 weeks) <sup>‡</sup>	\$ 17,628	1.295	\$ 13,611

\* — 79.5% of patients are treatment-naïve; 20.5% treatment-experienced.

† — Regimen for treatment-naïve patients

‡ — Regimen for treatment-experienced patients

Incr Eff — Incremental Effectiveness

### 7.3.2 Scenario Analyses

In this section, we present the results associated with varying four key assumptions underpinning the model. These are (1) a higher prevalence of patients in stage F4; (2) the costs of annual medical care increase as patients progress from stages F0 – F3; (3) an increase in discontinuation rates to reflect “real world” experience; and (4) variation in the average age of the cohort. Results are presented here based on the “Treat all” strategy for treatment-naïve patients. Results of these scenario analyses, including results for treatment-experienced patients as well as the “F3, F4” strategy for all patients, are presented in Appendix Tables F1-F4.

#### ***Distribution among fibrosis stages***

In the base case, the distribution of patients across F0-F4 is 17%, 35%, 22%, 14%, and 12%, respectively. In this revised scenario, the prevalence of F4 is increased from 12% to 20% by reducing prevalence in each of the other stages by two percentage points. PR (48 weeks), LDV/SOF (8/12 weeks) and LDV/SOF (12 weeks) are the only options that are not both more costly and less effective than their comparators. Results are very similar to the base-case analysis, both in terms of comparisons of these regimens to each other as well as to the within-regimen comparisons of “treat all” vs. “treat at F3, F4”. There are a number of reasons for this relatively small change in results.

First, only 8% of individuals were reclassified to F4, leaving 92% in the same fibrosis stages. Second, the differences in the regimens are generally stable across fibrosis stages, so that their comparison is not materially affected by the modest shift in fibrosis stage distribution. Finally, the added costs and benefits of treating early continue to apply to the individuals who are still in the pre-F3 stages.

### ***Equal costs for medical care for patients in stage F0-F3***

In the base case, we assumed equal annual medical care costs for patients in stages F0 through F2 of \$810, followed by increases to \$2,150 and \$2,516 in stages F3 and F4, respectively. In this scenario, we assume equal costs for each stage of \$1,023 per year for F0 – F3, followed by the same increase to \$2,516 in F4. As with the scenario analysis above, findings were essentially identical to the base-case. This is not surprising, since annual medical care costs make up a relatively small proportion of total costs in relation to the costs of drug treatment and downstream complications.

### ***Discontinuation rates***

Discontinuation rates were increased by 50% for Interferon-based treatment in the treatment-experienced cohort and doubled for all other treatments (in both treatment-naïve and -experienced cohorts). Note that, in some instances, the meta-analysis from which the SVRs were derived resulted in a base case discontinuation rate of “0.” In such cases, for this scenario analysis, we selected the lowest non-zero value from a comparable therapy.

In this scenario, PR (48 weeks), LDV/SOF (8/12 weeks), and LDV/SOF (12 weeks) had ICERs of \$20,160, \$15,736 and \$411,658 respectively, versus \$11,385, \$20,132, and \$283,927 respectively in the base case. The relatively large change in the LDV/SOF 12-week ratio is likely due to a greater absolute difference in discontinuation rates after doubling (2.6% for the 12-week regimen vs. 0.4% for 8 weeks).

### ***Age of cohort is 50 years***

A younger cohort will have a longer average life expectancy, and thus potentially more QALYs of benefit from treatment, but also potentially higher lifetime medical care costs as more individuals live long enough to progress to more advanced disease. In this scenario, we assumed that the patients were 10 years younger than those in our base case analysis and had accordingly higher rates of disease progression.<sup>166</sup> PR (48 weeks), LDV/SOF (8/12 weeks), and LDV/SOF (12 weeks) had ICERs of \$5,141, \$12,562, and \$201,418, respectively. Cost-effectiveness of the “treat all” vs. “treat at F3, F4” was somewhat improved, however, as a result of greater slowing of disease progression with effective treatment.

### 7.3.3 Sensitivity Analyses

Both one-way and multi-way sensitivity results are presented for treatment-naïve patients in this section; we did not conduct similar analyses for treatment-experienced patients given the similarity in base-case results. Under each of these headings, results for the “treat all” approach are presented first, followed by results when treatment is initiated only at stages F3 and F4. In the “Tornado diagrams”, we present only those variables that significantly affected results.

#### *One-way sensitivity analyses*

We present the results of one-way sensitivity analyses by means of tornado diagrams. These diagrams show the low to high range of ICER values for uncertainty in each variable, over the range displayed in the legend. The longer the bar associated with each variable, the greater its influence on the ICER. Only the 12 most influential input variables are displayed.

Importantly, none of the variations in parameter estimates we tested resulted in an incremental cost-effectiveness ratio above \$50,000 per QALY gained. For example, for the treatment-naïve, “treat all” strategy, the ICER of LDV/SOF (8/12 weeks) versus PR (48 weeks) varied from “cost saving” (i.e., more effective, less expensive) to approximately \$48,000 per QALY gained as the weekly drug cost varied from \$3,937 to \$11,812. The weekly cost of drugs for PR (48 weeks) had the second-largest effect on the ICER. Other inputs had much smaller effects (see Figure 6 on the following page); for example, varying the percentage of patients eligible for the 8-week LDV/SOF regimen from 30% to 90% caused the ICER to range from \$15,000 to \$26,000 per QALY gained relative to PR alone.

Comparing the same regimens of LDV/SOF (8/12 weeks) versus PR (48 weeks) but assuming a “treat at F3, F4” strategy, the weekly cost of drugs remained the most important variables in determining cost-effectiveness (see Figure 7 on page 63).

Figure 6: One-way Sensitivity Analyses for Treatment-naïve Patients and “Treat All” Strategy





Figure 7: One-way Sensitivity Analyses for Treatment-naïve Patients and “Treat at F3, F4” Strategy




### ***Multi-way probabilistic sensitivity analyses***

Multi-way sensitivity analyses are presented by means of cost-effectiveness acceptability curves. We show the distribution of ICERs across 10,000 model runs for LDV/SOF (8/12 weeks) versus PR (48 weeks), varying the base case assumptions for all variables in the model. We used the same range of the input variables employed in the one-way sensitivity analyses (i.e., either published confidence intervals, or 50% - 150% of the base case value if confidence intervals were unavailable). In Figure 8 below and Figure 9 on page 65, the horizontal axis represents the ICERs for LDV/SOF vs. PR, which can be taken to represent possible levels at which a health care system is “willing to pay” for the additional health gain of a QALY. The vertical axis is the percent of the model runs that produced an ICER at or below that particular level, indicating the percent likelihood that LDV/SOF would be considered “cost-effective” at that particular willingness to pay level.

#### ***Treat all***

Under this strategy, approximately 98% of the simulations yielded an acceptable cost-effectiveness ratio at a willingness to pay threshold of \$50,000 per QALY gained, suggesting that the finding that LDV/SOF is cost-effective at that threshold is robust. At \$150,000 per QALY gained, effectively 100% of the simulations would yield an acceptable ICER (see Figure 8 below).


**Figure 8: Cost-effectiveness Acceptability Curve for LDV/SOF 8/12 weeks, Treatment-naïve, “Treat All” (Compared to PR Alone)**



### ***Treat at F3 and F4 only***

Similar to the “treat all” strategy, over 99% of simulations for the “treat at F3, F4” strategy also yield an ICER of \$50,000 or less (see Figure 9 below).

**Figure 9: Cost-effectiveness Acceptability Curve for LDV/SOF 8/12 weeks, Treatment-naïve, “Treat at F3, F4 Only” (Compared to PR Alone)**



## **7.4 Health-System Value Analysis: Methods**

As mentioned in the beginning of this section, we also assessed the potential budgetary impact of new hepatitis C therapy over three periods of follow-up: one, five, and 20 years after treatment initiation. As with the cost-effectiveness analyses, the regimen of interest for genotype 1 was the LDV/SOF strategy (8/12 weeks for treatment-naïve, 12/24 for treatment-experienced), as this represents the cost-effective strategy that is currently available and most likely to receive widespread use. For each of these time points, we used outputs from the care value model to inform expected numbers (per 1,000 treated) of patients experiencing HCV-related complications (cirrhosis, decompensated cirrhosis, hepatocellular carcinoma, liver transplant) and dying of HCV-related causes. Costs of treatment and all other care were calculated on a per patient basis, as were total costs. Results for treatment-naïve and treatment-experienced patients were combined and weighted according to an assumed distribution of 79% and 21% for these two subpopulations respectively, as used in the care value analysis.<sup>176</sup>

We then combined these results with findings from the initial CTAF review for genotypes 2 and 3<sup>180</sup> to assess the one-, five-, and 20-year budgetary impact to California state agencies (Medi-Cal and the Department of Corrections) of adopting LDV/SOF for genotype 1 and the most effective therapies available for genotypes 2 and 3 (SOF + R for 12 weeks for genotype 2 and 24 weeks for genotype 3). The number of individuals with chronic hepatitis C who rely on these state entities was recently estimated to total 93,000,<sup>177</sup> of which 70%, 16%, and 12% were assumed to have genotypes 1, 2, and 3 respectively.<sup>178</sup>

Finally, we conducted analyses to examine the drug prices at which benchmark thresholds of insurer premium increases would not be crossed. In conversation with a variety of health plan professionals and pharmacy benefit managers, we were advised that these thresholds tend to fall in the range of a 0.5-1.0% increase in the per-member per-month (PMPM) premium. Payers believe that the introduction of a single intervention that could potentially cause an increase in PMPM beyond this level requires some form of management in order to modulate the immediate budget impact. If a budget impact of this magnitude cannot be managed, payers believe that there is a significant likelihood that care of equal or greater value will be displaced and/or that health insurance premiums will rise in a fashion that would adversely affect access to affordable care for all patients. The base PMPM was assumed to be \$611, based on a recent reporting of Medi-Cal rates from the state Department of Health Care Services (DHCS).

In addition to a full analysis of the prevalent population, the latter two analyses were also conducted under a scenario in which only those currently at F3 and F4 would be prioritized for treatment. All budget impact analyses were conducted using Microsoft Excel.



## 7.5 Health-System Value Analysis: Results

### Budgetary Impact: Per 1,000 Patients Treated

Findings for the performance of LDV/SOF vs. PR are presented in Table 21 below; results are weighted for the combined treatment-naïve and treatment-experienced populations (individual results for these populations are presented in Appendix Tables G1 and G2). As shown in the table, LDV/SOF produces incremental clinical benefits very soon after treatment initiation; for example, compared with PR alone, LDV/SOF prevents approximately six cases of cirrhosis and two HCV-related deaths per 1,000 patients treated in the first year alone.

**Table 21. Clinical Outcomes (per 1,000 patients treated) and Costs for LDV/SOF and PR Therapy over One, Five, and 20 Years of Follow-up**

Timeframe/Regimen	Liver-Related Complications				HCV	Costs (per patient, \$)		
	Cirrhosis	Decompensation	HCC	Transplant	Death	Treatment	Other	Total
<u>1 Year</u>								
PR	6.8	3.5	1.8	0.0	5.4	\$34,966	\$1,636	\$36,602
LDV/SOF	0.8	0.6	1.2	0.0	3.4	\$84,341	\$696	\$85,037
Difference (LS-PR)	(5.9)	(3.0)	(0.6)	0.0	(2.0)	\$49,375	(\$940)	\$48,435
<u>5 Years</u>								
PR	34.8	18.7	11.9	0.4	35.3	\$34,966	\$6,681	\$41,647
LDV/SOF	6.1	3.4	6.7	0.3	18.7	\$84,341	\$3,260	\$87,601
Difference (LS-PR)	(28.8)	(15.3)	(5.1)	(0.1)	(16.5)	\$49,375	(\$3,421)	\$45,954
<u>20 Years</u>								
PR	120.9	66.8	45.3	4.9	248.8	\$34,966	\$23,442	\$58,409
LDV/SOF	21.5	11.8	23.0	1.5	109.1	\$84,341	\$10,214	\$94,555
Difference (LS-PR)	(99.4)	(55.0)	(22.3)	(3.3)	(139.7)	\$49,375	(\$13,229)	\$36,146

LS-PR: Difference between LDV/SOF and PR therapy

However, treatment costs are more than doubled with the newer regimen, and only a small portion of costs are offset by reduced complications. The incremental cost required to avert one HCV-related death at one year is approximately \$24 million (data not shown).

Benefits are more fully realized at later time points. At five years, LDV/SOF would avoid 44 cases of cirrhosis (15 of which would be decompensated), five cases of HCC, and 17 HCV-related deaths per 1,000 treated. Cost offsets would total approximately 7% of incremental treatment costs, but the cost to prevent one HCV-related death would still be nearly \$3 million. At 20 years, there would be a nearly six-fold reduction in the incidence of cirrhosis, HCC incidence would be reduced by about half, and 140 HCV-related deaths would be averted per 1,000 treated. Over 25% of treatment costs would be offset by these reductions, and the cost per HCV death averted would be reduced to \$260,000.

### **Budgetary Impact: Medi-Cal/Department of Corrections Population**

Our estimates of the budgetary impact of adoption of new hepatitis C treatments to Medi-Cal and the Department of Corrections is summarized in Figure 10 on the following page and in detail in Appendix Table G3. As described previously, LDV/SOF 8/12 or 12/24 was assumed to be the therapy of choice for genotype 1, while SOF + R for 12 weeks and 24 weeks was assumed for genotypes 2 and 3, respectively. A total of 91,140 of the 93,000 total patients would have chronic hepatitis C and genotypes 1, 2, and 3, 50% of whom would be expected to be aware of infection and present for treatment (n=45,570). Total health plan expenditures for all medical care would be approximately \$56 billion (i.e., \$611 PMPM).

The availability of new HCV treatments would increase costs by approximately \$1.6 billion, \$545 million, and \$901 million for genotypes 1, 2, and 3 respectively (see Figure 10), resulting in a total increase of \$3 billion, or \$33 PMPM. This represents a 5% increase over the base PMPM of \$611. Cost offsets after five years would total \$254 million, reducing net expenditures modestly to \$2.8 billion. More substantial offsets after 20 years (\$1.2 billion) would reduce net expenditures further to \$1.8 billion.

**Figure 10. Budgetary Impact of New Hepatitis C Treatments in the Medi-Cal/Department of Corrections Hepatitis C Population in California, with and without Cost Offsets from Reduced Liver-related Complications**





Figure 11 on the following page illustrates the budgetary impact with treatment commenced only for patients at fibrosis levels of F3 and F4 (approximately one-quarter of the potential patient pool). The initial expenditures for new therapies are reduced to approximately \$800 million (~\$9 PMPM, a 1.4% increase). Total net expenditures after 20 years are \$475 million, an increase of less than 1%.

**Figure 11. Budgetary Impact of New Hepatitis C Treatments in the Medi-Cal/Department of Corrections Hepatitis C Population in California, with and without Cost Offsets from Reduced Liver-related Complications: Treatment of Patients Currently at F3 and F4 Only**



### Drug Pricing to Meet Per-Member Per-Month Benchmarks

As mentioned previously, PMPM increases of 0.5%-1.0% in a given year were used in this report as a range of potential budget impact that is likely to warrant specific efforts to manage the costs of a new health care intervention. We examined the incremental drug expenditures at which PMPM increases of 0.5% and 1.0% would be met for genotype 1, the patient subpopulation of interest in this review. Historical treatment costs were estimated based on the cost of PR (approximately \$42,000 per treatment course) weighted by the assumed proportion of patients eligible for such therapy (60%); no treatment at baseline was assumed for the 40% of patients who would be ineligible for interferon-based therapy. Thus, historical treatment costs were estimated to total approximately \$25,000 per patient with genotype 1 disease.

Based on the assumed baseline PMPM in this analysis (\$611) as well as the size of the population to be treated (approximately 33,000 patients in the Medi-Cal/Department of Corrections population in California if 50% of genotype 1 patients present for treatment), a course of treatment with a new agent would need to be priced at \$34,000 - \$42,000 to meet the 0.5% and 1% thresholds respectively.

We also conducted a hypothetical analysis of the number of treatment-naïve Medi-Cal/Department of Corrections patients who could be treated without exceeding a 1% PMPM threshold, based on the current wholesale acquisition costs of LDV/SOF (approximately \$63,000 and \$95,000 for 8 and 12 weeks, respectively). As with other model analyses, we assumed that 79% of genotype 1 patients presenting for treatment would be treatment-naïve (i.e., ~26,000 of 33,000 in the Medi-Cal/Department of Corrections population), and that 67% of treatment-naïve non-cirrhotic patients would receive 8 weeks of treatment.

Based on these assumptions, only two-thirds of these patients (approximately 16,500 of the 26,000 patients presenting for care) could receive treatment at these prices if the one-year PMPM increase were to be held to less than 1% (i.e.,  $\leq \$6.11$ ), leaving nearly 10,000 Medi-Cal/Department of Corrections patients without access to new therapy. When considering a 0.5% threshold for PMPM increase ( $\leq \$3.06$ ), less than half of eligible patients (12,600 of 26,000) could be treated at current prices.

We conducted an alternative analysis in which the percentage of treatment-naïve non-cirrhotic patients eligible for 8 weeks of therapy was adjusted upward to 90%. Even with this adjustment, the percentages of genotype 1 patients who could receive treatment increases to only 54% and 71% at the 0.5% and 1% PMPM thresholds respectively, leaving nearly 12,000 and 8,000 patients without access to treatment.

By contrast, if the population of treatment-naïve genotype 1 patients is restricted to those with F3 and F4 stage disease ( $n \sim 6,700$ ), LDV/SOF could replace historical PR therapy in all of these patients at current prices and remain under the 1% threshold for PMPM increase. When considering a 0.5% increase in PMPM ( $\$3.06$ ), LDV/SOF could replace PR in 91% of F3/F4 patients ( $n \sim 6,100$ ) at current prices. *(Note: if the percentage eligible for 8-week therapy is increased to 90%, then all F3/F4 patients could be treated below the 0.5% PMPM increase threshold.)*

## 7.6 Summary

Using the best available information on the costs and health consequences of drug therapies for the most common form of chronic hepatitis C (genotype 1), we modeled the net costs, health benefits (expressed in QALYs), and incremental cost-effectiveness of a range of sofosbuvir-based therapies as well as pegylated interferon and ribavirin alone. We also assessed these results in a comparison of a policy of treating HCV patients in all fibrosis stages against a policy of treating only those who reach F3 and F4, thus delaying the treatment for those initially in stages F0-F2. While estimates of what might be considered cost-effective vary, it is reasonable to rate an ICER of under \$150,000 to be “cost-effective” and ICERs under \$50,000 to be “very cost-effective”. In the base-case analysis we found that LDV/SOF regimens for treatment-naïve and treatment-experienced patients were very cost-effective, producing ICERs  $\leq \$20,000$  per QALY gained regardless of the comparison (e.g.,

PR alone vs. next-least costly alternative, treat all vs. treat at F3, F4, weighted estimates for a combined treatment-naïve and treatment-experienced cohort).

Our analysis also found that, while treating patients at all fibrosis stages was more expensive in comparison to waiting to treat until patients reached F3 or F4, it was also more effective. For example, treating all naïve patients with LDV/SOF 8/12 or LDV/SOF 12 as well as PR alone produced ICERs <\$40,000 per QALY gained in comparison to treating only at F3/F4. Among treatment-experienced patients, differences in effectiveness were more pronounced, with over two years of quality-adjusted life expectancy gained for sofosbuvir-based regimens relative to PR alone (generating ICERs of \$10,000-\$20,000 per QALY gained). Comparisons of the “treat all” vs. “treat at F3, F4” approaches in the treatment-experienced subgroup generated more costs (in part because sofosbuvir-based regimens are longer) but still produced estimates of cost-effectiveness of ~\$50,000 per QALY gained. Model findings were robust to a range of sensitivity analyses, with changes in model results greatest in relation to variation in the weekly prices of sofosbuvir and PR therapy.

These findings stand in contrast, however, to those of our budget impact analysis, which suggest that the introduction of LDV/SOF would increase the cost of treatment over PR alone by \$40,000-\$75,000 per patient depending on the duration of therapy. Some of these costs would be offset by reductions in the rate of serious liver complications but would offset 30-40% of additional treatment costs at most. As a result, the budgetary impact to the nearly 100,000 Californians being treated for HCV with state funds (i.e., Medi-Cal and Department of Corrections) would be substantial. Treatment costs would increase by \$1.6 billion for genotype 1 alone if 50% of infected patients are treated; when estimates for genotypes 2 and 3 from our March 2014 report are included, the total budgetary impact would be over \$3 billion, or \$33 per member per month (PMPM).

Based on a recent estimate of PMPM costs for Medi-Cal (\$611), this represents a 5% increase, far above the 0.5-1% increase that most insurers believe is the upper limit for a manageable increase in expenditures. This increase is reduced somewhat when downstream cost offsets are considered, but never approaches the 0.5-1% threshold. In fact, a new agent would need to be priced at \$34,000 - \$42,000 per course of treatment to fall within this range (approximately \$9,000-\$17,000 above the baseline cost of PR therapy). At current prices, LDV/SOF 8/12 could only be offered to approximately half of eligible patients presenting for treatment. If treatment were restricted only to patients at fibrosis stages F3 and F4, however, the budgetary impact is less pronounced. Treatment costs would rise by approximately \$800 million in the Medi-Cal/Department of Corrections population (~\$9 PMPM, a 1.4% increase) and would be \$475 million after 20-year cost offsets were considered.

We note some limitation of our analyses. First, we did not model the effects of HCV treatments on patients co-infected with HIV or in those treated following liver transplant. Clinical consequences and costs might be very different in these small but important subgroups. The analytic perspective

was that of a third-party payer, and we therefore did not include the costs of transportation or other incidental costs associated with seeking and obtaining medical care, nor did we incorporate patients' financial contributions (e.g., copayments, deductibles) into these calculations. The FDA approval for the combination of sofosbuvir and simeprevir came after our analyses had been completed; as such, our modeled duration of therapy in treatment-experienced individuals was half that of the approved duration (12 vs. 24 weeks). While adjustment of treatment duration would have increased the cost of treatment for this combination in treatment-experienced individuals, it would not have appreciably changed major findings, namely that SMV/SOF is less effective and more expensive than LDV/SOF regimens.

We also did not include the benefits resulting from reduced secondary transmission of HCV due to reduced community HCV burden, nor did we model the risk of re-infection or relapse following SVR and the associated costs and health outcomes. The simplified "snapshot" approach in the budget impact analysis also did not consider relapse, reinfection, or even incident infection in patients not treated at baseline. Finally, we obtained data from a variety of sources, many of them not perfectly suited to the demands of our models. For example, estimates of effectiveness as measured by SVR were derived from clinical trial results. "Real world" effectiveness might diverge significantly from these estimates.

Nevertheless, our findings have important implications. In particular, model results suggest that the introduction of LDV/SOF for both treatment-naïve and treatment-experienced individuals would confer substantial clinical benefits in comparison to historical treatment standards and even in relation to other sofosbuvir-based regimens. While the use of this new regimen would increase treatment costs, such use appears to be cost-effective. However, the additional expenditures required to treat all patients with genotype 1 infection (even if only 50% of them are aware of their infection) are substantial; when added to the additional expenditures already required for genotypes 2 and 3, this represents a per-member per-month premium increase that is fivefold higher than frequently-discussed manageable thresholds for new interventions. It is clear that patients, physicians, insurers, and health systems will have to grapple with the budget impact of new, highly effective, and expensive treatments for hepatitis C. Whether this will result in prioritization of clinical care, new contracting and financing tactics, evolving market dynamics, or policy actions remains to be seen.

\*\*\*\*

**This is the first review of these technologies by the California Technology Assessment Forum and the second review of treatment alternatives for chronic hepatitis C.**

# References

---

1. Kim WR. The burden of hepatitis C in the United States. *Hepatology (Baltimore, Md.)*. 2002; 36(5 Suppl 1):S30-34.
2. United States Department of Veterans Affairs. National Hepatitis C Program Office. Interferon and Ribavirin Treatment Side Effects. <http://www.hepatitis.va.gov/provider/reviews/treatment-side-effects.asp>. Accessed January 8, 2014.
3. Butt AA, Kanwal F. Boceprevir and telaprevir in the management of hepatitis C virus-infected patients. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*. 2012; 54(1):96-104.
4. Hagan LM, Yang Z, Ehteshami M, Schinazi RF. All-oral, interferon-free treatment for chronic hepatitis C: cost-effectiveness analyses. *Journal of Viral Hepatitis*. 2013; 20(12):847-857.
5. Reau NS, Jensen DM. Sticker shock and the price of new therapies for hepatitis C: Is it worth it? *Hepatology (Baltimore, Md.)*. 2014.
6. Hepatitis C--global prevalence (update). *Releve epidemiologique hebdomadaire / Section d'hygiene du Secretariat de la Societe des Nations = Weekly epidemiological record / Health Section of the Secretariat of the League of Nations*. 1999; 74(49):425-427.
7. Armstrong GL, Wasley A, Simard EP, McQuillan GM, Kuhnert WL, Alter MJ. The prevalence of hepatitis C virus infection in the United States, 1999 through 2002. *Annals of Internal Medicine*. 2006; 144(10):705-714.
8. Chak E, Talal AH, Sherman KE, Schiff ER, Saab S. Hepatitis C virus infection in USA: an estimate of true prevalence. *Liver International: Official Journal of the International Association for the Study of the Liver*. 2011; 31(8):1090-1101.
9. Holmberg SD, Spradling PR, Moorman AC, Denniston MM. Hepatitis C in the United States. *The New England Journal of Medicine*. 2013; 368(20):1859-1861.
10. Kershenobich D, Razavi HA, Cooper CL, et al. Applying a system approach to forecast the total hepatitis C virus-infected population size: model validation using US data. *Liver International: Official Journal of the International Association for the Study of the Liver*. 2011; 31 Suppl 2:4-17.
11. Chinnadurai R, Velazquez V, Grakoui A. Hepatic transplant and HCV: a new playground for an old virus. *American Journal of Transplantation: Official Journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2012; 12(2):298-305.
12. Simmonds P, Holmes EC, Cha TA, et al. Classification of hepatitis C virus into six major genotypes and a series of subtypes by phylogenetic analysis of the NS-5 region. *The Journal of General Virology*. 1993; 74 ( Pt 11):2391-2399.



13. Backus LI, Boothroyd DB, Phillips BR, Belperio P, Halloran J, Mole LA. A sustained virologic response reduces risk of all-cause mortality in patients with hepatitis C. *Clinical Gastroenterology and Hepatology: The Official Clinical Practice Journal of the American Gastroenterological Association*. 2011; 9(6):509-516.e501.
14. Blatt LM, Mutchnick MG, Tong MJ, et al. Assessment of hepatitis C virus RNA and genotype from 6807 patients with chronic hepatitis C in the United States. *Journal of Viral Hepatitis*. 2000; 7(3):196-202.
15. Germer JJ, Mandrekar JN, Bendel JL, Mitchell PS, Yao JD. Hepatitis C virus genotypes in clinical specimens tested at a national reference testing laboratory in the United States. *Journal of Clinical Microbiology*. 2011; 49(8):3040-3043.
16. Manos MM, Shvachko VA, Murphy RC, Arduino JM, Shire NJ. Distribution of hepatitis C virus genotypes in a diverse US integrated health care population. *Journal of Medical Virology*. 2012; 84(11):1744-1750.
17. Nainan OV, Alter MJ, Kruszon-Moran D, et al. Hepatitis C virus genotypes and viral concentrations in participants of a general population survey in the United States. *Gastroenterology*. 2006; 131(2):478-484.
18. Zein NN, Rakela J, Krawitt EL, Reddy KR, Tominaga T, Persing DH. Hepatitis C virus genotypes in the United States: epidemiology, pathogenicity, and response to interferon therapy. Collaborative Study Group. *Annals of Internal Medicine*. 1996; 125(8):634-639.
19. Denniston MM, Klevens RM, McQuillan GM, Jiles RB. Awareness of infection, knowledge of hepatitis C, and medical follow-up among individuals testing positive for hepatitis C: National Health and Nutrition Examination Survey 2001-2008. *Hepatology (Baltimore, Md.)*. 2012; 55(6):1652-1661.
20. Moorman AC, Gordon SC, Rupp LB, et al. Baseline characteristics and mortality among people in care for chronic viral hepatitis: the chronic hepatitis cohort study. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*. 2013; 56(1):40-50.
21. Moyer VA. Screening for hepatitis C virus infection in adults: U.S. Preventive Services Task Force recommendation statement. *Annals of Internal Medicine*. 2013; 159(5):349-357.
22. Smith BD, Morgan RL, Beckett GA, Falck-Ytter Y, Holtzman D, Ward JW. Hepatitis C virus testing of persons born during 1945-1965: recommendations from the Centers for Disease Control and Prevention. *Annals of Internal Medicine*. 2012; 157(11):817-822.
23. Farci P, Alter HJ, Wong D, et al. A long-term study of hepatitis C virus replication in non-A, non-B hepatitis. *The New England Journal of Medicine*. 1991; 325(2):98-104.
24. Alter MJ, Margolis HS, Krawczynski K, et al. The natural history of community-acquired hepatitis C in the United States. The Sentinel Counties Chronic non-A, non-B Hepatitis Study Team. *The New England Journal of Medicine*. 1992; 327(27):1899-1905.

25. Barrera JM, Bruguera M, Ercilla MG, et al. Persistent hepatitis C viremia after acute self-limiting posttransfusion hepatitis C. *Hepatology (Baltimore, Md.)*. 1995; 21(3):639-644.
26. Thein HH, Yi Q, Dore GJ, Krahn MD. Estimation of stage-specific fibrosis progression rates in chronic hepatitis C virus infection: a meta-analysis and meta-regression. *Hepatology (Baltimore, Md.)*. 2008; 48(2):418-431.
27. Freeman AJ, Dore GJ, Law MG, et al. Estimating progression to cirrhosis in chronic hepatitis C virus infection. *Hepatology (Baltimore, Md.)*. 2001; 34(4 Pt 1):809-816.
28. Degenhardt L, Randall D, Hall W, Law M, Butler T, Burns L. Mortality among clients of a state-wide opioid pharmacotherapy program over 20 years: risk factors and lives saved. *Drug and Alcohol Dependence*. 2009; 105(1-2):9-15.
29. Grebely J, Dore GJ. What is killing people with hepatitis C virus infection? *Seminars in Liver Disease*. 2011; 31(4):331-339.
30. Grebely J, Raffa JD, Lai C, et al. Impact of hepatitis C virus infection on all-cause and liver-related mortality in a large community-based cohort of inner city residents. *Journal of Viral Hepatitis*. 2011; 18(1):32-41.
31. Bonis PA, Tong MJ, Blatt LM, Conrad A, Griffith JL. A predictive model for the development of hepatocellular carcinoma, liver failure, or liver transplantation for patients presenting to clinic with chronic hepatitis C. *The American Journal of Gastroenterology*. 1999; 94(6):1605-1612.
32. Ghany MG, Kim HY, Stoddard A, Wright EC, Seeff LB, Lok AS. Predicting clinical outcomes using baseline and follow-up laboratory data from the hepatitis C long-term treatment against cirrhosis trial. *Hepatology (Baltimore, Md.)*. 2011; 54(5):1527-1537.
33. Ghany MG, Lok AS, Everhart JE, et al. Predicting clinical and histologic outcomes based on standard laboratory tests in advanced chronic hepatitis C. *Gastroenterology*. 2010; 138(1):136-146.
34. Tong MJ, el-Farra NS, Reikes AR, Co RL. Clinical outcomes after transfusion-associated hepatitis C. *The New England Journal of Medicine*. 1995; 332(22):1463-1466.
35. Graham CS, Baden LR, Yu E, et al. Influence of human immunodeficiency virus infection on the course of hepatitis C virus infection: a meta-analysis. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*. 2001; 33(4):562-569.
36. Hourigan LF, Macdonald GA, Purdie D, et al. Fibrosis in chronic hepatitis C correlates significantly with body mass index and steatosis. *Hepatology (Baltimore, Md.)*. 1999; 29(4):1215-1219.
37. Hutchinson SJ, Bird SM, Goldberg DJ. Influence of alcohol on the progression of hepatitis C virus infection: a meta-analysis. *Clinical Gastroenterology and Hepatology: The Official Clinical Practice Journal of the American Gastroenterological Association*. 2005; 3(11):1150-1159.

38. Minola E, Prati D, Suter F, et al. Age at infection affects the long-term outcome of transfusion-associated chronic hepatitis C. *Blood*. 2002; 99(12):4588-4591.
39. Poynard T, Bedossa P, Opolon P. Natural history of liver fibrosis progression in patients with chronic hepatitis C. The OBSVIRC, METAVIR, CLINIVIR, and DOSVIRC groups. *Lancet*. 1997; 349(9055):825-832.
40. Ratziu V, Munteanu M, Charlotte F, Bonyhay L, Poynard T. Fibrogenic impact of high serum glucose in chronic hepatitis C. *Journal of Hepatology*. 2003; 39(6):1049-1055.
41. Fried MW, Buti M, Dore GJ, et al. Once-daily simeprevir (TMC435) with pegylated interferon and ribavirin in treatment-naïve genotype 1 hepatitis C: The randomized PILLAR study. *Hepatology (Baltimore, Md.)*. 2013; 58(6):1918-1929.
42. Thorlund K, Druyts E, Mills EJ. SVR12 is higher than SVR24 in treatment-naïve hepatitis C genotype 1 patients treated with peginterferon plus ribavirin. *Clinical Epidemiology*. 2014; 6(1):49-58.
43. Yoshida EM, Sulkowski MS, Gane EJ, et al. Concordance of Sustained Virologic Response 4, 12, and 24 Weeks Post-Treatment With Sofosbuvir-Containing Regimens for Hepatitis C Virus. *Hepatology (Baltimore, Md.)*. 2014.
44. Ng V, Saab S. Effects of a sustained virologic response on outcomes of patients with chronic hepatitis C. *Clinical Gastroenterology and Hepatology: The Official Clinical Practice Journal of the American Gastroenterological Association*. 2011; 9(11):923-930.
45. Hezode C, Fontaine H, Dorival C, et al. Triple therapy in treatment-experienced patients with HCV-cirrhosis in a multicentre cohort of the French Early Access Programme (ANRS CO20-CUPIC) - NCT01514890. *Journal of Hepatology*. 2013; 59(3):434-441.
46. Brennan TA, Lotvin A, Shrank W. Analysis of "real world" Sovaldi (sofosbuvir) use and discontinuation rates. *CVS Health*. 2014.  
<http://www.cvshealth.com/sites/default/files/hepatitisCutilization.pdf>
47. Nelson D. Real world experience with DAA regimens. AASLD/EASL Special Conference on Hepatitis C; 2014; New York City, NY.
48. Fried MW, Shiffman ML, Reddy KR, et al. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *The New England Journal of Medicine*. 2002; 347(13):975-982.
49. Kramer JR, Kanwal F, Richardson P, Mei M, El-Serag HB. Gaps in the achievement of effectiveness of HCV treatment in national VA practice. *Journal of Hepatology*. 2012; 56(2):320-325.
50. Zeuzem S, Hultcrantz R, Bourliere M, et al. Peginterferon alfa-2b plus ribavirin for treatment of chronic hepatitis C in previously untreated patients infected with HCV genotypes 2 or 3. *Journal of Hepatology*. 2004; 40(6):993-999.
51. Gaetano JN, Reau N. Hepatitis C: management of side effects in the era of direct-acting antivirals. *Current Gastroenterology Reports*. 2013; 15(1):305.

52. Ghany MG, Nelson DR, Strader DB, Thomas DL, Seeff LB. An update on treatment of genotype 1 chronic hepatitis C virus infection: 2011 practice guideline by the American Association for the Study of Liver Diseases. *Hepatology (Baltimore, Md.)*. 2011; 54(4):1433-1444.
53. Myers RP, Ramji A, Bilodeau M, Wong S, Feld JJ. An update on the management of hepatitis C: consensus guidelines from the Canadian Association for the Study of the Liver. *Canadian Journal of Gastroenterology = Journal Canadien de Gastroenterologie*. 2012; 26(6):359-375.
54. Ramachandran P, Fraser A, Agarwal K, et al. UK consensus guidelines for the use of the protease inhibitors boceprevir and telaprevir in genotype 1 chronic hepatitis C infected patients. *Alimentary Pharmacology & Therapeutics*. 2012; 35(6):647-662.
55. Hezode C. Boceprevir and telaprevir for the treatment of chronic hepatitis C: safety management in clinical practice. *Liver International: Official Journal of the International Association for the Study of the Liver*. 2012; 32 Suppl 1:32-38.
56. Reesink HW, Zeuzem S, Weegink CJ, et al. Rapid decline of viral RNA in hepatitis C patients treated with VX-950: a phase Ib, placebo-controlled, randomized study. *Gastroenterology*. 2006; 131(4):997-1002.
57. Susser S, Welsch C, Wang Y, et al. Characterization of resistance to the protease inhibitor boceprevir in hepatitis C virus-infected patients. *Hepatology (Baltimore, Md.)*. 2009; 50(6):1709-1718.
58. Lawitz E, Sulkowski MS, Ghalib R, et al. Simeprevir plus sofosbuvir, with or without ribavirin, to treat chronic infection with hepatitis C virus genotype 1 in non-responders to pegylated interferon and ribavirin and treatment-naïve patients: the COSMOS randomised study. *Lancet*. 2014.
59. Sulkowski MS, Gardiner DF, Rodriguez-Torres M, et al. Daclatasvir plus sofosbuvir for previously treated or untreated chronic HCV infection. *The New England Journal of Medicine*. 2014; 370(3):211-221.
60. Forns X, Lawitz E, Zeuzem S, et al. Simeprevir with peginterferon and ribavirin leads to high rates of SVR in patients with HCV genotype 1 who relapsed after previous therapy: a phase 3 trial. *Gastroenterology*. 2014; 146(7):1669-1679.e1663.
61. Jacobson IM, Dore GJ, Foster GR, et al. Simeprevir with pegylated interferon alfa 2a plus ribavirin in treatment-naïve patients with chronic hepatitis C virus genotype 1 infection (QUEST-1): a phase 3, randomised, double-blind, placebo-controlled trial. *Lancet*. 2014; 384(9941):403-413.
62. Manns M, Marcellin P, Poordad F, et al. Simeprevir with pegylated interferon alfa 2a or 2b plus ribavirin in treatment-naïve patients with chronic hepatitis C virus genotype 1 infection (QUEST-2): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet*. 2014; 384(9941):414-426.

63. Zeuzem S, Berg T, Gane E, et al. Simeprevir increases rate of sustained virologic response among treatment-experienced patients with HCV genotype-1 infection: a phase IIb trial. *Gastroenterology*. 2014; 146(2):430-441.e436.
64. Hayashi N, Izumi N, Kumada H, et al. Simeprevir with peginterferon/ribavirin for treatment-naive hepatitis C genotype 1 patients in Japan: CONCERTO-1, a phase III trial. *Journal of Hepatology*. 2014; 61(2):219-227.
65. Hayashi N, Seto C, Kato M, Komada Y, Goto S. Once-daily simeprevir (TMC435) with peginterferon/ribavirin for treatment-naive hepatitis C genotype 1-infected patients in Japan: the DRAGON study. *Journal of Gastroenterology*. 2014; 49(1):138-147.
66. Izumi N, Hayashi N, Kumada H, et al. Once-daily simeprevir with peginterferon and ribavirin for treatment-experienced HCV genotype 1-infected patients in Japan: the CONCERTO-2 and CONCERTO-3 studies. *Journal of Gastroenterology*. 2014; 49(5):941-953.
67. Kumada H, Hayashi N, Izumi N, et al. Simeprevir (TMC435) once daily with peginterferon-alpha-2b and ribavirin in patients with genotype 1 hepatitis C virus infection: The CONCERTO-4 study. *Hepatology Research: The Official Journal of the Japan Society of Hepatology*. 2014.
68. Kowdley KV, Lawitz E, Crespo I, et al. Sofosbuvir with pegylated interferon alfa-2a and ribavirin for treatment-naive patients with hepatitis C genotype-1 infection (ATOMIC): an open-label, randomised, multicentre phase 2 trial. *Lancet*. 2013; 381(9883):2100-2107.
69. Lawitz E, Lalezari JP, Hassanein T, et al. Sofosbuvir in combination with peginterferon alfa-2a and ribavirin for non-cirrhotic, treatment-naive patients with genotypes 1, 2, and 3 hepatitis C infection: a randomised, double-blind, phase 2 trial. *The Lancet Infectious Diseases*. 2013; 13(5):401-408.
70. Lawitz E, Mangia A, Wyles D, et al. Sofosbuvir for previously untreated chronic hepatitis C infection. *The New England Journal of Medicine*. 2013; 368(20):1878-1887.
71. Gane EJ, Stedman CA, Hyland RH, et al. Nucleotide polymerase inhibitor sofosbuvir plus ribavirin for hepatitis C. *The New England Journal of Medicine*. 2013; 368(1):34-44.
72. Lalezari J, Nelson DR, Hyland RH, et al. Once daily sofosbuvir plus ribavirin for 12 and 24 weeks in treatment-naïve patients with HCV infection: The QUANTUM study. *Journal of Hepatology*. 2013; 58(Suppl 1):S346.
73. Osinusi A, Meissner EG, Lee YJ, et al. Sofosbuvir and ribavirin for hepatitis C genotype 1 in patients with unfavorable treatment characteristics: a randomized clinical trial. *JAMA: The Journal of the American Medical Association*. 2013; 310(8):804-811.
74. Afdhal N, Reddy KR, Nelson DR, et al. Ledipasvir and sofosbuvir for previously treated HCV genotype 1 infection. *The New England Journal of Medicine*. 2014; 370(16):1483-1493.
75. Afdhal N, Zeuzem S, Kwo P, et al. Ledipasvir and sofosbuvir for untreated HCV genotype 1 infection. *The New England Journal of Medicine*. 2014; 370(20):1889-1898.

76. Gane EJ, Stedman CA, Hyland RH, et al. Efficacy of nucleotide polymerase inhibitor sofosbuvir plus the NS5A inhibitor ledipasvir or the NS5B non-nucleoside inhibitor GS-9669 against HCV genotype 1 infection. *Gastroenterology*. 2014; 146(3):736-743.e731.
77. Kowdley KV, Gordon SC, Reddy KR, et al. Ledipasvir and sofosbuvir for 8 or 12 weeks for chronic HCV without cirrhosis. *The New England Journal of Medicine*. 2014; 370(20):1879-1888.
78. Lawitz E, Poordad FF, Pang PS, et al. Sofosbuvir and ledipasvir fixed-dose combination with and without ribavirin in treatment-naïve and previously treated patients with genotype 1 hepatitis C virus infection (LONESTAR): an open-label, randomised, phase 2 trial. *Lancet*. 2014; 383(9916):515-523.
79. Osinusi A, Kohli A, Marti MM, et al. Re-treatment of Chronic Hepatitis C Virus Genotype 1 Infection After Relapse: An Open-Label Pilot Study. *Annals of Internal Medicine*. 2014; 161(9):634-638.
80. Gane E, Hyland RH, An D, et al. Sofosbuvir/ledipasvir fixed dose combination is safe and effective in difficult-to-treat populations including genotype-3 patients, decompensated genotype-1 patients, and genotype-1 patients with prior sofosbuvir treatment experience. 49th European Association for the Study of the Liver International Liver Congress (EASL 2014); April 9-13, 2014, 2014; London, England.
81. Kohli A, Sims Z, Marti MM, et al. Combination Oral Hepatitis C Antiviral Therapy for 6 or 12 weeks: Results from the SYNERGY Trial. Conference on Retroviruses and Opportunistic Infection; March 3-6, 2014, 2014; Boston, MA.
82. Chayama K, Takahashi S, Toyota J, et al. Dual therapy with the nonstructural protein 5A inhibitor, daclatasvir, and the nonstructural protein 3 protease inhibitor, asunaprevir, in hepatitis C virus genotype 1b-infected null responders. *Hepatology (Baltimore, Md.)*. 2012; 55(3):742-748.
83. Kumada H, Suzuki Y, Ikeda K, et al. Daclatasvir plus asunaprevir for chronic HCV genotype 1b infection. *Hepatology (Baltimore, Md.)*. 2014; 59(6):2083-2091.
84. Lok AS, Gardiner DF, Hezode C, et al. Randomized trial of daclatasvir and asunaprevir with or without PegIFN/RBV for hepatitis C virus genotype 1 null responders. *Journal of Hepatology*. 2014; 60(3):490-499.
85. Lok AS, Gardiner DF, Lawitz E, et al. Preliminary study of two antiviral agents for hepatitis C genotype 1. *The New England Journal of Medicine*. 2012; 366(3):216-224.
86. Manns M, Pol S, Jacobson IM, et al. All-oral daclatasvir plus asunaprevir for hepatitis C virus genotype 1b: a multinational, phase 3, multicohort study. *Lancet*. 2014.
87. Suzuki Y, Ikeda K, Suzuki F, et al. Dual oral therapy with daclatasvir and asunaprevir for patients with HCV genotype 1b infection and limited treatment options. *Journal of Hepatology*. 2013; 58(4):655-662.



88. Andreone P, Colombo MG, Enejosa JV, et al. ABT-450, ritonavir, ombitasvir, and dasabuvir achieves 97% and 100% sustained virologic response with or without ribavirin in treatment-experienced patients with HCV genotype 1b infection. *Gastroenterology*. 2014; 147(2):359-365.e351.
89. Feld JJ, Kowdley KV, Coakley E, et al. Treatment of HCV with ABT-450/r-ombitasvir and dasabuvir with ribavirin. *The New England Journal of Medicine*. 2014; 370(17):1594-1603.
90. Ferenci P, Bernstein D, Lalezari J, et al. ABT-450/r-ombitasvir and dasabuvir with or without ribavirin for HCV. *The New England Journal of Medicine*. 2014; 370(21):1983-1992.
91. Kowdley KV, Lawitz E, Poordad F, et al. Phase 2b trial of interferon-free therapy for hepatitis C virus genotype 1. *The New England Journal of Medicine*. 2014; 370(3):222-232.
92. Poordad F, Hezode C, Trinh R, et al. ABT-450/r-ombitasvir and dasabuvir with ribavirin for hepatitis C with cirrhosis. *The New England Journal of Medicine*. 2014; 370(21):1973-1982.
93. Zeuzem S, Jacobson IM, Baykal T, et al. Retreatment of HCV with ABT-450/r-ombitasvir and dasabuvir with ribavirin. *The New England Journal of Medicine*. 2014; 370(17):1604-1614.
94. Hezode C, Hirschfield GM, Ghesquiere W, et al. Daclatasvir plus peginterferon alfa and ribavirin for treatment-naïve chronic hepatitis C genotype 1 or 4 infection: a randomised study. *Gut*. 2014.
95. Lawitz E, Gaultier I, Poordad F, Cohen DE, Menon R. ABT-450/ritonavir (ABT-450/R) combined with pegylated interferon alpha-2a and ribavirin (SOC) after 3-day monotherapy in genotype 1 HCV-infected treatment-naïve subjects: 12-week interim efficacy and safety results. *Journal of Hepatology*. 2011; 54(Suppl. 1):S482.  
<http://onlinelibrary.wiley.com/doi/10.1002/hep.24444>  
[http://ac.els-cdn.com/S0168827811612221/1-s2.0-S0168827811612221-main.pdf?\\_tid=b0d8b288-3ead-11e4-b102-00000aabb0f26&acdnat=1410987824\\_f08bc26764857c2fffaa0f36dc39d2a2](http://ac.els-cdn.com/S0168827811612221/1-s2.0-S0168827811612221-main.pdf?_tid=b0d8b288-3ead-11e4-b102-00000aabb0f26&acdnat=1410987824_f08bc26764857c2fffaa0f36dc39d2a2)
96. Lenz O, de Bruijne J, Vijgen L, et al. Efficacy of re-treatment with TMC435 as combination therapy in hepatitis C virus-infected patients following TMC435 monotherapy. *Gastroenterology*. 2012; 143(5):1176-1178.e1171-1176.
97. Lawitz E, Poordad F, Kowdley KV, et al. A phase 2a trial of 12-week interferon-free therapy with two direct-acting antivirals (ABT-450/r, ABT-072) and ribavirin in IL28B C/C patients with chronic hepatitis C genotype 1. *Journal of Hepatology*. 2013; 59(1):18-23.
98. Lawitz E, Sullivan G, Rodriguez-Torres M, et al. Exploratory trial of ombitasvir and ABT-450/r with or without ribavirin for HCV genotype 1, 2, and 3 infection. *The Journal of Infection*. 2014.
99. Pol S, Ghalib RH, Rustgi VK, et al. Daclatasvir for previously untreated chronic hepatitis C genotype-1 infection: a randomised, parallel-group, double-blind, placebo-controlled, dose-finding, phase 2a trial. *The Lancet Infectious Diseases*. 2012; 12(9):671-677.

100. Poordad F, Lawitz E, Kowdley KV, et al. Exploratory study of oral combination antiviral therapy for hepatitis C. *The New England Journal of Medicine*. 2013; 368(1):45-53.
101. Rodriguez-Torres M, Lawitz E, Kowdley KV, et al. Sofosbuvir (GS-7977) plus peginterferon/ribavirin in treatment-naïve patients with HCV genotype 1: a randomized, 28-day, dose-ranging trial. *Journal of Hepatology*. 2013; 58(4):663-668.
102. Suzuki F, Toyota J, Ikeda K, et al. A randomized trial of daclatasvir with peginterferon alfa-2b and ribavirin for HCV genotype 1 infection. *Antiviral Therapy*. 2014.
103. Wyles DL, Rodriguez-Torres M, Lawitz E, et al. All-oral combination of ledipasvir, vedoprevir, tegobuvir, and ribavirin in treatment-naïve patients with genotype 1 HCV infection. *Hepatology (Baltimore, Md.)*. 2014; 60(1):56-64.
104. Dieterich D, Rockstroh JK, Orkin C, et al. Simeprevir (TMC435) with Peginterferon/Ribavirin in Patients Coinfected with HCV Genotype-1 and HIV-1: A Phase III Study. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*. 2014.
105. Osinusi A, Townsend K, Nelson A, et al. Use of sofosbuvir/ledipasvir fixed dose combination for treatment of HCV genotype-1 infection in patients coinfecting with HIV (interim results). 49th Annual Meeting of the European Association for the Study of the Liver (EASL); April 10-13, 2014, 2014; London, United Kingdom.
106. Sulkowski MS, Naggie S, Lalezari J, et al. Sofosbuvir and ribavirin for hepatitis C in patients with HIV coinfection. *JAMA: The Journal of the American Medical Association*. 2014; 312(4):353-361.
107. Charlton M, Gane E, Manns MP, et al. Sofosbuvir and Ribavirin for Treatment of Compensated Recurrent Hepatitis C Virus Infection After Liver Transplantation. *Gastroenterology*. 2014.
108. Curry MP, Forns X, Chung RT, et al. Sofosbuvir and Ribavirin Prevent Recurrence of HCV Infection after Liver Transplantation: An Open-Label Study. *Gastroenterology*. 2014.
109. Pellicelli AM, Montalbano M, Lionetti R, et al. Sofosbuvir plus daclatasvir for post-transplant recurrent hepatitis C: Potent antiviral activity but no clinical benefit if treatment is given late. *Digestive and Liver Disease: Official Journal of the Italian Society of Gastroenterology and the Italian Association for the Study of the Liver*. 2014; 46(10):923-927.
110. Ollendorf D, Pearson SD. ICER Evidence Rating Matrix: A User's Guide. 2013. <http://www.icer-review.org/wp-content/uploads/2013/04/Rating-Matrix-User-Guide-Exec-Summ-FINAL.pdf>.
111. Cardoso AC, Moucari R, Figueiredo-Mendes C, et al. Impact of peginterferon and ribavirin therapy on hepatocellular carcinoma: incidence and survival in hepatitis C patients with advanced fibrosis. *Journal of Hepatology*. 2010; 52(5):652-657.



112. George SL, Bacon BR, Brunt EM, Mihindukulasuriya KL, Hoffmann J, Di Bisceglie AM. Clinical, virologic, histologic, and biochemical outcomes after successful HCV therapy: a 5-year follow-up of 150 patients. *Hepatology (Baltimore, Md.)*. 2009; 49(3):729-738.
113. Giannini EG, Basso M, Savarino V, Picciotto A. Sustained virological response to pegylated interferon and ribavirin is maintained during long-term follow-up of chronic hepatitis C patients. *Alimentary Pharmacology & Therapeutics*. 2010; 31(4):502-508.
114. Trapero-Marugan M, Mendoza J, Chaparro M, et al. Long-term outcome of chronic hepatitis C patients with sustained virological response to peginterferon plus ribavirin. *World Journal of Gastroenterology: WJG*. 2011; 17(4):493-498.
115. Koh C, Heller T, Haynes-Williams V, et al. Long-term outcome of chronic hepatitis C after sustained virological response to interferon-based therapy. *Alimentary Pharmacology & Therapeutics*. 2013; 37(9):887-894.
116. Marcellin P, Boyer N, Gervais A, et al. Long-term histologic improvement and loss of detectable intrahepatic HCV RNA in patients with chronic hepatitis C and sustained response to interferon-alpha therapy. *Annals of Internal Medicine*. 1997; 127(10):875-881.
117. Arora S, O'Brien C, Zeuzem S, et al. Treatment of chronic hepatitis C patients with persistently normal alanine aminotransferase levels with the combination of peginterferon alpha-2a (40 kDa) plus ribavirin: impact on health-related quality of life. *Journal of Gastroenterology and Hepatology*. 2006; 21(2):406-412.
118. Spiegel BM, Younossi ZM, Hays RD, Revicki D, Robbins S, Kanwal F. Impact of hepatitis C on health related quality of life: a systematic review and quantitative assessment. *Hepatology (Baltimore, Md.)*. 2005; 41(4):790-800.
119. Morgan RL, Baack B, Smith BD, Yartel A, Pitasi M, Falck-Ytter Y. Eradication of hepatitis C virus infection and the development of hepatocellular carcinoma: a meta-analysis of observational studies. *Annals of Internal Medicine*. 2013; 158(5 Pt 1):329-337.
120. Morgan TR, Ghany MG, Kim HY, et al. Outcome of sustained virological responders with histologically advanced chronic hepatitis C. *Hepatology (Baltimore, Md.)*. 2010; 52(3):833-844.
121. van der Meer AJ, Veldt BJ, Feld JJ, et al. Association between sustained virological response and all-cause mortality among patients with chronic hepatitis C and advanced hepatic fibrosis. *JAMA: The Journal of the American Medical Association*. 2012; 308(24):2584-2593.
122. Veldt BJ, Heathcote EJ, Wedemeyer H, et al. Sustained virologic response and clinical outcomes in patients with chronic hepatitis C and advanced fibrosis. *Annals of Internal Medicine*. 2007; 147(10):677-684.
123. McCombs J, Matsuda T, Tonnu-Mihara I, et al. The Risk of Long-term Morbidity and Mortality in Patients With Chronic Hepatitis C: Results From an Analysis of Data From a Department of Veterans Affairs Clinical Registry. *JAMA Internal Medicine*. 2013.

124. Ferreira Sda C, Carneiro Mde V, Souza FF, et al. Long-term follow-up of patients with chronic hepatitis C with sustained virologic response to interferon. *The Brazilian Journal of Infectious Diseases: An Official Publication of the Brazilian Society of Infectious Diseases*. 2010; 14(4):330-334.
125. Sonnenberg FA, Beck JR. Markov models in medical decision making: a practical guide. *Medical Decision Making: An International Journal of the Society for Medical Decision Making*. 1993; 13(4):322-338.
126. Gold M. *Cost-Effectiveness in Health and Medicine*. New York: Oxford University Press; 1996.
127. Bai Y, Zhao Y, Wang G, Wang H, Liu K, Zhao W. Cost-effectiveness of a hypertension control intervention in three community health centers in China. *Journal of Primary Care & Community Health*. Jul 1 2013; 4(3):195-201.
128. Alba AC, Alba LF, Delgado DH, Rao V, Ross HJ, Goeree R. Cost-effectiveness of ventricular assist device therapy as a bridge to transplantation compared with nonbridged cardiac recipients. *Circulation*. Jun 18 2013; 127(24):2424-2435.
129. Ubel PA, Hirth RA, Chernew ME, Fendrick AM. What is the price of life and why doesn't it increase at the rate of inflation? *Archives of Internal Medicine*. Jul 28 2003; 163(14):1637-1641.
130. Hirth RA, Chernew ME, Miller E, Fendrick AM, Weissert WG. Willingness to pay for a quality-adjusted life year: in search of a standard. *Medical Decision Making: An International Journal of the Society for Medical Decision Making*. Jul-Sep 2000; 20(3):332-342.
131. CHOosing Interventions that are Cost Effective (WHO-CHOICE). 2014.  
[http://www.who.int/choice/cost-effectiveness/inputs/health\\_service/en/](http://www.who.int/choice/cost-effectiveness/inputs/health_service/en/)
132. Hutubessy R, Chisholm D, Edejer TT. Generalized cost-effectiveness analysis for national-level priority-setting in the health sector. *Cost Effectiveness and Resource Allocation: C/E*. Dec 19 2003; 1(1):8.
133. Ditah I, Ditah F, Devaki P, et al. The Changing Epidemiology of Hepatitis C Virus Infection in the United States: National Health and Nutrition Examination Survey 2001 through 2010. *Journal of Hepatology*. Nov 27 2013.
134. Bedossa P, Poynard T. An algorithm for the grading of activity in chronic hepatitis C. The METAVIR Cooperative Study Group. *Hepatology*. Aug 1996; 24(2):289-293.
135. Seeff LB. Natural history of chronic hepatitis C. *Hepatology*. Nov 2002; 36(5 Suppl 1):S35-46.
136. Gilead Sciences Inc. SOVALDI (sofosbuvir) Prescribing Information. Gilead Sciences Inc, Foster City, CA 94404.
137. Thein H-H, Yi Q, Dore GJ, Krahn MD. Estimation of stage-specific fibrosis progression rates in chronic hepatitis C virus infection: a meta-analysis and meta-regression. *Hepatology*. 2008; 48(2):418-431.

138. Liu S, Cipriano LE, Holodniy M, Owens DK, Goldhaber-Fiebert JD. New protease inhibitors for the treatment of chronic hepatitis C: a cost-effectiveness analysis. *Ann Intern Med.* Feb 21 2012; 156(4):279-290.
139. Yousuf M, Nakano Y, Tanaka E, Sodeyama T, Kiyosawa K. Persistence of viremia in patients with type-C chronic hepatitis during long-term follow-up. *Scandinavian Journal of Gastroenterology.* 1992; 27(9):812-816.
140. Hagan LM, Yang Z, Ehteshami M, Schinazi RF. All-oral, interferon-free treatment for chronic hepatitis C: cost-effectiveness analyses. *Journal of Viral Hepatitis.* Dec 2013; 20(12):847-857.
141. Coffin PO, Scott JD, Golden MR, Sullivan SD. Cost-effectiveness and population outcomes of general population screening for hepatitis C. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America.* 2012; 54(9):1259-1271.
142. Razavi H, Elkhoury AC, Elbasha E, et al. Chronic hepatitis C virus (HCV) disease burden and cost in the United States. *Hepatology.* Jun 2013; 57(6):2164-2170.
143. El-Kamary SS, Jhaveri R, Shardell MD. All-cause, liver-related, and non-liver-related mortality among HCV-infected individuals in the general US population. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America.* Jul 15 2011; 53(2):150-157.
144. Veldt BJ, Saracco G, Boyer N, et al. Long term clinical outcome of chronic hepatitis C patients with sustained virological response to interferon monotherapy. *Gut.* Oct 2004; 53(10):1504-1508.
145. Arias E. United States life tables, 2009. *National vital statistics reports: from the Centers for Disease Control and Prevention, National Center for Health Statistics, National Vital Statistics System.* Jan 6 2014; 62(7):1-63.
146. Micromedex. Red Book Online. 2014. Accessed July 11th, 2014.
147. Centers for Medicare & Medicaid Services. CMS - Healthcare Common Procedure Coding System (HCPCS) codes. 2014. Accessed July 21, 2014.
148. Rein DB, Wittenborn JS. *The Cost-Effectiveness of Birth Cohort and Universal Hepatitis C Antibody Screening in U.S. Primary Care Settings - Technical Report.* Research Triangle Park, NC: RTI International; 2011.
149. Carlson JJ, Kowdley KV, Sullivan SD, Ramsey SD, Veenstra DL. An evaluation of the potential cost-effectiveness of non-invasive testing strategies in the diagnosis of significant liver fibrosis. *Journal of Gastroenterology and Hepatology.* May 2009; 24(5):786-791.
150. Backx M, Lewszuk A, White JR, et al. The cost of treatment failure: resource use and costs incurred by hepatitis C virus genotype 1-infected patients who do or do not achieve sustained virological response to therapy. *Journal of Viral Hepatitis.* Mar 2014; 21(3):208-215.

151. Manos MM, Darbinian J, Rubin J, et al. The effect of hepatitis C treatment response on medical costs: a longitudinal analysis in an integrated care setting. *Journal of Managed Care Pharmacy: JMCP*. Jul-Aug 2013; 19(6):438-447.
152. McAdam-Marx C, McGarry LJ, Hane CA, Biskupiak J, Deniz B, Brixner DI. All-cause and incremental per patient per year cost associated with chronic hepatitis C virus and associated liver complications in the United States: a managed care perspective. *Journal of managed care pharmacy: JMCP*. Sep 2011; 17(7):531-546.
153. D'Ambrosio R, Aghemo A, Rumi MG, et al. A morphometric and immunohistochemical study to assess the benefit of a sustained virological response in hepatitis C virus patients with cirrhosis. *Hepatology*. Aug 2012; 56(2):532-543.
154. Mallet V, Gilgenkrantz H, Serpaggi J, et al. Brief communication: the relationship of regression of cirrhosis to outcome in chronic hepatitis C. *Annals of Internal Medicine*. Sep 16 2008; 149(6):399-403.
155. Maylin S, Martinot-Peignoux M, Moucari R, et al. Eradication of hepatitis C virus in patients successfully treated for chronic hepatitis C. *Gastroenterology*. Sep 2008; 135(3):821-829.
156. Pol S, Carnot F, Nalpas B, et al. Reversibility of hepatitis C virus-related cirrhosis. *Hum Pathol*. 2004; 35(1):107-112.
157. Serpaggi J, Carnot F, Nalpas B, et al. Direct and indirect evidence for the reversibility of cirrhosis. *Human Pathology*. 12// 2006; 37(12):1519-1526.
158. Drummond, MF, Sculpher, MJ, Torrance, GW, et al. *Methods for the Economic Evaluation of Health Care Programmes*. Third Edition ed: Oxford University Press; 2005.
159. Salomon JA, Weinstein MC, Hammitt JK, Goldie SJ. Cost-effectiveness of treatment for chronic hepatitis C infection in an evolving patient population. *JAMA*. Jul 9 2003; 290(2):228-237.
160. Thein HH, Krahn M, Kaldor JM, Dore GJ. Estimation of utilities for chronic hepatitis C from SF-36 scores. *Am J Gastroenterol*. Mar 2005; 100(3):643-651.
161. Gao X, Stephens JM, Carter JA, Haider S, Rustgi VK. Impact of adverse events on costs and quality of life in protease inhibitor-based combination therapy for hepatitis C. *Expert Review Of Pharmacoeconomics & Outcomes research*. Jun 2012; 12(3):335-343.
162. McHutchison JG, Lawitz EJ, Shiffman ML, Muir AJ, et al. Peginterferon alfa-2b or alfa-2a with ribavirin for treatment of hepatitis C infection. *N Eng J Med*. Aug 2009; 361(6):580-93.
163. Lawitz E, Mangia A, Wyles D, et al. Sofosbuvir for previously untreated chronic hepatitis C infection. *N Engl J Med*. 2013; 368(20):1878-1887.
164. Lawitz E, Sulkowski MS, Ghalib R, et al. Simeprevir plus sofosbuvir, with or without ribavirin, to treat chronic infection with hepatitis C virus genotype 1 in non-responders to pegylated interferon and ribavirin and treatment-naïve patients: the COSMOS randomised study. *Lancet*. Jul 26 2014.

165. McCombs J, Matsuda T, Tonnu-Mihara I, et al. The risk of long-term morbidity and mortality in patients with chronic hepatitis C: results from an analysis of data from a Department of Veterans Affairs Clinical Registry. *JAMA Internal Medicine*. Feb 1 2014; 174(2):204-212.
166. Thein HH, Yi Q, Dore GJ, Krahn MD. Estimation of stage-specific fibrosis progression rates in chronic hepatitis C virus infection: a meta-analysis and meta-regression. *Hepatology*. Aug 2008; 48(2):418-431
167. Hagan LM, Sulkowski MS, Schinazi RF. Cost analysis of sofosbuvir/ribavirin versus sofosbuvir/simeprevir for genotype 1 hepatitis C virus in interferon-ineligible/intolerant individuals. *Hepatology* 2014; 60:37-45.
168. Saab S, Gordon SC, Park H, et al. Cost-effectiveness of sofosbuvir plus peginterferon/ribavirin in the treatment of chronic hepatitis C virus genotype 1 infection. *Aliment Pharmacol Ther* 2014; 40:657-75.
169. Petta S, Cabibbo G, Enea M, et al. Cost-effectiveness of sofosbuvir-based triple therapy for untreated patients with genotype 1 chronic hepatitis C. *Hepatology* 2014; 59:1692-1705.
170. Liu S, Watcha D, Holodny M, Goldhaber-Fiebert J. Sofosbuvir-based treatment regimens for chronic, genotype 1 hepatitis C virus infection in U.S. incarcerated populations. *Ann Intern Med* 2014; 161:546-53.
171. Leleu H, Blachier M, Rosa I. Cost-effectiveness of sofosbuvir in the treatment of patients with hepatitis C. *J Viral Hepat* 2014 Sep 15. doi: 10.1111/jvh.12311. [Epub ahead of print].
172. San Miguel R, Gimeno-Ballester V, Blazquez A, Mar J. Cost-effectiveness analysis of sofosbuvir-based regimens for chronic hepatitis C. *Gut* 2014 Oct 13. pii: gutjnl-2014-307772. doi: 10.1136/gutjnl-2014-307772. [Epub ahead of print].
173. Hagan LM, Yang Z, Ehteshami M, Schinazi RF. All-oral, interferon-free treatment for chronic hepatitis C: cost-effectiveness analysis. *J Viral Hepat* 2013; 20:847-57.
174. Younossi ZM, Singer ME, Mir HM, Henry L, Hunt S. Impact of interferon free regimens on clinical and cost outcomes for chronic hepatitis C genotype 1 patients. *J Hepatol* 2014 Mar; 60:530-7.
175. Deuffic-Burban S, Schwarzingier M, Obach D, et al. Should we await IFN-free regimens to treat HCV genotype 1 treatment-naïve patients? A cost-effectiveness analysis (ANRS 95141). *J Hepatol* 2014 Jul; 61:7-14.
176. McCombs J, Matsuda T, Tonnu-Mihara I, et al. The risk of long-term morbidity and mortality in patients with chronic hepatitis C: Results from an analysis of data from a Department of Veterans Affairs clinical registry. *JAMA Int Med* 2014 Feb 1; 174:204-12.
177. Express Scripts, Inc. State Governments May Spend \$55 Billion on Hepatitis C Medications. Accessed at: [http://lab.express-scripts.com/insights/specialty-medications/state-governments-may-spend-\\$55-billion-on-hepatitis-c-medications](http://lab.express-scripts.com/insights/specialty-medications/state-governments-may-spend-$55-billion-on-hepatitis-c-medications), November 2014.

178. Nainan OV, Alter MJ, Kruszon-Moran D, et al. Hepatitis C virus genotypes and viral concentrations in participants of a general population survey in the United States. *Gastroenterology* Aug 2006; 131:478-484.
179. California Department of Health Care Services. 6-month ACA Expansion Capitation Rate Summary - with SB78. January 1, 2014 – June 30, 2014. Accessed at: [http://www.dhcs.ca.gov/dataandstats/reports/Documents/MMCD\\_Fin\\_Rpts/Medicaid\\_ACA\\_Expansion\\_Rates.pdf](http://www.dhcs.ca.gov/dataandstats/reports/Documents/MMCD_Fin_Rpts/Medicaid_ACA_Expansion_Rates.pdf), November 2014.
180. Tice JA, Ollendorf D, Pearson SD. The Comparative Clinical Effectiveness and Value of Simeprevir and Sofosbuvir in the Treatment of Chronic Hepatitis C Infection: A Technology Assessment. March 2014. <http://ctaf.org/reports/treatments-hepatitis-c>
181. Pearlman BL, Traub N. Sustained virologic response to antiviral therapy for chronic hepatitis C virus infection: a cure and so much more. *Clin Infect Dis*. 2011; 52(7):889-900.
182. Bruno S, Zuin M, Crosignani A, Rossi S, Zadra F, Roffi L, et al. Predicting mortality risk in patients with compensated HCV-induced cirrhosis: a long-term prospective study. *Am J Gastroenterol*. 2009; 104(5):1147-58.

# **APPENDICES**

## Appendix A: Coverage Policies

Appendix Table A1: Coverage Policies for LDV/SOF

	Medi-Cal	<a href="#">Aetna</a>	<a href="#">Anthem</a>	CVS/Caremark	<a href="#">Health Net</a>	Humana	<a href="#">UHC</a>
METAVIR or equivalent score	N/A	Covered with documentation of HCV diagnosis, genotype, and subtype	Covered if $\geq$ F3	N/A	Covered if $\geq$ F2	N/A	Covered if $\geq$ F3
Patients with severe renal impairment	N/A	--	Not covered	N/A	--	N/A	--
Extrahepatic manifestations	N/A	--	Covered	N/A	--	N/A	Covered
Decompensated liver disease	N/A	Not covered	Not covered	N/A	Not covered	N/A	--
Post-liver transplant	N/A	Not covered	Covered	N/A	--	N/A	Covered
Hepatocellular carcinoma	N/A	Not covered	--	N/A	--	N/A	--
Eligible if treatment experienced?	N/A	Not eligible if previous SOF failure	Not eligible if previous LDV or SOF failure	N/A	Not eligible if previous SOF failure	N/A	Yes, including any protease inhibitor or SOF failure
Treatment continuation based on reduced HCV RNA levels	N/A	Yes	--	N/A	--	N/A	--
Treatment restrictions related to abuse of illicit drugs and/or alcohol	N/A	--	Yes	N/A	--	N/A	Yes
Require specialist to prescribe or consult	N/A	--	--	N/A	Yes	N/A	Yes
Other criteria	N/A	For patients meeting clinical criteria, use of LDV/SOF is required unless patient is contraindicated or intolerant to any of its ingredients	Not to be used in combination with other NS5B polymerase or NS5A inhibitors	N/A	Non-FDA-approved indications are covered only with sufficient documentation in published literature	N/A	--



Maximum duration authorized	N/A	8 weeks: tx-naïve w/o cirrhosis, viral load <6M; 12 weeks: tx-naïve w/o cirrhosis and viral load ≥6M OR tx-naïve w/ cirrhosis OR tx-experienced w/o cirrhosis; 24 weeks: tx-experienced w/ cirrhosis	8 weeks: tx-naïve w/o cirrhosis, viral load <6M; 12 weeks: tx-naïve w/o cirrhosis and viral load ≥6M OR tx-naïve w/ cirrhosis OR tx-experienced w/o cirrhosis; 24 weeks: tx-experienced w/ cirrhosis	N/A	8 weeks: tx-naïve w/o cirrhosis, viral load <6M; 12 weeks: tx-naïve w/o cirrhosis and viral load ≥6M OR tx-naïve w/ cirrhosis OR tx-experienced w/o cirrhosis; 24 weeks: tx-experienced w/ cirrhosis	N/A	8 weeks: tx-naïve w/o cirrhosis, viral load <6M; 12 weeks: tx-naïve w/o cirrhosis and viral load ≥6M OR post-liver transplant OR tx-naïve w/ cirrhosis OR tx-experienced w/o cirrhosis; 24 weeks: tx-experienced w/ cirrhosis
Published/revised/effective date	N/A	10/31/2014	10/15/2014	N/A	10/28/2014	N/A	10/15/2014

Abbreviations: HCC = hepatocellular carcinoma; P = pegylated interferon; R = ribavirin; tx = treatment; -- = not specified in coverage policy; N/A = no online coverage policy available

Note: The information in this table is extracted from publicly available documents as of November 3, 2014 and is not intended to be a definitive source on coverage policies, as these are being updated regularly and contain details that cannot reasonably be reflected in this summary table. Interested parties should obtain current, specific coverage policy information from individual payers.

**Appendix Table A2: Coverage Policies for Sofosbuvir + PR**

	<a href="#">Medi-Cal</a>	<a href="#">Aetna</a>	<a href="#">Anthem</a>	<a href="#">CVS/Caremark</a>	<a href="#">Health Net</a>	<a href="#">Humana</a>	<a href="#">UHC</a>
METAVIR or equivalent score	Covered if $\geq$ F3 or if F0-F2 with severe extrahepatic manifestations	Covered with documentation of HCV diagnosis, genotype, and subtype	Covered if $\geq$ F3	-	Covered if $\geq$ F2	-	Covered if $\geq$ F3
Patients with severe renal impairment	Not covered	--	Not covered	Not covered	--	--	--
Extrahepatic manifestations	Covered	--	Covered	--	--	--	Covered
Decompensated liver disease	Not covered	Not covered	Not covered	Not covered	Not covered	Not covered	Not covered
Post-liver transplant	Must meet DHCS investigational services criteria	Covered	SOF covered but treatment regimen not specified	Not covered	Covered	Covered for patients with compensated liver disease	Covered
Hepatocellular carcinoma	Not covered	Not covered	Not covered	Not covered	Not covered	--	Not covered
Eligible if treatment experienced?	Yes	Yes in most cases, see policy for details	Not eligible if previous PR + protease or polymerase inhibitor failure	--	--	--	Not eligible if previous SOF failure
Treatment continuation based on reduced HCV RNA levels	Recommended	Yes	--	--	--	--	--
Treatment restrictions related to abuse of illicit drugs and/or alcohol	Yes	--	Yes	--	--	--	Yes
Require specialist to prescribe or consult	Recommended	--	--	--	Yes	--	Yes

Other criteria	All non-FDA approved indications must meet DHCS investigational services criteria	Intolerance/contraindication to or nonfulfillment of criteria for LDV/SOF required	--	--	Failure/contraindication to LDV/SOF required; non-FDA-approved use must be supported by published literature	Investigational/experimental SOF regimens must be supported by published literature or CMS compendia	--
Maximum duration authorized	12 weeks	24 weeks for post-liver transplant; otherwise 12 weeks	12 weeks	12 weeks	12 weeks	12 weeks	12 weeks
Published/revised/effective date	6/30/2014	10/31/2014	10/17/2014	--	10/28/2014	3/6/2014	9/1/2014

Abbreviations: HCC = hepatocellular carcinoma; P = pegylated interferon; R = ribavirin; tx = treatment; -- = not specified in coverage policy; N/A = no online coverage policy available

Note: The information in this table is extracted from publicly available documents as of November 3, 2014 and is not intended to be a definitive source on coverage policies, as these are being updated regularly and contain details that cannot reasonably be reflected in this summary table. Interested parties should obtain current, specific coverage policy information from individual payers.

**Appendix Table A3: Coverage Policies for Sofosbuvir + R**

	<a href="#">Medi-Cal</a>	<a href="#">Aetna</a>	<a href="#">Anthem</a>	<a href="#">CVS/Caremark</a>	<a href="#">Health Net</a>	<a href="#">Humana</a>	<a href="#">UHC</a>
METAVIR or equivalent score	Covered if $\geq$ F3 or if F0-F2 with severe extrahepatic manifestations	Covered with documentation of HCV diagnosis, genotype, and subtype	Covered if $\geq$ F3	--	Covered if $\geq$ F2	--	Covered if $\geq$ F3
Patients with severe renal impairment	Not covered	--	Not covered	Not covered	--	--	--
Extrahepatic manifestations	Covered	--	Covered	--	--	--	Covered
Decompensated liver disease	Covered, patient must be referred to specialist	Covered	Covered if decompensation is reason for interferon-ineligibility	Covered if decompensation is reason for interferon-ineligibility	Not covered	Covered	Covered
Post-liver transplant	Must meet DHCS investigational services criteria	Covered	SOF covered but treatment regimen not specified	Not covered	Covered	Covered for patients with decompensated liver disease	Covered
Hepatocellular carcinoma	Covered	Covered if awaiting liver transplant	Covered if awaiting liver transplant	Covered if awaiting liver transplant	Covered if awaiting liver transplant	--	Covered if patient is on waiting list for liver transplant and being managed in a liver transplant center
Eligible if treatment experienced?	Yes in most cases, see other criteria for details	--	Not eligible if previous PR + protease or polymerase inhibitor failure	Not eligible if previous SOF failure	--	Yes	Not eligible if previous SOF failure
Treatment continuation based on reduced HCV RNA levels	Recommended	Yes	--	--	--	--	--
Treatment restrictions related to abuse of illicit drugs and/or alcohol	Yes	--	Yes	--	--	--	Yes

Require specialist to prescribe or consult	Recommended	--	--	--	Yes	--	Yes
Other criteria	Must be interferon-ineligible; must meet DCHS investigational services criteria for non-FDA approved indications including treatment-experienced, advanced fibrosis/compensated cirrhosis, interferon-eligible	Intolerance/contraindication to or nonfulfillment of criteria for LDV/SOF required; must be interferon-ineligible	Must be interferon-ineligible	Must be interferon-ineligible	Failure/contraindication to LDV/SOF required; must be interferon-ineligible; non-FDA approved use must be supported by published literature	Investigational/experimental SOF regimens must be supported by published literature or CMS compendia	Must be interferon-ineligible; documented contraindication to SMV required unless patient has HCC or decompensated liver disease
Maximum duration authorized	24-48 weeks or until liver transplant for HCC; otherwise 24 weeks	48 weeks or until liver transplantation for HCC or decompensated cirrhosis; otherwise 24 weeks	48 weeks or until liver transplantation for HCC; otherwise 24 weeks	48 weeks or until liver transplantation for HCC; otherwise 24 weeks	48 weeks or until liver transplantation for HCC; otherwise 24 weeks	48 weeks or until liver transplantation for decompensated cirrhosis or for decompensated liver disease post-liver transplant; otherwise 24 weeks	48 weeks for HCC or decompensated liver disease; otherwise 24 weeks
Published/revised/effective date	6/30/2014	10/31/2014	10/17/2014	--	10/28/2014	3/6/2014	9/1/2014

Abbreviations: HCC = hepatocellular carcinoma; P = pegylated interferon; R = ribavirin; tx = treatment; -- = not specified in coverage policy; N/A = no online coverage policy available

Note: The information in this table is extracted from publicly available documents as of November 3, 2014 and is not intended to be a definitive source on coverage policies, as these are being updated regularly and contain details that cannot reasonably be reflected in this summary table. Interested parties should obtain current, specific coverage policy information from individual payers.

**Appendix Table A4: Coverage Policies for Simeprevir/Sofosbuvir ± R**

	<a href="#">Medi-Cal</a>	<a href="#">Aetna</a>	<a href="#">Anthem</a>	<a href="#">CVS/Caremark</a>	Health Net	<a href="#">Humana</a>	<a href="#">UHC</a>
METAVIR or equivalent score	Covered if ≥F3 or if F0-F2 with severe extrahepatic manifestations and interferon-ineligible	Covered with documentation of HCV diagnosis, genotype, and subtype	Covered if ≥F3	Covered if ≥F3	N/A	--	Covered if ≥F3
Patients with severe renal impairment	Not covered	--	--	--	N/A	--	--
Extrahepatic manifestations	Covered	--	Covered	--	N/A	--	Covered
Decompensated liver disease	Covered, patient must be referred to specialist	Not covered	Covered if decompensation is reason for interferon-ineligibility	Covered if decompensation is reason for interferon-ineligibility	N/A	Covered if decompensation is reason for interferon-ineligibility	Not covered
Post-liver transplant	Must meet DHCS investigational services criteria	Covered	SOF covered but treatment regimen not specified	Covered only if treatment-naïve post-transplant	N/A	Covered for patients with compensated liver disease	Covered
Hepatocellular carcinoma	Not covered	Not covered	Not covered	Covered if HCC is reason for interferon-ineligibility	N/A	--	--
Eligible if treatment experienced?	Yes	Yes if previous PR failure	Not eligible if previous PR + protease or polymerase inhibitor failure	Yes if previous failure of PR therapy without a protease inhibitor	N/A	Yes	Not eligible if previous SOF failure, unless discontinuation due to PR intolerance
Treatment continuation based on reduced HCV RNA levels	Recommended	Yes	--	Yes	N/A	--	--
Treatment restrictions related to abuse of illicit drugs and/or alcohol	Yes	--	Yes	--	N/A	--	Yes

Require specialist to prescribe or consult	Recommended	--	--	--	N/A	--	Yes
Other criteria	All non-FDA-approved indications must meet DHCS investigational services criteria	Intolerance/contraindication to or nonfulfillment of criteria for LDV/SOF required; must be interferon-ineligible or post-liver transplant	Patient must be interferon-ineligible OR have had a previous partial or nonresponse to PR therapy	Patient must be treatment naïve and interferon-ineligible OR have had a previous PR failure	N/A	Patient must be interferon-ineligible or treatment experienced; investigational/experimental SOF regimens must be supported by published literature or CMS compendia	Must be interferon-ineligible
Maximum duration authorized	12 weeks	12-24 weeks for post-liver transplant; otherwise 12 weeks	12 weeks	24 weeks for post-liver transplant; otherwise 12 weeks	N/A	12-24 weeks for post-liver transplant; otherwise 12 weeks	12 weeks
Published/revised/effective date	6/30/2014	10/31/2014	10/17/2014	--	N/A	3/6/2014	9/1/2014

Abbreviations: HCC = hepatocellular carcinoma; P = pegylated interferon; R = ribavirin; tx = treatment; -- = not specified in coverage policy; N/A = no online coverage policy available

Note: The information in this table is extracted from publicly available documents as of November 3, 2014 and is not intended to be a definitive source on coverage policies, as these are being updated regularly and contain details that cannot reasonably be reflected in this summary table. Interested parties should obtain current, specific coverage policy information from individual payers.

**Appendix Table A5: Coverage Policies for Simeprevir + PR**

	<a href="#">Medi-Cal</a>	<a href="#">Aetna</a>	<a href="#">Anthem</a>	<a href="#">CVS/Caremark</a>	<a href="#">Health Net</a>	<a href="#">Humana</a>	<a href="#">UHC</a>
METAVIR or equivalent score	Covered if $\geq$ F3 or if F0-F2 with severe extrahepatic manifestations	Covered with documentation of HCV diagnosis, genotype, and subtype	Covered if $\geq$ F3	--	Covered if $\geq$ F2	--	--
Patients with severe renal impairment	--	--	--	--	--	--	--
Extrahepatic manifestations	Covered	--	--	--	--	--	--
Decompensated liver disease	Not covered	Not covered	Not covered	Not covered	Not covered	Not covered	Not covered
Genotype 1a NS3 Q80k polymorphism	Not covered	Not covered	Not covered	Not covered	Not covered	Not covered	Not recommended
Post-liver transplant	--	Not covered	--	Not covered	--	--	--
Hepatocellular carcinoma	Not covered	--	--	--	--	--	--
Eligible if treatment experienced?	Not eligible if previous protease inhibitor failure	Not eligible if previous protease inhibitor failure	Not eligible if previous PR + protease or polymerase inhibitor failure	Not eligible if previous PR + protease inhibitor failure	Not eligible if previous protease inhibitor failure	Not eligible if previous protease inhibitor failure	Not eligible if previous protease inhibitor failure
Treatment continuation based on reduced HCV RNA levels	Yes	Yes	--	Yes	Yes	Yes	--
Treatment restrictions related to abuse of illicit drugs and/or alcohol	Yes	--	Yes	--	--	--	--
Require specialist to prescribe or consult	--	--	--	--	--	--	--



Other criteria	Prior treatment failure with any protease inhibitor precludes use of SMV; all non-FDA-approved indications must meet DHCS investigational services criteria	Intolerance/contraindication to or nonfulfillment of criteria for LDV/SOF required	Not for use in combination with other protease inhibitors	--	Failure/contraindication to LDV/SOF required; non-FDA approved use must be supported by published literature	Investigational/experimental SMV regimens must be supported by published literature or CMS compendia	--
Maximum duration authorized	SMV up to 12 weeks, R up to 48 weeks	SMV up to 12 weeks, R up to 48 weeks	--	SMV up to 12 weeks, R up to 48 weeks	SMV up to 12 weeks, R up to 48 weeks	SMV up to 12 weeks, R up to 48 weeks	SMV up to 12 weeks, R up to 48 weeks
Published/revised/effective date	6/30/2014	10/31/2014	7/2/2014	--	10/16/2014	10/2/2014	9/1/2014

Abbreviations: HCC = hepatocellular carcinoma; P = pegylated interferon; R = ribavirin; tx = treatment; -- = not specified in coverage policy; N/A = no online coverage policy available

Note: The information in this table is extracted from publicly available documents as of November 3, 2014 and is not intended to be a definitive source on coverage policies, as these are being updated regularly and contain details that cannot reasonably be reflected in this summary table. Interested parties should obtain current, specific coverage policy information from individual payers.

## Appendix B: Search Strategies

### PubMed (NLM), run date 9/10/14

sofosbuvir OR simeprevir OR daclatasvir OR ombitasvir OR abt-450\* AND English[la] NOT (review[pt] OR editorial[pt] OR news[pt]) AND (clinical trial[pt] OR clinical trials as topic[mh] OR random\* OR study OR trial OR trials)  
157 refs

### Cochrane Library (Wiley), run date 9/10/14

sofosbuvir OR simeprevir OR daclatasvir OR ombitasvir OR "abt-450" OR "abt-450r"

All Results (58)

Cochrane Reviews (0) All Review Protocol Other Reviews (1) Trials (47) Methods Studies (0)  
Technology Assessments (9) Economic Evaluations (1) Cochrane Groups (0)

Cochrane Central Register of Controlled Trials (Central) Issue 8 of 12, August 2014

### Embase (Elsevier), run date 9/10/14

sofosbuvir or simeprevir or daclatasvir or ombitasvir or 'abt-450' or 'abt-450r' and [english]/lim and ('clinical trial'/de or 'clinical trial (topic)'/de or 'controlled study'/de or 'double blind procedure'/de or 'major clinical study'/de or 'multicenter study'/de or 'multicenter study (topic)'/de or 'phase 2 clinical trial'/de or 'phase 2 clinical trial (topic)'/de or 'phase 3 clinical trial'/de or 'phase 3 clinical trial (topic)'/de or 'randomized controlled trial'/de or 'randomized controlled trial (topic)'/de or random\* or study or trial or trials) not ('conference abstract'/it OR 'conference review'/it or 'editorial'/it or 'review'/it or 'short survey'/it)  
404 refs

## Appendix C: Supplemental Tables from Chapter 6

### Simeprevir + PR

**Appendix Table C1. Clinical Trials of Simeprevir + PR in Patients Infected with HCV Genotype 1**

Study	Publication	Study drugs	Control	Treatment Naïve	Prevalence of Cirrhosis (%)
<i>Phase 2</i>					
PILLAR	Fried 2013 <sup>41</sup>	SMV12/24 + PR48	PR48	Yes	0
ASPIRE	Zeuzem 2014 <sup>63</sup>	SMV12/24/48 + PR48	PR48	No	18
<i>Phase 3</i>					
QUEST 1	Jacobson 2014 <sup>61</sup>	SMV12 + PR24/48	PR48	Yes	12
QUEST 2	Manns 2014 <sup>62</sup>	SMV12 + PR24/48	PR48	Yes	9
PROMISE	Forns 2013 <sup>60</sup>	SMV12 + PR24/48	PR48	No	15
<i>Japan</i>					
CONCERTO-1	Hayashi 2014b <sup>64</sup>	SMV12 + PR24/48	PR48	Yes	0
CONCERTO-2	Izumi 2014 <sup>66</sup>	SMV12 + PR24/48 or SMV24 + PR24/48	PR48	No	0
CONCERTO-3	Izumi 2014 <sup>66</sup>	SMV12 + PR24/48	None	No	0
CONCERTO-4	Kumada 2014 <sup>67</sup>	SMV12 + PR24/48	None	Both	0
DRAGON	Hayashi 2014a <sup>65</sup>	SMV12 + PR24	PR48	Yes	0

**Appendix Table C2. Summary of the Outcomes of Simeprevir + PR in Patients Infected with HCV Genotype 1**

*FDA approved or probable treatment dose/duration only*

Study	Treatment Naïve	Cirrhosis	Treatment	N	SVR (%)	DR (%)
QUEST 1	Yes	No	SMV12 + PR24/48	233	82.4	8.2
QUEST 2	Yes	No	SMV12 + PR24/48	240	82.5	4.6
QUEST 1	Yes	Yes	SMV12 + PR24/48	31	58.1	6.5
QUEST 2	Yes	Yes	SMV12 + PR24/48	17	64.7	5.9
ASPIRE	No	No	SMV12 + PR48	53	66.0	7.5
PROMISE	No	No	SMV12 + PR24/48	221	80.1	0.9
ASPIRE	No	Yes	SMV12 + PR48	13	69.2	7.7
PROMISE	No	Yes	SMV12 + PR24/48	39	74.4	20.5

## **Sofosbuvir + PR**

**Appendix Table C3. Clinical Trials of Sofosbuvir + PR in Patients Infected with HCV Genotype 1**

Study	Publication	Study drugs	Control	Treatment Naïve	Prevalence of Cirrhosis (%)
<i>Phase 2</i>					
PROTON	Lawitz 2013a <sup>69</sup>	SOF12 + PR24/48	PR24/48	Yes	0
ATOMIC	Kowdley 2013 <sup>68</sup>	SOF12 + PR12 or SOF24 + PR24	None	Yes	0
<i>Phase 3</i>					
NEUTRINO	Lawitz 2013b <sup>70</sup>	SOF12 + PR12	None	Yes	17

**Appendix Table C4. Summary of the Outcomes of Sofosbuvir + PR in Patients Infected with HCV Genotype 1**

*FDA approved or probable treatment dose/duration only*

Study	Treatment Naïve	Cirrhosis	Treatment	N	SVR (%)	DR (%)
PROTON	Yes	No	SOF12 + PR24/48	47	89.4	17.0
ATOMIC	Yes	No	SOF12 + PR12	52	88.5	9.6
NEUTRINO	Yes	No	SOF12 + PR12	249	92.8	9.6
NEUTRINO	Yes	Yes	SOF12 + PR12	43	81.4	11.6

## **Sofosbuvir + R**

**Appendix Table C5. Clinical Trials of Sofosbuvir + R in Patients Infected with HCV Genotype 1**

Study	Publication	Study drugs	Control	Treatment Naïve	Prevalence of Cirrhosis (%)
<i>Phase 2</i>					
QUANTUM	Abstract <sup>72</sup>	SOF24 + R24	None	Yes	6
NIH SPARE	Osinusi 2013 <sup>73</sup>	SOF24 + R24	None	Yes	23
ELECTRON	Gane 2013 <sup>71</sup>	SOF12 + R12	None	Both	0

**Appendix Table C6. Summary of the Outcomes of sofosbuvir + R in Patients Infected with HCV Genotype 1**

*FDA approved or probable treatment dose/duration only*

Study	Treatment Naïve	Cirrhosis	Treatment	N	SVR (%)	DR (%)
NIH SPARE	Yes	No	SOF24 + R24	10	90.0	10.0
NIH SPARE	Yes	No	SOF24 + R24	19	73.7	10.5
QUANTUM	Yes	No	SOF24 + R24	19	47.4	5.3
NIH SPARE	Yes	Yes	SOF24 + R24	6	50.0	0.0

## Simeprevir/Sofosbuvir

**Appendix Table C7. Clinical Trials of Simeprevir/Sofosbuvir in Patients Infected with HCV Genotype 1**

Study	Publication	Study drugs	Control	Treatment Naïve	Prevalence of Cirrhosis (%)
<i>Phase 2</i>					
COSMOS	Lawitz 2014 <sup>58</sup>	SMV/SOF12 ± R12 or SMV/SOF24 ± R24	None	Both	25

**Appendix Table C8. Summary of the Outcomes of Simeprevir/Sofosbuvir in Patients Infected with HCV Genotype 1**

*FDA approved or probable treatment dose/duration only*

Study	Treatment Naïve	Cirrhosis	Treatment	N	SVR (%)	DR (%)
COSMOS	Yes	No	SMV/SOF12	4	100	0
COSMOS	Yes	Yes	SMV/SOF24	6	100	16.7
COSMOS	No	No	SMV/SOF12	14	92.9	0
COSMOS	No	No	SMV/SOF12	3	100	0
COSMOS	No	Yes	SMV/SOF24	4	100	0

## Ledipasvir/Sofosbuvir

**Appendix Table C9. Clinical Trials of Ledipasvir/Sofosbuvir in Patients Infected with HCV Genotype 1**

Study	Publication	Study drugs	Control	Treatment Naïve	Prevalence of Cirrhosis (%)
<i>Phase 2</i>					
LONESTAR	Lawitz 2014 <sup>78</sup>	LDV/SOF8 or 12 ± R	None	Both	22
ELECTRON	Gane 2014 <sup>76</sup>	LDV/SOF6 or 12 ± R ± GS-9669	None	Both	17
ELECTRON 2	Abstract <sup>80</sup>	LDV/SOF12	None	No	0
NIH SPARE 2	Osinusi 2014 <sup>79</sup>	LDV/SOF12	None	No	50
SYNERGY	Abstract <sup>81</sup>	LDV/SOF12 ± GS-9451	None	Yes	30
<i>Phase 3</i>					
ION-1	Afdhal 2014 <sup>75</sup>	LDV/SOF12 ± R12 or LDV/SOF24 ± R24	None	Yes	16
ION-2	Afdhal 2014 <sup>74</sup>	LDV/SOF12 ± R12 or LDV/SOF24 ± R24	None	No	20
ION-3	Kowdley 2014 <sup>77</sup>	LDV/SOF8 ± R8 or LDV/SOF12 ± R12	None	No	0

**Appendix Table C10. Summary of the Outcomes of Ledipasvir/Sofosbuvir in Patients Infected with HCV Genotype 1**

*FDA approved or probable treatment dose/duration only*

Study	Treatment Naïve	Cirrhosis	Treatment	N	SVR (%)	DR (%)
LONESTAR	Yes	No	LDV/SOF8	20	95.0	0.0
ION-3	Yes	No	LDV/SOF8	215	94.0	0.9
LONESTAR	Yes	No	LDV/SOF12	19	94.7	5.3
SYNERGY	Yes	No	LDV/SOF12	17	100	0.0
ION-1	Yes	No	LDV/SOF12	180	99.4	0.6
ION-3	Yes	No	LDV/SOF12	216	95.4	4.2
SYNERGY	Yes	Yes	LDV/SOF12	3	100	0.0
ELECTRON-2	Yes	Yes	LDV/SOF12	20	65.0	0.0
ION-1	Yes	Yes	LDV/SOF12	34	94.1	2.9
LONESTAR	No	No	LDV/SOF12	8	100	0.0
ION-2	No	No	LDV/SOF12	87	95.4	0.0
ION-2	No	Yes	LDV/SOF24	22	100	0

## Daclatasvir/Sofosbuvir

**Appendix Table C11. Clinical Trials of Daclatasvir/Sofosbuvir in Patients Infected with HCV Genotype 1**

Study	Publication	Study drugs	Control	Treatment Naïve	Prevalence of Cirrhosis (%)
<i>Phase 2</i>					
AI444040	Sulkowski 2014 <sup>59</sup>	DCV/SOF12 ± R12 or DCV/SOF24 ± R24	None	Both	16

**Appendix Table C12. Summary of the Outcomes of Daclatasvir/Sofosbuvir in Patients Infected with HCV Genotype 1**

*FDA approved or probable treatment dose/duration only*

Study	Treatment Naïve	Cirrhosis	Treatment	N	SVR (%)	DR (%)
AI444040	Yes	No	DCV/SOF12	35	100	0.0
AI444040	Yes	No	DCV/SOF24	25	100	4.0
AI444040	Yes	Yes	DCV/SOF12	6	100	0.0
AI444040	Yes	Yes	DCV/SOF24	4	100	0.0
AI444040	No	No	DCV/SOF24	18	92.9	0.0
AI444040	No	Yes	DCV/SOF24	3	100	0.0



## Daclatasvir/Asunaprevir

**Appendix Table C13. Clinical Trials of Daclatasvir/Asunaprevir in Patients Infected with HCV Genotype 1**

Study	Publication	Study drugs	Control	Treatment Naïve	Prevalence of Cirrhosis (%)
<i>Phase 2</i>					
NCT01012895	Lok 2012 <sup>85</sup>	DCV/ASV24 ± PR24	None	No	0
	Lok 2014 <sup>84</sup>	DCV/ASV24 ± PR24	None	No	0
<i>Phase 3</i>					
HALLMARK-DUAL GT1b only	Manns 2014 <sup>86</sup>	DCV/ASV24	Placebo	Both	30
<i>Japan</i>					
GT1b only	Chayama 2012 <sup>82</sup>	DCV/ASV24	None	No	0
GT1b only	Suzuki 2013 <sup>87</sup>	DCV/ASV24	None	Both	0
GT1b only	Kumada 2014 <sup>83</sup>	DCV/ASV24	None	Both	10

The dosing used in the US Phase 3 clinical trial HALLMARK-DUAL<sup>86</sup> (daclatasvir 60 mg once daily plus asunaprevir 100 mg twice daily) was only used in one of the other clinical trials.<sup>83</sup> It would likely have been the recommended dosing schedule for the combination if BMS had not withdrawn its application from the FDA. In early studies, asunaprevir was dosed at 600 mg twice daily and reduced to 200 mg twice daily due to elevations in liver enzymes.<sup>82,85,87</sup> It is worth noting that the combination of daclatasvir and asunaprevir was not as effective in HCV genotype 1a and the later, larger Phase 3 studies are limited to genotype 1b. This is the primary genotype in Japan.

**Appendix Table C14. Summary of the Outcomes of Daclatasvir/Asunaprevir in Patients Infected with HCV Genotype 1**

*FDA approved or probable treatment dose/duration only*

Study	Treatment Naïve	Cirrhosis	Treatment	N	SVR (%)	DR (%)
Kumada 2014	Yes	No	DCV/ASV24	124	87.1	10.4
HALLMARK-DUAL	Yes	No	DCV/ASV24	171	89.5	7.4
Kumada 2014	Yes	Yes	DCV/ASV24	11	90.9	10.4
HALLMARK-DUAL	Yes	Yes	DCV/ASV24	32	90.6	7.4
Kumada 2014	No	No	DCV/ASV24	76	78.9	16.1
HALLMARK-DUAL	No	No	DCV/ASV24	142	79.6	13.7
Kumada 2014	No	Yes	DCV/ASV24	11	90.9	16.1
HALLMARK-DUAL	No	Yes	DCV/ASV24	63	87.3	13.7

## Paritaprevir, Ritonavir, Ombitasvir, and Dasabuvir (3D) ± Ribavirin

**Appendix Table C15. Clinical Trials of 3D ± R in Patients Infected with HCV Genotype 1**

Study	Publication	Study drugs	Control	Treatment Naïve	Prevalence of Cirrhosis (%)
<i>Phase 2</i>					
AVIATOR	Kowdley 2014 <sup>91</sup>	3D12 ± R12 or 3D24 ± R24  14 groups	None	Both	0
<i>Phase 3</i>					
PEARL-II GT1b only	Andreone 2014 <sup>88</sup>	3D12 ± R12	None	No	0
PEARL-III GT1b only	Ferenci 2014 <sup>90</sup>	3D12 ± R12	None	Yes	0
PEARL-IV GT1a only	Ferenci 2014 <sup>90</sup>	3D12 ± R12	None	Yes	0
SAPPHIRE-I	Feld 2014 <sup>89</sup>	3D12 + R12	Placebo	Yes	0
SAPPHIRE-II	Zeuzem 2014 <sup>93</sup>	3D12 + R12	Placebo	No	0
TURQUOISE-II	Poordad 2014 <sup>92</sup>	3D12 + R12 or 3D24 + R24	None	No	100

**Appendix Table C16. Summary of the Outcomes of 3D + R in Patients Infected with HCV Genotype 1**

*FDA approved or probable treatment dose/duration only*

Study	Treatment Naïve	Cirrhosis	Treatment	N	SVR (%)	DR (%)
AVIATOR	Yes	No	3D24 + R24	40	90.0	7.5
AVIATOR	Yes	No	3D12 + R12	40	95.0	5.0
PEARL-III	Yes	No	3D12 + R12	210	99.5	0.5
PEARL-IV	Yes	No	3D12 + R12	100	97.0	0.0
SAPPHIRE-I	Yes	No	3D12 + R12	473	96.2	1.9
TURQUOISE-II	Yes	Yes	3D12 + R12	86	94.2	2.3
TURQUOISE-II	Yes	Yes	3D24 + R24	74	94.6	5.4
AVIATOR	No	No	3D12 + R12	22	95.5	0.0
AVIATOR	No	No	3D24 + R24	20	100	0.0
PEARL-II	No	No	3D12 + R12	95	95.8	4.2
SAPPHIRE-II	No	No	3D12 + R12	297	96.3	1.7
TURQUOISE-II	No	Yes	3D12 + R12	122	90.2	1.6
TURQUOISE-II	No	Yes	3D24 + R24	98	96.9	5.1

## **HIV Co-infection**

**Appendix Table C17. Clinical Trials of the Treatment of HCV in HIV Co-infected Patients**

Study	Publication	Study drugs	Control	Treatment Naïve	Prevalence of Cirrhosis (%)
C212	Dieterich 2014 <sup>104</sup>	SMV12 + PR24/48	None	Both	13
PHOTON-1	Sulkowski 2014 <sup>106</sup>	SOF24 + R24	None	Yes for GT1	4
ERADICATE	Abstract <sup>105</sup>	LDV/SOF12	None	Yes	0

**Appendix Table C18. Summary of the Outcomes in Patients Co-infected with HCV Genotype 1 and HIV**

*FDA approved or probable treatment dose/duration only*

Study	Treatment Naïve	Cirrhosis	Treatment	N	SVR (%)	DR (%)
C212	Yes	~6%	SMV12 + PR24/48	53	79.2	17.0
C212	No	~11%	SMV12 + PR24/48	53	67.9	34.0
C212	Both	Yes	SMV12 + PR48	9	77.8	-
PHOTON-1	Yes	No	SOF24 + R24	109	77.1	11.4
PHOTON-1	Yes	Yes	SOF24 + R24	5	60.0	11.4
ERADICATE	Yes	No	LDV/SOF12	10	100	0.0

## Pre- or post-liver transplant

**Appendix Table C19. Clinical Trials of the Treatment of HCV Pre- or Post-liver Transplant**

Study	Publication	Study drugs	Control	Treatment Naïve	Prevalence of Cirrhosis (%)
P7977-2025 Pre-transplant	Curry 2014 <sup>108</sup>	SOF48 + R48	None	Both	100% with HCC
Post-transplant	Charlton 2014 <sup>107</sup>	SOF24 + R24	None	Both	40
Post-transplant	Pellicelli 2014 <sup>109</sup>	DCV/SOF24 ± R24	None	NR	75

**Appendix Table C20. Summary of the Outcomes in Patients with HCV Genotype 1 Pre- and Post-liver Transplant**

*FDA approved or probable treatment dose/duration only*

Study	Transplant	Cirrhosis	Treatment	N	SVR (%)	DR (%)
Curry 2014 <sup>108</sup>	Pre	100% with HCC	SOF48 + R48	45	17/31 54.8% 12 weeks after transplant	24.6
Charlton 2014 <sup>107</sup>	Post	40	SOF24 + R24	53	67.9	34.0
Pellicelli 2014 <sup>109</sup>	Post	75	DCV/SOF24 ± R24	12	5/5 SVR4	3 unrelated deaths

## Appendix D: Supplementary Tables for Chapter 7

**Table D1: METAVIR Score for Classification of Liver Damage Due to HCV and Distribution of Fibrosis Stages in CHC Population**

Stage of Fibrosis	Histological definition	Distribution of fibrosis	Reference
F0	No fibrosis	0.17 (0.14-0.19)	134
F1	Portal fibrosis without septa	0.35 (0.26-0.39)	134
F2	Portal fibrosis with rare septa	0.22 (0.18-0.24)	134
F3	Numerous septa without cirrhosis	0.14 (0.12-0.15)	134
F4 (CC)	Compensated Cirrhosis	0.12 (0.11-0.13)	134

CHC – Chronic Hepatitis C; F0-F4 – METAVIR fibrosis score; CC – Compensated Cirrhosis

**Table D2: Chronic Hepatitis C Annual Post-SVR Transition Probabilities, and Regression Proportions**

Source State	Target State	Base case	Lower limit	Upper	Reference
CHC Progression Post-SVR					
F0	F1	0.010023	0.005012	0.015035	Calculated*
F1	F2	0.007282	0.003641	0.010923	Calculated*
F2	F3	0.01028	0.00514	0.01542	Calculated*
F3	F4	0.009937	0.004969	0.014906	Calculated*
	Decompensated Cirrhosis	0.001028	0.0005	0.0015	140
	Hepatocellular Carcinoma	0.004753	0.001	0.007	140
F4	Decompensated Cirrhosis	0.003342	0.002	0.005	140
	Hepatocellular Carcinoma	0.012449	0.006	0.019	140
Decompensated Cirrhosis	Hepatocellular Carcinoma	0.010	0.008	0.017	126
	Liver Transplant	0.012	0.007	0.016	141
	Death	0.09	0.07	0.15	126
Fibrosis Regression Post-SVR (Proportions)					
F1	F0	0.35	0.17	0.52	142-145
F2	F0	0.12	0.06	0.18	142-145
	F1	0.58	0.29	0.87	142-145
F3	F1	0.24	0.12	0.36	142-145
	F2	0.46	0.23	0.69	142-145
F4	F1	0.09	0.05	0.14	142-149
	F2	0.14	0.07	0.21	142-149
	F3	0.22	0.11	0.33	142-146,148,150

\* – calculated post-SVR F0 to F4 transition probabilities (using non-SVR probabilities from meta-analysis by Thein et al.) based on a 91% reduction observed in progression from F3 to decompensated cirrhosis post-SVR

**Table D3: SVR and Discontinuation Rates of Sofosbuvir-based Treatments**

Therapy	Subgroup	Treatment Duration	SVR (95% CI)	DR (95% CI)
<b>SOF + PR</b>	Naïve, no cirrhosis	12 weeks	.920 (.888-.948)	.103 (.072-.139)
	Naïve, + cirrhosis	12 weeks	.814 (.666-.916)	.116 (.039-.251)
	Experienced, no cirrhosis	12 weeks	.780 (0.390-1.00) †	.103 (.072-.139) ‡
	Experienced, + cirrhosis	12 weeks	.710 (.570-0.830)	.116 (.039-.251) ‡
<b>SOF + R</b>	Naïve, no cirrhosis	24 weeks	.750 (.675-.819)	.078 (.036-.131)
	Naïve, + cirrhosis	24 weeks	.545 (.227-.484)	.000 (.000-.013)
<b>SMV/SOF</b>	Naïve, no cirrhosis	12 weeks	1.00 (.398-1.00)	.000 (.000-.602)
	Naïve, + cirrhosis	12 weeks	.667 (.094-.992)	.333 (.008-.906)
	Experienced, no cirrhosis	12 weeks	.970 (.781-1.00)	.000 (.000-.083)
	Experienced, + cirrhosis	12 weeks	1.00 (.398-1.00)	.000 (.000-.602)
<b>LDV/SOF*</b>	Naïve, no cirrhosis	8 weeks	.948 (.913-.976)	.002 (.000-.018)
	Naïve, no cirrhosis	12 weeks	.985 (.968-.997)	.013 (.002-.029)
	Naïve, + cirrhosis	12 weeks	.892 (.778-.974)	.000 (.000-.043)
	Experienced, no cirrhosis	12 weeks	.977 (.924-1.00)	.000 (.000-.004)
	Experienced, + cirrhosis	24 weeks	1.00 (.846-1.00)	.000 (.000-.154)

\* – For base-case 67% of patients were allocated to receive LDV/SOF 8, while the remaining received 12 weeks of LDV/SOF therapy. This value was varied in probabilistic sensitivity analysis using a range of (30% to 90%).

† – CI selected by authors (lower limit 50% of base-case, upper limit 100%)

‡ – Due to lack of data, discontinuation rates modeled to be the same as treatment-naïve group.

**Table D4: SVR and Discontinuation Rates for PR (48 weeks)**

<b>Treatment Characteristics</b>	<b>Base case (%)</b>	<b>Lower limit*</b>	<b>Upper limit*</b>	<b>Reference</b>
<i>Treatment-naïve</i>				
<i>SVR – Overall</i>	54.6	27	82	153, 154
Discontinuation rate	24.2	12	36	154, 155
EVR12	79.9	40	100†	153, 154
SVR followed by EVR12	68.3	34	85†	153, 154
<i>Treatment-experienced</i>				
<i>SVR – Overall</i>	16.5	8	25	156, 157
Discontinuation rate	64.6	32	97	157
<b>SVR by fibrosis</b>	<b>Base case (%)</b>	<b>Lower limit*</b>	<b>Upper limit*</b>	<b>Reference</b>
<i>Prior Relapse (0.53)</i>				
<i>Overall for Prior Relapse</i>	22.1	11	33	156, 157
F0-F1	35	18	53	156, 157
F2	27.8	14	42	156, 157
F3	13.3	7	20	156, 157
F4	6.7	3	10	156, 157
<i>Partial Response (0.19)</i>				
<i>Overall for Partial Response</i>	18.2	9	27	156, 157
F0-F1	0	0	10‡	156, 157
F2	42.9	21	64	156, 157
F3	0	0	10‡	156, 157
F4	20	10	30	156, 157
<i>Null Response (0.28)</i>				
<i>Overall for Null Response</i>	5.4	3	8	156, 157
F0-F1	0.0	0	10‡	156, 157
F2	7.7	4	12	156, 157
F3	0	0	10‡	156, 157
F4	10	5	15	156, 157

\* – Lower and upper bounds are 50% to 150% of base-case, unless otherwise noted.

† – Lower and upper bounds are 50% to 125% of base-case.

‡ – Upper limit selected by authors.

EVR = Early Virologic Response

**Table D5: Frequency, by Week, of Follow-up/Testing/Management of Each Treatment Modality**

Test and Office Visit	8-week therapy	12-week therapies					24-week therapies				48-week therapy
		SOF + PR	LDV/SOF	SMV/SOF	3D	DCV/SOF	SOF + R	LDV/SOF	3D	DCV/SOF	PR
<b>Anti-HCV (antibody) test</b>	0 (#1)†,‡	0 (#1)					0 (#1)				0 (#1)
<b>Genotype assay</b>	0 (#1) ‡	0 (#1)					0 (#1)				0 (#1)
<b>Fibrosis assessment</b>	0 (#1) ‡	0 (#1)					0 (#1)				0 (#1)
<b>HCV RNA quantification</b>	0, 4, 8, 12 (#4) ‡	0, 4, 12, 24 (#4)					0, 4, 24, 36 (#4)				0, 4, 12, 24, 48, 60 (#6) §
<b>CBC w/Differential</b>	0, 4, 8, 12 (#4) ‡	0, 4, 8, 12, 24 (#5)					0, 4, 8, 12, 16, 24, 36 (#7)				0, 4, 8, 12, 16, 24, 48, 60 (#8)
<b>Hepatic function panel</b>	0, 4, 8, 12 (#4) ‡	0, 4, 8, 12, 24 (#5)					0, 4, 8, 12, 16, 24, 36 (#7)				0, 4, 8, 12, 16, 24, 48, 60 (#8)
<b>Office visit (outpatient)</b>	0, 4, 8, 12 (#4) ‡	0, 4, 8, 12, 24 (#5)					0, 4, 8, 12, 16, 24, 36 (#7)				0, 4, 8, 12, 16, 24, 48, 60 (#8)

# – indicates the quantity of tests or office visits over the course of treatment.

\* – Treatment-naïve.

† – Week (0, 2, 4, etc.) at which the test or office visit takes place.

‡ – Per AASLD guidelines and an additional test at 12-weeks after end-of-treatment.<sup>159</sup>

§ – Increased number of tests based on response-guided therapy criteria for PR therapy.<sup>160</sup>



**Table D6: Utility Loss with CHC Treatment**


Treatment Modality	Annualized utility loss	Base case (during treatment)	Lower limit*	Upper limit*	Reference
<i>Utility penalties during treatment</i>					
PR (48 weeks)	-0.1931	-0.1782	-0.2896	0	Calculated
SOF + PR (12-weeks)	-0.1657	-0.0382	-0.2486	0	Calculated
SOF + R (24 weeks)	-0.0852	-0.0393	-0.1279	0	Calculated
SMV/SOF (12 weeks)	-0.0873	-0.0202	-0.1310	0	Calculated
LDV/SOF (8 weeks) †	-0.0754	-0.0116	-0.1130	0	Calculated
LDV/SOF (12 weeks)†	-0.0754	-0.0174	-0.1130	0	Calculated
LDV/SOF (24 weeks)†	-0.0754	-0.0348	-0.1130	0	Calculated

\* – Lower Limit is 50% more than the annualized base-case. Upper Limit is no utility loss.

† – Annualized disutility represents an average of disutility across the various treatment durations.

## Appendix E: Explanation of Disease Progression and Markov Model Details

Figure E1: Hepatitis C Natural History Markov Representation



**Figure E1 description:** Patients enter the Markov model either when they receive no treatment, after unsuccessful therapy, or treatment discontinuation, in stages F0 through F4. The black arrows indicate annual progression of liver damage. The one time “no-progression” proportion from F0 fibrosis state is


removed from the progression cascade after that proportion of patients accrue the cost of treatment under the "treat all" strategy.

### **Post-SVR progression and regression model**

To account for progression of liver disease and liver regeneration following SVR, this model allows patients to attain a worse or better health status after HCV eradication.<sup>181</sup> A graphical representation of post-SVR HCV history and health states is available in Figure E2. Following SVR, there exists the possibility of progression from F3 and F4 states to more advanced liver complications.<sup>4,182</sup> Therefore, patients will cycle through a set of post-SVR Markov states that have different transition probabilities than those of the natural history Markov states. The annual post-SVR progression probabilities are shown in Appendix Table D2 and are derived from published literature.<sup>4,141</sup>

Additionally, it is possible for the liver damage caused by HCV to be reversed, at least partially, in a subset of the patients following successful therapy.<sup>4,153-157</sup> The data for regression are determined from the literature as a proportion of patients achieving regression post-SVR.<sup>4,153-157</sup> Therefore, the model assumes that immediately after SVR, a certain percent of patients from F1 to F4 states regress to a lower fibrotic state as indicated by the proportions listed in Appendix Table D2.

**Figure E2: Hepatitis C Post-SVR Markov Representation Showing Progression and Regression of CHC Following Successful Treatment**



**Figure E2 description:** Patients enter the Markov model after successful therapy in stages F0 through F4. Blue arrows indicate proportional regression from source state to a lower fibrosis state. The regression data covers a wide time range, between 1 and 10 years post regression. In this model, the regression transition occurs immediately after successful treatment. Red arrows indicate annual progression of liver damage after achieving SVR.

Figure E3: Simplified Tree Structure of the HCV Cost-effectiveness Model




Figure E3 depicts a simplified tree of the model for illustrative purposes. This structure shows only five of 26 Markov states representing 15 health states. See Appendix Figures E1 and E2 for details of the Markov model. The Markov model is the same for all treatment policies. The policy analysis starts at the node marked with an “M.” At the “M” node, a policy to treat all immediately or to wait until patients progress to F3 or F4 stages is selected. The terminal nodes (red diamonds) indicate the transition to other Markov states depending on the outcome of the cycle.

## Appendix F: Scenario Analyses

**Table F1: Scenario 1: Increased Discontinuation Rates**

**Scenario Analysis** 1. PR treatment experienced DR increased by 1.5  
**Discontinuation rates** 2. All others DR doubled; including all treatment-naïve.  
 3. For treatments with base-case of 0, lowest non-zero value of DR from another treatment selected (0.013)

Incremental comparison of regimens							Vs. PR			Vs. No Treatment			Treat All vs. Treat at F3, F4		
Strategy	Cost	Incr Cost	Eff	Incr Eff	ICER	Comment	Cost	Eff	ICER	Cost	Eff	ICER	Cost	Eff	ICER
<b>Tx naïve, treat all</b>							<b>Tx naïve, treat all</b>						<b>Tx naïve, treat all</b>		
No Treatment	\$ 45,313	\$ -	11.82	0.00	-	undominated	\$ (19,562)	-0.970	\$ 20,160				\$ 13,315	0.22	\$ 60,028
PR (48 weeks)	\$ 64,875	\$ 19,562	12.79	0.97	\$ 20,160	ext. dominated				\$ 19,562	0.97	\$ 20,160	\$ 25,665	0.72	\$ 35,639
LDV/SOF (8/12 weeks)	\$ 90,947	\$ 45,634	14.72	2.90	\$ 15,736	undominated	\$ 26,072	1.930	\$ 13,511	\$ 45,634	2.90	\$ 15,736	\$ 35,554	0.59	\$ 60,573
SOF + PR (12 weeks)	\$ 106,417	\$ 15,470	14.19	-0.53	\$ (29,139)	abs. dominated	\$ 41,542	1.399	\$ 29,700	\$ 61,104	2.37	\$ 25,793	\$ 27,889	0.73	\$ 38,065
LDV/SOF (12 weeks)	\$ 108,436	\$ 17,489	14.77	0.04	\$ 411,659	undominated	\$ 43,561	1.972	\$ 22,089	\$ 63,123	2.94	\$ 21,453	\$ 62,852	0.73	\$ 85,861
SMV/SOF (12 weeks)	\$ 161,849	\$ 53,412	14.50	-0.27	\$ (199,644)	abs. dominated	\$ 96,974	1.705	\$ 56,891	\$ 116,536	2.67	\$ 43,566	\$ 68,744	0.52	\$ 132,383
SOF + R (24 weeks)	\$ 182,165	\$ 73,729	13.82	-0.95	\$ (77,799)	abs. dominated	\$ 117,290	1.024	\$ 114,494	\$ 136,852	1.99	\$ 68,606			
<b>Tx naïve, treat at F3, F4</b>							<b>Tx naïve, treat at F3, F4</b>								
No Treatment	\$ 45,313	\$ -	11.82	0.00	\$ -	undominated	\$ (6,247)	-0.749	\$ 8,346						
PR (48 weeks)	\$ 51,560	\$ 6,247	12.57	0.75	\$ 8,346	undominated				\$ 6,247	0.75	\$ 8,346			
LDV/SOF (8/12 weeks)	\$ 65,282	\$ 13,721	14.00	1.43	\$ 9,587	undominated	\$ 13,721	1.431	\$ 9,587	\$ 19,969	2.18	\$ 9,161			
SOF + PR (12 weeks)	\$ 70,863	\$ 5,581	13.60	-0.40	\$ (14,034)	abs. dominated	\$ 19,302	1.034	\$ 18,676	\$ 25,550	1.78	\$ 14,337			
LDV/SOF (12 weeks)	\$ 80,547	\$ 15,265	14.03	0.03	\$ 509,551	undominated	\$ 28,986	1.461	\$ 19,837	\$ 35,234	2.21	\$ 15,945			
SMV/SOF (12 weeks)	\$ 98,997	\$ 18,450	13.77	-0.27	\$ (69,134)	abs. dominated	\$ 47,437	1.194	\$ 39,717	\$ 53,684	1.94	\$ 27,631			
SOF + R (24 weeks)	\$ 113,421	\$ 32,875	13.30	-0.73	\$ (44,771)	abs. dominated	\$ 61,861	0.727	\$ 85,097	\$ 68,108	1.48	\$ 46,160			
<b>Tx exp, treat all</b>							<b>Tx exp, treat all</b>						<b>Tx exp, treat all</b>		
No Treatment	\$ 45,313	\$ -	11.82	0.00	\$ -	undominated	\$ (24,755)	0.144	\$ (172,066)				\$ 11,699	-0.06	\$ (185,847)
PR (48 weeks)	\$ 70,069	\$ 24,755	11.68	-0.14	\$ (172,066)	abs. dominated				\$ 24,755	-0.14	\$ (172,066)	\$ 35,434	0.57	\$ 62,258
SOF + PR (12 weeks)	\$ 110,196	\$ 64,883	13.83	2.01	\$ 32,273	ext. dominated	\$ 40,128	2.154	\$ 18,627	\$ 64,883	2.01	\$ 32,273	\$ 38,785	0.74	\$ 52,713
LDV/SOF (12/24 weeks)	\$ 119,079	\$ 73,766	14.76	2.93	\$ 25,147	undominated	\$ 49,010	3.077	\$ 15,927	\$ 73,766	2.93	\$ 25,147	\$ 63,199	0.68	\$ 93,022
SMV/SOF (12 weeks)	\$ 164,649	\$ 45,570	14.62	-0.13	\$ (344,119)	abs. dominated	\$ 94,580	2.945	\$ 32,117	\$ 119,336	2.80	\$ 42,605			
<b>Tx exp, treat at F3, F4</b>							<b>Tx exp, treat at F3, F4</b>								
No Treatment	\$ 45,313	\$ -	11.82	0.00	\$ -	undominated	\$ (13,057)	0.081	\$ (161,345)						
PR (48 weeks)	\$ 58,370	\$ 13,057	11.74	-0.08	\$ (161,345)	abs. dominated				\$ 13,057	-0.08	\$ (161,345)			
SOF + PR (12 weeks)	\$ 74,762	\$ 29,449	13.26	1.44	\$ 20,432	ext. dominated	\$ 16,392	1.522	\$ 10,769	\$ 29,449	1.44	\$ 20,432			
LDV/SOF (12/24 weeks)	\$ 80,294	\$ 5,532	14.02	0.76	\$ 7,314	undominated	\$ 21,924	2.279	\$ 9,622	\$ 34,981	2.20	\$ 15,918			
SMV/SOF (12 weeks)	\$ 101,450	\$ 21,156	13.94	-0.08	\$ (278,207)	abs. dominated	\$ 43,080	2.203	\$ 19,559	\$ 56,137	2.12	\$ 26,460			

DR — Discontinuation Rate; Incr Net Cost — Incremental Net Cost; Eff — Effectiveness; Incr Eff — Incremental Effectiveness

**Table F2: Scenario 2: Increased Portion of Cohort at Initial Stage F4**

**Scenario Analysis** 1. Scenario Analysis Distribution: F0, F1, F2, F3, F4 = 0.15, 0.33, 0.20, 0.12, 0.20 = 1.0  
**Increased F4 Prevalence**

Incremental comparison of regimens							Vs. PR			Vs. No Treatment			Treat All vs. Treat at F3, F4		
Strategy	Cost	Incr Cost	Eff	Incr Eff	ICER	Comment	Cost	Eff	ICER	Cost	Eff	ICER	Cost	Eff	ICER
<b>Tx naïve, treat all</b>							<b>Tx naïve, treat all</b>						<b>Tx naïve, treat all</b>		
No Treatment	\$ 50,227	\$ -	11.46	0.00	\$ -	undominated	-\$15,582	-1.559	\$ 9,993				\$12,938	0.340	\$ 38,048
PR (48 weeks)	\$ 65,809	\$ 15,582	13.02	1.56	\$ 9,993	undominated				\$ 15,582	1.559	\$ 9,993	\$23,544	0.668	\$ 35,259
LDV/SOF (8/12 weeks)	\$ 94,994	\$ 29,185	14.44	1.42	\$ 20,486	undominated	\$29,185	1.425	\$ 20,486	\$ 44,767	2.984	\$ 15,003	\$34,150	0.613	\$ 55,676
SOF + PR (12 weeks)	\$ 110,430	\$ 15,436	14.20	-0.24	\$ (64,235)	abs. dominated	\$44,621	1.184	\$ 37,676	\$ 60,203	2.744	\$ 21,943	\$25,574	0.684	\$ 37,362
LDV/SOF (12 weeks)	\$ 111,026	\$ 16,032	14.50	0.06	\$ 285,937	undominated	\$45,217	1.481	\$ 30,537	\$ 60,799	3.040	\$ 20,000	\$58,330	0.693	\$ 84,164
SMV/SOF (12 weeks)	\$ 164,881	\$ 53,854	14.34	-0.16	\$ (345,454)	abs. dominated	\$99,072	1.325	\$ 74,781	\$ 114,654	2.884	\$ 39,754	\$65,532	0.525	\$ 124,933
SOF + R (24 weeks)	\$ 190,636	\$ 79,610	13.62	-0.88	\$ (90,777)	abs. dominated	\$124,828	0.604	\$ 206,759	\$ 140,409	2.163	\$ 64,914			
<b>Tx naïve, treat at F3, F4</b>							<b>Tx naïve, treat at F3, F4</b>								
No Treatment	\$ 50,227	\$ -	11.46	0.00	\$ -	undominated	-\$2,644	-1.219	\$ 2,169						
PR (48 weeks)	\$ 52,871	\$ 2,644	12.68	1.22	\$ 2,169	undominated				\$ 2,644	1.219	\$ 2,169			
LDV/SOF (8/12 weeks)	\$ 71,450	\$ 18,579	13.77	1.10	\$ 16,937	undominated	\$18,579	1.097	\$ 16,937	\$ 21,223	2.316	\$ 9,163			
SOF + PR (12 weeks)	\$ 76,280	\$ 4,830	13.59	-0.19	\$ (25,976)	abs. dominated	\$23,409	0.911	\$ 25,695	\$ 26,053	2.130	\$ 12,230			
LDV/SOF (12 weeks)	\$ 85,452	\$ 14,002	13.81	0.04	\$ 356,143	undominated	\$32,581	1.136	\$ 28,674	\$ 35,225	2.355	\$ 14,954			
SMV/SOF (12 weeks)	\$ 106,551	\$ 21,099	13.65	-0.16	\$ (128,301)	abs. dominated	\$53,680	0.972	\$ 55,237	\$ 56,324	2.191	\$ 25,706			
SOF + R (24 weeks)	\$ 125,105	\$ 39,653	13.10	-0.72	\$ (55,302)	abs. dominated	\$72,234	0.419	\$ 172,296	\$ 74,878	1.638	\$ 45,700			
<b>Tx exp, treat all</b>							<b>Tx exp, treat all</b>						<b>Tx exp, treat all</b>		
No Treatment	\$ 50,227	\$ -	11.46	0.00	\$ -	undominated	-\$26,777	-0.302	\$ 88,615				\$11,405	0.210	\$ 54,254
PR (48 weeks)	\$ 77,004	\$ 26,777	11.76	0.30	\$ 88,615	ext. dominated				\$ 26,777	0.302	\$ 88,615	\$34,025	0.595	\$ 57,208
SOF + PR (12 weeks)	\$ 114,883	\$ 37,880	13.80	2.04	\$ 18,601	ext. dominated	\$37,880	2.036	\$ 18,601	\$ 64,656	2.339	\$ 27,648	\$35,970	0.697	\$ 51,643
LDV/SOF (12/24 weeks)	\$ 128,890	\$ 14,006	14.56	0.76	\$ 18,358	undominated	\$51,886	2.799	\$ 18,535	\$ 78,663	3.101	\$ 25,363	\$58,658	0.643	\$ 91,189
SMV/SOF (12 weeks)	\$ 167,600	\$ 38,710	14.43	-0.13	\$ (294,135)	abs. dominated	\$90,596	2.668	\$ 33,960	\$ 117,373	2.970	\$ 39,521			
<b>Tx exp, treat at F3, F4</b>							<b>Tx exp, treat at F3, F4</b>								
No Treatment	\$ 50,227	\$ -	11.46	0.00	\$ -	undominated	-\$15,371	-0.092	\$ 167,174						
PR (48 weeks)	\$ 65,598	\$ 15,371	11.55	0.09	\$ 167,174	ext. dominated				\$ 15,371	0.092	\$ 167,174			
SOF + PR (12 weeks)	\$ 80,859	\$ 15,260	13.20	1.65	\$ 9,238	undominated	\$15,260	1.652	\$ 9,238	\$ 30,632	1.744	\$ 17,566			
LDV/SOF (12/24 weeks)	\$ 92,920	\$ 12,061	13.86	0.66	\$ 18,242	undominated	\$27,321	2.313	\$ 11,812	\$ 42,693	2.405	\$ 17,752			
SMV/SOF (12 weeks)	\$ 108,942	\$ 16,022	13.79	-0.08	\$ (204,469)	abs. dominated	\$43,344	2.235	\$ 19,396	\$ 58,715	2.327	\$ 25,236			

DR — Discontinuation Rate; Incr Net Cost — Incremental Net Cost; Eff — Effectiveness; Incr Eff — Incremental Effectiveness

**Table F3: Scenario 3: Modified F0-F3 Costs**

**Scenario Analysis** 1. Applying the same costs across F0-F3 (\$900 weighted by frequency = \$1,023/stage)  
**F0-F3 costs** 2. F4 costs remain the same (\$2,516/year)

Incremental comparison of regimens							Vs. PR			Vs. No Treatment			Treat All vs. Treat at F3, F4		
Strategy	Cost	Incr Cost	Eff	Incr Eff	ICER	Comment	Cost	Eff	ICER	Cost	Eff	ICER	Cost	Eff	ICER
<b>Tx naïve, treat all</b>							<b>Tx naïve, treat all</b>						<b>Tx naïve, treat all</b>		
No Treatment	\$ 44,582	\$ -	11.82	0.00	-	undominated	\$ (17,626)	-1.51	\$ 11,649						
PR (48 weeks)	\$ 62,208	\$ 17,626	13.34	1.51	\$ 11,649	undominated				\$17,626	1.51	\$ 11,649	\$13,361	0.37	\$ 36,259
LDV/SOF (8/12 weeks)	\$ 90,962	\$ 28,754	14.75	1.41	\$ 20,347	undominated	\$ 28,754	1.41	\$ 20,347	\$46,380	2.93	\$ 15,849	\$24,122	0.72	\$ 33,301
SOF + PR (12 weeks)	\$ 107,868	\$ 16,906	14.52	-0.23	\$ (73,376)	abs. dominated	\$ 45,660	1.18	\$ 38,604	\$63,286	2.70	\$ 23,475	\$35,764	0.67	\$ 53,754
LDV/SOF (12 weeks)	\$ 108,608	\$ 17,646	14.81	0.06	\$ 284,212	undominated	\$ 46,400	1.48	\$ 31,452	\$64,026	2.99	\$ 21,425	\$26,344	0.74	\$ 35,479
SMV/SOF (12 weeks)	\$ 163,336	\$ 54,727	14.74	-0.08	\$ (719,492)	abs. dominated	\$ 101,127	1.40	\$ 72,275	\$118,753	2.91	\$ 40,776	\$61,956	0.75	\$ 82,411
SOF + R (24 weeks)	\$ 186,333	\$ 77,724	13.99	-0.82	\$ (94,800)	abs. dominated	\$ 124,124	0.66	\$ 189,391	\$141,750	2.17	\$ 65,366	\$70,280	0.57	\$ 123,533
<b>Tx naïve, treat at F3, F4</b>							<b>Tx naïve, treat at F3, F4</b>								
No Treatment	\$ 44,582	\$ -	11.82	0.00	\$ -	undominated	\$ (4,266)	-1.14	\$ 3,726						
PR (48 weeks)	\$ 48,848	\$ 4,266	12.97	1.14	\$ 3,726	undominated				\$4,266	1.14	\$ 3,726			
LDV/SOF (8/12 weeks)	\$ 66,840	\$ 17,992	14.02	1.06	\$ 17,018	undominated	\$ 17,992	1.06	\$ 17,018	\$22,258	2.20	\$ 10,108			
SOF + PR (12 weeks)	\$ 72,105	\$ 5,264	13.85	-0.17	\$ (30,722)	abs. dominated	\$ 23,257	0.89	\$ 26,251	\$27,522	2.03	\$ 13,554			
LDV/SOF (12 weeks)	\$ 82,264	\$ 15,424	14.07	0.04	\$ 351,184	undominated	\$ 33,416	1.10	\$ 30,345	\$37,682	2.25	\$ 16,778			
SMV/SOF (12 weeks)	\$ 101,380	\$ 19,115	13.98	-0.09	\$ (224,046)	abs. dominated	\$ 52,532	1.02	\$ 51,711	\$56,797	2.16	\$ 26,288			
SOF + R (24 weeks)	\$ 116,053	\$ 33,788	13.42	-0.65	\$ (52,283)	abs. dominated	\$ 67,205	0.45	\$ 147,723	\$71,470	1.60	\$ 44,679			
<b>Tx exp, treat all</b>							<b>Tx exp, treat all</b>						<b>Tx exp, treat all</b>		
No Treatment	\$ 44,582	\$ -	11.82	0.00	\$ -	undominated	\$ (27,063)	-0.31	\$ 88,254						
PR (48 weeks)	\$ 71,645	\$ 27,063	12.13	0.31	\$ 88,254	ext. dominated				\$27,063	0.31	\$ 88,254	\$12,353	0.23	\$ 54,077
SOF + PR (12 weeks)	\$ 112,017	\$ 40,372	14.11	1.98	\$ 20,357	ext. dominated	\$ 40,372	1.98	\$ 20,357	\$67,435	2.29	\$ 29,449	\$35,909	0.65	\$ 55,656
LDV/SOF (12/24 weeks)	\$ 119,586	\$ 7,569	14.84	0.72	\$ 10,467	undominated	\$ 47,941	2.71	\$ 17,714	\$75,004	3.01	\$ 24,894	\$37,706	0.76	\$ 49,906
SMV/SOF (12 weeks)	\$ 165,749	\$ 46,163	14.70	-0.14	\$ (341,330)	abs. dominated	\$ 94,104	2.57	\$ 36,601	\$121,167	2.88	\$ 42,105	\$62,430	0.70	\$ 89,470
<b>Tx exp, treat at F3, F4</b>							<b>Tx exp, treat at F3, F4</b>								
No Treatment	\$ 44,582	\$ -	11.82	0.00	\$ -	undominated	\$ (14,710)	-0.08	\$ 188,073						
PR (48 weeks)	\$ 59,292	\$ 14,710	11.90	0.08	\$ 188,073	ext. dominated				\$14,710	0.08	\$ 188,073			
SOF + PR (12 weeks)	\$ 76,108	\$ 16,816	13.47	1.57	\$ 10,735	ext. dominated	\$ 16,816	1.57	\$ 10,735	\$31,526	1.64	\$ 19,169			
LDV/SOF (12/24 weeks)	\$ 81,880	\$ 5,772	14.08	0.61	\$ 9,419	undominated	\$ 22,588	2.18	\$ 10,365	\$37,298	2.26	\$ 16,522			
SMV/SOF (12 weeks)	\$ 103,319	\$ 21,439	14.00	-0.08	\$ (276,706)	abs. dominated	\$ 44,027	2.10	\$ 20,948	\$58,737	2.18	\$ 26,944			

DR — Discontinuation Rate; Incr Net Cost — Incremental Net Cost; Eff — Effectiveness; Incr Eff — Incremental Effectiveness



**Table F4: Scenario 4: Age Set to 50**

**Scenario Analysis** 1. Age set to 50 years old, all other values held constant  
**Age**

Incremental comparison of regimens							Vs. PR			Vs. No Treatment			Treat All vs. Treat at F3, F4		
Strategy	Cost	Incr Cost	Eff	Incr Eff	ICER	Comment	Cost	Eff	ICER	Cost	Eff	ICER	ICER	Eff	ICER
<b>Tx naïve, treat all</b>							<b>Tx naïve, treat all</b>						<b>Tx naïve, treat all</b>		
No Treatment	\$ 60,082	\$ -	14.11	0.00	\$ -	undominated	\$ (10,913)	-2.12	\$ 5,141	\$ 10,913	2.12	\$ 5,141	\$ 10,913	2.12	\$ 5,141
PR (48 weeks)	\$ 70,995	\$ 10,913	16.23	2.12	\$ 5,141	undominated				\$ 34,214	3.98	\$ 8,602	\$ 19,568	0.95	\$ 20,695
LDV/SOF (8/12 weeks)	\$ 94,297	\$ 23,302	18.08	1.85	\$ 12,562	undominated	\$ 23,302	1.85	\$ 12,562	\$ 51,621	4.06	\$ 12,702	\$ 31,579	1.26	\$ 24,982
LDV/SOF (12 weeks)	\$ 111,704	\$ 17,407	18.17	0.09	\$ 201,418	undominated	\$ 40,709	1.94	\$ 20,970	\$ 51,903	3.67	\$ 14,132	\$ 18,197	0.58	\$ 31,436
SOF + PR (12 weeks)	\$ 111,985	\$ 282	17.78	-0.39	\$ (720)	abs. dominated	\$ 40,990	1.55	\$ 26,444	\$ 106,521	3.97	\$ 26,850	\$ 55,391	0.98	\$ 56,399
SMV/SOF (12 weeks)	\$ 166,603	\$ 54,900	18.07	-0.10	\$ (567,135)	abs. dominated	\$ 95,609	1.84	\$ 51,834	\$ 132,747	2.97	\$ 44,769	\$ 62,797	0.74	\$ 84,926
SOF + R (24 weeks)	\$ 192,829	\$ 81,126	17.07	-1.10	\$ (73,829)	abs. dominated	\$ 121,834	0.84	\$ 144,611						
<b>Tx naïve, treat at F3, F4</b>							<b>Tx naïve, treat at F3, F4</b>								
PR (48 weeks)	\$ 59,419	\$ -	15.74	0.00	\$ -	undominated	\$ (664)	1.63	\$ (407)	\$ 664	-1.63	\$ (407)			
No Treatment	\$ 60,082	\$ 664	14.11	-1.63	\$ (407)	abs. dominated				\$ 15,310	1.40	\$ 10,917			
LDV/SOF (8/12 weeks)	\$ 74,728	\$ 15,310	17.14	1.40	\$ 10,917	undominated	\$ 14,646	3.03	\$ 4,830	\$ 20,706	1.17	\$ 17,693			
SOF + PR (12 weeks)	\$ 80,125	\$ 5,397	16.91	-0.23	\$ (23,251)	abs. dominated	\$ 20,042	2.80	\$ 7,158	\$ 34,369	1.46	\$ 23,472			
LDV/SOF (12 weeks)	\$ 93,788	\$ 19,059	17.20	0.06	\$ 308,063	undominated	\$ 33,705	3.09	\$ 10,894	\$ 51,794	1.36	\$ 38,212			
SMV/SOF (12 weeks)	\$ 111,212	\$ 17,424	17.09	-0.11	\$ (160,087)	abs. dominated	\$ 51,130	2.99	\$ 17,128	\$ 70,614	0.60	\$ 118,457			
SOF + R (24 weeks)	\$ 130,033	\$ 36,245	16.33	-0.87	\$ (41,749)	abs. dominated	\$ 69,950	2.23	\$ 31,428						
<b>Tx exp, treat all</b>							<b>Tx exp, treat all</b>						<b>Tx exp, treat all</b>		
No Treatment	\$ 60,082	\$ -	14.11	0.00	\$ -	undominated	\$ (25,146)	-0.49	\$ 51,314	\$ 25,146	0.49	\$ 51,314	\$ 9,611	0.35	\$ 27,164
PR (48 weeks)	\$ 85,229	\$ 25,146	14.60	0.49	\$ 51,314	ext. dominated				\$ 57,590	3.12	\$ 18,437	\$ 31,487	0.86	\$ 36,661
SOF + PR (12 weeks)	\$ 117,673	\$ 32,444	17.23	2.63	\$ 12,320	ext. dominated	\$ 32,444	2.63	\$ 12,320	\$ 62,831	4.10	\$ 15,311	\$ 33,410	0.98	\$ 34,228
LDV/SOF (12/24 weeks)	\$ 122,913	\$ 5,240	18.21	0.98	\$ 5,348	undominated	\$ 37,684	3.61	\$ 10,429	\$ 109,363	3.91	\$ 27,966	\$ 55,843	0.91	\$ 61,243
SMV/SOF (12 weeks)	\$ 169,445	\$ 46,532	18.02	-0.19	\$ (241,216)	abs. dominated	\$ 84,216	3.42	\$ 24,621						
<b>Tx exp, treat at F3, F4</b>							<b>Tx exp, treat at F3, F4</b>								
No Treatment	\$ 60,082	\$ -	14.11	0.00	\$ -	undominated	\$ (15,535)	-0.14	\$ 114,039	\$ 15,535	0.14	\$ 114,039			
PR (48 weeks)	\$ 75,618	\$ 15,535	14.24	0.14	\$ 114,039	ext. dominated				\$ 26,103	2.26	\$ 11,526			
SOF + PR (12 weeks)	\$ 86,185	\$ 10,568	16.37	2.13	\$ 4,965	ext. dominated	\$ 10,568	2.13	\$ 4,965	\$ 29,421	3.13	\$ 9,407			
LDV/SOF (12/24 weeks)	\$ 89,503	\$ 3,318	17.23	0.86	\$ 3,846	undominated	\$ 13,885	2.99	\$ 4,642	\$ 53,520	3.00	\$ 17,847			
SMV/SOF (12 weeks)	\$ 113,602	\$ 24,099	17.10	-0.13	\$ (187,341)	abs. dominated	\$ 37,984	2.86	\$ 13,269						

DR — Discontinuation Rate; Incr Net Cost — Incremental Net Cost; Eff — Effectiveness; Incr Eff — Incremental Effectiveness

## Appendix G: Budgetary Impact Tables

**Table G1. Clinical and Economic Impact of Alternative Treatment Regimens for Hepatitis C, per 1,000 Patients Treated (Treatment-naïve)**

Timeframe/Regimen	Liver-Related Complications				HCV	Costs (per patient, \$)		
	Cirrhosis	Decompensation	HCC	Transplant	Death	Treatment	Other	Total
Treatment-Naïve								
1 Year								
PR	5.6	3.0	1.5	0.0	5.4	\$35,743	\$1,549	\$37,292
LDV/SOF 8/12	0.9	0.7	1.3	0.0	3.3	\$78,095	\$731	\$78,826
Difference (LS-PR)	(4.7)	(2.3)	(0.2)	0.0	(2.1)	\$42,352	(\$818)	\$41,534
5 Years								
PR	29.4	16.0	10.6	0.3	32.1	\$35,743	\$6,136	\$41,879
LDV/SOF 8/12	6.2	3.9	6.8	0.3	19.1	\$78,095	\$3,288	\$81,383
Difference (LS-PR)	(23.2)	(12.1)	(3.8)	0.0	(13.0)	\$42,352	(\$2,848)	\$39,504
20 Years								
PR	104.2	56.9	41.0	4.2	226.6	\$35,743	\$21,236	\$56,979
LDV/SOF 8/12	22.1	13.1	23.2	1.7	110.8	\$78,095	\$10,394	\$88,489
Difference (LS-PR)	(82.1)	(43.8)	(17.8)	(2.5)	(115.8)	\$42,352	(\$10,842)	\$31,510

HCC: Hepatocellular carcinoma; HCV: hepatitis C virus; LS-PR: difference between LDV/SOF and PR therapy

**Table G2. Clinical and Economic Impact of Alternative Treatment Regimens for Hepatitis C, per 1,000 Patients Treated (Treatment-experienced)**

Timeframe/Regimen	Cirrhosis	Liver-Related Complications			HCV Death	Costs (per patient, \$)		
		Decompensation	HCC	Transplant		Treatment	Other	Total
Treatment-Experienced								
1 Year								
PR	11.2	5.6	2.8	0.0	5.5	\$32,044	\$1,963	\$34,007
LDV/SOF 12/24	0.6	0.1	0.8	0.0	3.7	\$107,838	\$563	\$108,401
Difference (LS-PR)	(10.6)	(5.5)	(2.0)	0.0	(1.8)	\$75,794	(\$1,400)	\$74,394
5 Years								
PR	55.2	29.0	16.6	0.7	47.2	\$32,044	\$8,731	\$40,775
LDV/SOF 12/24	5.5	1.5	6.4	0.2	17.4	\$107,838	\$3,153	\$110,991
Difference (LS-PR)	(49.7)	(27.5)	(10.2)	(0.5)	(29.8)	\$75,794	(\$5,578)	\$70,216
20 Years								
PR	183.7	104.0	61.5	7.3	332.4	\$32,044	\$31,743	\$63,787
LDV/SOF 12/24	19.2	6.8	22.1	0.9	102.6	\$107,838	\$9,536	\$117,374
Difference (LS-PR)	(164.5)	(97.2)	(39.4)	(6.4)	(229.8)	\$75,794	(\$22,207)	\$53,587

HCC: Hepatocellular carcinoma; HCV: hepatitis C virus; LS-PR: difference between LDV/SOF and PR therapy

**Table G3. Budget Impact of New Treatment Regimens for Chronic Hepatitis C in the Medi-Cal/Department of Corrections Population in California**

Analysis Step	All Patients				Fibrosis Level 3-4 Only			
	Genotype 1	Genotype 2	Genotype 3	Total	Genotype 1	Genotype 2	Genotype 3	Total
(1) HCV Prevalence	65,100	14,880	11,160	91,140	16,926	3,869	2,902	23,696
(2) # Treated (50%)	32,550	7,440	5,580	45,570	8,463	1,934	1,451	11,848
(3) Interferon Eligibility								
Eligible (60%)	19,530	4,464	3,348	27,342	5,078	1,161	870	7,109
Ineligible (40%)	13,020	2,976	2,232	18,228	3,385	774	580	4,739
(4) Current Total Expenditures (All Care)				\$ 56,456,400,000				\$ 56,456,400,000
PMPM				\$ 611				\$ 611
(5) Increase in HCV Treatment Costs*								
Total \$	\$ 1,607,150,391	\$ 544,712,160	\$ 900,556,200	\$ 3,052,418,751	\$ 417,859,102	\$ 141,625,162	\$ 234,144,612	\$ 793,628,875
PMPM	\$ 17.39	\$ 5.90	\$ 9.75	\$ 33.03	\$ 4.52	\$ 1.53	\$ 2.53	\$ 8.59
% Change	3%	1%	2%	5%	1%	0%	0%	1%
(6) Cost Offsets from New HCV Treatments								
5 Years	\$ (111,363,315)	\$ (85,121,040)	\$ (57,496,320)	\$ (253,980,675)	\$ (28,954,462)	\$ (22,131,470)	\$ (14,949,043)	\$ (66,034,976)
20 Years	\$ (430,592,558)	\$ (475,244,880)	\$ (321,017,400)	\$ (1,226,854,838)	\$ (111,954,065)	\$ (123,563,669)	\$ (83,464,524)	\$ (318,982,258)
(7) Total Net Budgetary Impact								
5 Years	\$ 1,495,787,076	\$ 459,591,120	\$ 843,059,880	\$ 2,798,438,076	\$ 388,904,640	\$ 119,493,691	\$ 219,195,569	\$ 727,593,900
% Change	3%	1%	1%	5%	1%	0%	0%	1%
20 Years	\$ 1,176,557,834	\$ 69,467,280	\$ 579,538,800	\$ 1,825,563,914	\$ 305,905,037	\$ 18,061,493	\$ 150,680,088	\$ 474,646,618
% Change	2%	0%	1%	3%	1%	0%	0%	1%

\*Based on average treatment cost for ledipasvir+sofosbuvir (genotype 1, for 8, 12, or 24 weeks) and sofosbuvir+ribavirin (genotypes 2 and 3, for 12 and 24 weeks respectively)

PMPM: Per-member per-month; HCV: hepatitis C virus