




Tirzepatide for Type 2 Diabetes

Evidence Report

January 6, 2022

Prepared for



AUTHORS:

ICER Staff and Consultants	University of Washington Modeling Group
<p>Grace A. Lin, MD Medical Director for Health Technology Assessment ICER Associate Professor of Medicine and Health Policy University of California, San Francisco</p> <p>Dmitriy Nikitin, MSPH Research Lead, Evidence Synthesis ICER</p> <p>Ashton Moradi, PharmD, MS Health Economist ICER</p> <p>Serina Herron-Smith, BA Senior Research Assistant, Evidence Synthesis ICER</p> <p>Steven D. Pearson, MD, MSc President ICER</p> <p>Jon D. Campbell, PhD, MS Senior Vice President for Health Economics ICER</p>	<p>Elizabeth Brouwer, PhD, MPH Research Scientist The Comparative Health Outcomes, Policy, and Economics (CHOICE) Institute Department of Pharmacy University of Washington</p> <p>Yilin Chen, MPH PhD Student The CHOICE Institute Department of Pharmacy University of Washington</p> <p>Ryan N. Hansen, PharmD, PhD Associate Professor The CHOICE Institute Department of Pharmacy University of Washington</p> <p><i>The role of the University of Washington is limited to the development of the cost-effectiveness model, and the resulting ICER reports do not necessarily represent the views of the University of Washington.</i></p>

DATE OF

PUBLICATION: January 6, 2022

How to cite this document: Lin GA, Brouwer E, Nikitin D, Moradi A, Chen Y, Herron-Smith S, Hansen RN, Pearson SD, Campbell JD. Tirzepatide for Type 2 Diabetes; Evidence Report. Institute for Clinical and Economic Review, January 6, 2022. <https://icer.org/assessment/diabetes-type-2-2022/#timeline>.

Acknowledgements: Grace A. Lin served as the lead author for the report. Dmitriy Nikitin led the systematic review in collaboration with Serina Herron-Smith. Elizabeth Brouwer, Yilin Chen, and Ryan Hansen developed the economic model. Ashton Moradi provided oversight of the cost-effectiveness analyses and developed the budget impact model. Jon D. Campbell and Steven D. Pearson provided methodologic guidance on the clinical and economic evaluations. We would also like to thank Kelsey Gosselin, Maggie Houle, Victoria Lancaster, Rasheed Mohammed, Marina Richardson and Grace Sternklar for their contributions to this report.

About ICER

The Institute for Clinical and Economic Review (ICER) is an independent non-profit research organization that evaluates medical evidence and convenes public deliberative bodies to help stakeholders interpret and apply evidence to improve patient outcomes and control costs. Through all its work, ICER seeks to help create a future in which collaborative efforts to move evidence into action provide the foundation for a more effective, efficient, and just health care system. More information about ICER is available at <https://icer.org/>.

The funding for this report comes from government grants and non-profit foundations, with the largest single funder being the Arnold Ventures. No funding for this work comes from health insurers, pharmacy benefit managers, or life science companies. ICER receives approximately 19% of its overall revenue from these health industry organizations to run a separate Policy Summit program, with funding approximately equally split between insurers/PBMs and life science companies. Life science companies relevant to this review who participate in this program include Boehringer Ingelheim. For a complete list of funders and for more information on ICER's support, please visit <https://icer.org/who-we-are/independent-funding/>.

For drug topics, in addition to receiving recommendations [from the public](#), ICER scans publicly available information and also benefits from a collaboration with [IPD Analytics](#), an independent organization that performs analyses of the emerging drug pipeline for a diverse group of industry stakeholders, including payers, pharmaceutical manufacturers, providers, and wholesalers. IPD provides a tailored report on the drug pipeline on a courtesy basis to ICER but does not prioritize topics for specific ICER assessments.

About the New England CEPAC

The New England Comparative Effectiveness Public Advisory Council (New England CEPAC) – a core program of ICER – provides a public venue in which the evidence on the effectiveness and value of health care services can be discussed with the input of all stakeholders. The New England CEPAC seeks to help patients, clinicians, insurers, and policymakers interpret and use evidence to improve the quality and value of health care.

The New England CEPAC Panel is an independent committee of medical evidence experts from across New England, with a mix of practicing clinicians, methodologists, and leaders in patient engagement and advocacy. All Panel members meet strict conflict of interest guidelines and are convened to discuss the evidence summarized in ICER reports and vote on the comparative clinical effectiveness and value of medical interventions. More information about the New England CEPAC is available at <https://icer.org/who-we-are/people/independent-appraisal-committees/new-england-cepac/>.

The findings contained within this report are current as of the date of publication. Readers should be aware that new evidence may emerge following the publication of this report that could potentially influence the results. ICER may revisit its analyses in a formal update to this report in the future.

The economic models used in ICER reports are intended to compare the clinical outcomes, expected costs, and cost effectiveness of different care pathways for broad groups of patients. Model results therefore represent average findings across patients and should not be presumed to represent the clinical or cost outcomes for any specific patient. In addition, data inputs to ICER models often come from clinical trials; patients in these trials may differ in real-world practice settings.

In the development of this report, ICER’s researchers consulted with several clinical experts, patients, manufacturers, and other stakeholders. The following experts provided input that helped guide the ICER team as we shaped our scope and report. It is possible that expert reviewers may not have had the opportunity to review all portions of this report. None of these individuals is responsible for the final contents of this report, nor should it be assumed that they support any part of it. The report should be viewed as attributable solely to the ICER team and its affiliated researchers.

For a complete list of stakeholders from whom we requested input, please visit:

<https://icer.org/wp-content/uploads/2021/11/Key-Stakeholder-List.pdf>

Expert Reviewers

Todd Boudreaux

Director of Publishing

Beyond Type 1

Beyond Type 1 receives 3.5% of its funding from Eli Lilly and 1% from Novo Nordisk.

Joanna Mitri, MD, MS

Medical Director, Global Education and Care Division

Joslin Diabetes Center

Assistant Professor

Harvard Medical School

Dr. Mitri has received manufacturer support of research in the clinical area of this meeting, and her institution conducts clinical trials and educational programs that may be supported by health care companies. A household member of Dr. Mitri’s has received consulting fees from health care companies including AbbVie, Roche, Janssen Pharmaceuticals, Pharmacyclics, and BeiGene.

Hui Shao, MD, PhD

Assistant Professor, Pharmaceutical Outcomes and Policy Department

University of Florida

Dr. Shao holds a position with BRAVO4Health LLC which receives more than 25% of its funding from health care companies, including Sanofi and AstraZeneca.

Table of Contents

Executive Summary	ES
1. Background	1
2. Patient and Caregiver Perspectives	4
3. Comparative Clinical Effectiveness	6
3.1. Methods Overview	6
Scope of Review	6
Evidence Base	6
3.2. Results	11
Clinical Benefits	11
Harms	16
Subgroup Analyses and Heterogeneity	17
Uncertainty and Controversies	18
3.3. Summary and Comment	20
4. Long-Term Cost Effectiveness	23
4.1. Methods Overview	23
4.2. Key Model Assumptions and Inputs	26
4.3. Results	30
Base-Case Results	30
Sensitivity Analyses	31
Scenario Analyses	33
Threshold Analyses	33
Model Validation	34
Uncertainty and Controversies	34
4.4 Summary and Comment	35
5. Contextual Considerations and Potential Other Benefits	36
6. Health Benefit Price Benchmarks	38
7. Potential Budget Impact	39
7.1. Overview of Key Assumptions	39
7.2. Results	40

References	41
A. Background: Supplemental Information	A1
A1. Definitions.....	A1
A2. Drug Classes for the Treatment of Type 2 Diabetes	A4
A3. Potential Cost-Saving Measures in Type 2 Diabetes	A6
B. Patient Perspectives: Supplemental Information.....	B1
B1. Methods.....	B1
C. Clinical Guidelines	C1
D. Comparative Clinical Effectiveness: Supplemental Information	D1
D1. Detailed Methods	D1
PICOTS.....	D1
Comparators	D1
Data Sources and Searches	D5
Study Selection.....	D8
Data Extraction and Quality Assessment	D9
Assessment of Level of Certainty in Evidence	D10
Assessment of Bias.....	D10
Data Synthesis and Statistical Analyses	D10
D2. Network Meta-Analysis Supplemental Information.....	D10
D3. Additional Clinical Evidence.....	D13
Trials of Tirzepatide.....	D13
Trials of NMA Linkage Studies	D15
Cardiovascular Outcomes Trials (CVOT)	D16
D4. Evidence Tables	D18
D5. Ongoing Studies.....	D48
D6. Previous Systematic Reviews and Technology Assessments	D52
NICE Technology Assessments.....	D52
E. Long-Term Cost-Effectiveness: Supplemental Information	E1
E1. Detailed Methods.....	E1
Target Population.....	E2

E2. Model Inputs and Assumptions	E2
Model Inputs	E2
E3. Results	E9
Description evLYG Calculations.....	E9
E4. Sensitivity Analyses	E11
E5. Scenario Analyses.....	E13
E6. Model Validation.....	E15
F. Potential Budget Impact: Supplemental Information.....	F1
F1. Methods	F1
F2. Results	F3

List of Acronyms and Abbreviations Used in this Report

AACE	American Association of Clinical Endocrinologists
ACCORD	Action to Control Cardiovascular Risk in Diabetes
ACE	American College of Endocrinology
ADA	American Diabetes Association
AE	Adverse event
ASCVD	Atherosclerotic cardiovascular disease
BL	Baseline
BMI	Body Mass Index
bpm	Beats per minute
BT	Background therapy
CEPAC	Comparative Effectiveness Public Advisory Council
CGM	Continuous glucose monitor
CHF	Congestive heart failure
CI	Confidence interval
cm	Centimeter
CKD	Chronic kidney disease
CR	Credible range
CV	Cardiovascular
CVD	Cardiovascular disease
CVO	Cardiovascular outcomes
CVOT	Cardiovascular outcomes trial
DBP	Diastolic blood pressure
D/C	Discontinuation
dL	Deciliter
DPP-4	Dipeptidyl peptidase-4
DTSQc	Diabetes Treatment Satisfaction Questionnaire change version
DUA	Dulaglutide
Dx	Diagnosis
EASD	European Association of the Study of Diabetes
eGFR	Estimated glomerular filtration rate
ESC	European Society of Cardiology
evLY	Equal value of life years
evLYG	Equal value of life years gained
EMPA	Empagliflozin
ER	Emergency room
ESRD	End-stage renal disease
ETD	Estimated treatment difference
FDA	Food and Drug Administration
FPG	Fasting plasma glucose
GIP	Glucose-dependent insulintropic polypeptide
GLP-1	Glucagon-like peptide 1
GLP-1 RA	GLP-1 receptor agonist
HbA1c	Hemoglobin A1c/glycosylated hemoglobin
HDL	High density lipoprotein cholesterol
HF	Heart failure
HFpEF	Heart failure with preserved ejection fraction
HFREF	Heart failure with reduced ejection fraction
HR	Hazard ratio
HUI-3	Health Utilities Index Mark 3
ICER	Institute for Clinical and Economic Review

IGlar	Insulin glargine
IWQOL-Lite-CT	Impact of Weight on Quality of Life-Lite Clinical Trials Version
ITT	Intention to treat
kg	Kilogram
LDL	Low density lipoprotein cholesterol
LOCF	Last observation carried forward
LYG	Life years gained
m	Meter
MACE	Major adverse cardiovascular event
MACE-3	3-point MACE
MACE-4	4-point MACE
MAR	Missing at random
MET	Metformin
mg	Milligram
MI	Myocardial infarction
mITT	Modified intent to treat
mL	Milliliter
MM	Mixed-effects model
mmol	Millimole
mmHg	Millimeter of mercury
MMRM	Mixed-effects model repeated measures
mol	Mole
MTD	Maximum tolerated dose
n	Number of subjects with events
N	Number of subjects in full analysis set
NA	Not applicable
NHANES	National Health and Nutrition Examination Survey
NICE	National Institute for Health and Care Excellence
NR	Not reported
NMA	Network meta-analysis
NON	Noninferiority
OM2	Outcomes Model 2
OR	Odds ratio
OSEM	Oral semaglutide
OW	Once weekly
OWSA	One-way sensitivity analysis
PBO	Placebo
PICOTS	Population, Intervention, Comparator, and Study Design
PK	Pharmacokinetics
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-Analyses
PRO	Patient reported outcome
QALY	Quality-adjusted life year
QW	Once weekly
RA	Receptor agonist
RCT	Randomized controlled trial
REF	Reference
SBP	Systolic blood pressure
SD	Standard deviation
SEM	Semaglutide (injectable)
SF-36	Short Form Health Survey
SGLT-2	Sodium-dependent glucose co-transporter-2
SITA	Sitagliptin

SU	Sulfonylurea
TEAE	Treatment emergent adverse event
T2DM	Type 2 diabetes mellitus
Tx	Treatment
TZD	Thiazolidinedione
TZP	Tirzepatide
UKPDS	United Kingdom Prospective Diabetes Study
US	United States
USD	United States dollar
USPSTF	United States Preventive Services Task Force
VAS	Visual analogue scale
WAC	Wholesale acquisition cost

Executive Summary

Type 2 diabetes mellitus (T2DM) is characterized by the progressive loss of adequate insulin secretion from the pancreas and peripheral insulin resistance. It affects more than 34 million Americans, with minorities bearing a disproportionate burden of disease.¹ Chronic exposure to high blood glucose levels may damage both small (microvascular) and large (macrovascular) blood vessels, and can result in complications such as blindness, chronic kidney disease (CKD), and atherosclerotic cardiovascular disease (ASCVD). Consequently, the annual costs associated with T2DM exceeded \$300 billion in 2017.²

Patients with T2DM described the struggle of managing their disease, including struggles with glycemic control, losing weight, managing comorbidities and disease complications, and the expense of medications. Early and comprehensive education about diabetes self-management, along with access to and affordability of medications, were identified as critical factors in the success of managing T2DM over a patient's lifetime.

A measurable short-run goal of treating T2DM is glycemic control, with a goal glycated hemoglobin (HbA1c) of < 7.0% in most patients.³ Beyond lifestyle modifications, metformin is recommended as first-line therapy based on its efficacy and safety.⁴ Additional therapy is indicated if glycemic goals are not met with metformin alone. For patients with or at high risk of ASCVD, heart failure, or CKD, sodium glucose transporter-2 (SGLT-2) inhibitors and glucagon-like peptide-1 receptor agonists (GLP-1 RA) are preferred due to favorable cardiovascular and renal outcomes data.⁵

Even with current treatment options, nearly half of T2DM patients may not have adequate levels of glycemic control.¹ Tirzepatide (Eli Lilly), a novel, once-weekly injectable dual glucose-dependent insulinotropic polypeptide (GIP) receptor and GLP-1 RA combination drug, has been developed to treat patients with T2DM. A biologics license application with priority review was submitted to the FDA for T2DM on October 27, 2021, with a decision expected in mid-2022.

We compared the clinical and cost effectiveness of tirzepatide added on to background therapy compared with background therapy alone, or injectable semaglutide (Ozempic®, Novo Nordisk) or empagliflozin (Jardiance®, Boehringer Ingelheim and Eli Lilly) added on to background therapy. Treatment with tirzepatide 15 mg resulted in a statistically significant decrease in HbA1c of 2.5% and in weight of 10.9 kg compared with background therapy.⁶ Gastrointestinal symptoms were the most common adverse events; severe hypoglycemia was rare.

Tirzepatide also showed a greater reduction in HbA1c, weight, triglycerides, and blood pressure when compared head-to-head with injectable semaglutide in a Phase 3 randomized controlled trial.⁷ However, the tirzepatide group had a greater incidence of gastrointestinal side effects, severe adverse events, and discontinuation compared with semaglutide.

Due to a lack of head-to-head trials, tirzepatide and empagliflozin were compared through a network meta-analysis. Tirzepatide had a greater decrease in HbA1c, weight loss, lipids, and blood pressure compared with empagliflozin, though there was more uncertainty in these estimates given the indirect comparison. The point estimate decreases in HbA1c and weight loss fell in between the point estimates observed when comparing tirzepatide to injectable semaglutide and comparing tirzepatide to background therapy alone. We did not have data to compare adverse events between tirzepatide and empagliflozin.

Based on data from cardiovascular outcomes trials, semaglutide and empagliflozin have FDA indications for prevention of cardiovascular events.^{8,9} Empagliflozin has also been shown to improve outcomes in patients with CKD.¹⁰ The cardiovascular outcomes trial for tirzepatide is ongoing; however, data from SURPASS-4, a cardiovascular safety trial, showed no increase in cardiovascular events and a trend towards cardiovascular benefit.

Although tirzepatide shows an impressive impact on glucose-lowering and weight loss, given the established cardiovascular benefits of semaglutide and empagliflozin, establishing whether tirzepatide has similar benefit is imperative to reducing uncertainty in its comparative effectiveness. Additionally, although GLP-1 RAs have longer-term safety and cardiovascular data, the impact of the addition of GIP inhibition is currently unknown. Finally, the lack of head-to-head comparison makes it more difficult to fully assess whether tirzepatide provides superior benefit to empagliflozin.

Thus, compared to background therapy alone, we judge the net health benefits of tirzepatide to be incremental or better (B+). For tirzepatide compared with semaglutide, although tirzepatide had greater impact on glycemic control and weight, the lack of definitive cardiovascular outcomes data causes us to judge tirzepatide to have comparable or incremental net health benefits (C+). For tirzepatide compared with empagliflozin, the indirect comparison and lack of definitive cardiovascular or renal outcomes data, causes more uncertainty about the relative benefit (whether it has comparable, small or even substantial health benefit compared with empagliflozin), and thus we judge tirzepatide to have comparable or better net health benefits (C++).

Table ES1. Evidence Ratings

Treatment	Comparator	Evidence Rating
Tirzepatide	Background therapy	B+
Tirzepatide	Injectable Semaglutide	C+
Tirzepatide	Empagliflozin	C++

We developed an individual, patient-level, Monte Carlo-based microsimulation of costs, quality of life, clinical events, and mortality associated with T2DM among US adults using the United Kingdom Prospective Diabetes Study Outcomes Model 2 (UKPDS-OM2)¹¹ equations. Patients, with data from multiple National Health and Nutrition Examination Survey (NHANES) surveys, were simulated through the modeling steps for each comparator versus tirzepatide. The base-case analysis took a

health care sector perspective and thus focused on direct medical care costs only. Costs and outcomes were discounted at 3% per year.¹² All modeled therapies were informed by changes in intermediate outcomes: HbA1c, body weight, LDL, and SBP, predictors in the UKPDS-OM2 risk engine. Modeled cardiovascular and renal outcomes for therapies with existing long-term trials were adjusted to trial data using trial-based hazard ratios. We adjusted tirzepatide’s modeled composite MACE outcomes based on SURPASS-4 trial’s pooled dosing analysis hazard ratio and its uncertainty. Where possible, we compared the treatment-specific modeled events to that of comparable time horizons from long-term trials.

Tirzepatide had the highest average lifetime discounted quality-adjusted life-years (QALYs) of all considered therapies, however the QALY 95% credible ranges for active comparators overlapped. Equal value of life years (evLY) gained was not reported given no average increased survival when comparing tirzepatide with injectable semaglutide. Using a placeholder price equal to injectable semaglutide, the incremental costs per QALY gained for tirzepatide were around or under \$100,000 versus all comparators with mean differences in health gains and costs being smallest in comparison to injectable semaglutide. Uncertainty analyses suggested a wide range of plausible cost-effectiveness estimates for tirzepatide.

ICER’s Health Benefit Price Benchmarks (HBPBs) are defined as the target prices for a drug that would meet but not surpass benchmarks tied to incremental cost-effectiveness ratios of \$100,000 and \$150,000 per QALY or per evLY gained. Table ES2 illustrates the annual HBPBs for tirzepatide plus background therapy as compared to semaglutide plus background therapy, ranging from an annual price for tirzepatide of \$5,500 to \$5,700. Price reductions or discounts from a list price to reach a HBPB point estimate or range is not applicable for tirzepatide as its list price is currently not available. HBPBs based on evLY gained are not reported given no modeled tirzepatide survival gains versus injectable semaglutide.

Table ES2. Annual Cost-Effectiveness Health Benefit Price Benchmarks for Tirzepatide

Outcome for Annual HBPB Calculation	Annual WAC	Annual Price at \$100,000 Threshold	Annual Price at \$150,000 Threshold	Discount from WAC to Reach Threshold Prices
Tirzepatide plus Background Therapy vs. Semaglutide plus Background Therapy				
QALYs Gained	NA*	\$5,500	\$5,700	NA*

HBPB: health benefit price benchmark, QALY: quality-adjusted life year, WAC: wholesale acquisition cost,
*Not applicable (NA) as placeholder prices were used

In summary, tirzepatide improves blood glucose levels and results in weight loss to a greater extent than other diabetes medications. Although early results are promising, definitive data are needed to understand tirzepatide’s impact on cardiovascular and renal outcomes. Additionally, because T2DM is more prevalent among minorities, health gains from a successful treatment that has consistent benefits across racial subgroups would provide proportionally greater benefit to those racial groups. Studies have not adequately enrolled minority populations to demonstrate such a

consistent effect. Based on current evidence and when compared to injectable semaglutide, the estimated annualized health benefit price benchmark range for tirzepatide is \$5,500 to \$5,700. This range factors in assumptions about long-term cardiovascular benefits that have not been directly demonstrated yet in clinical trials.

1. Background

More than 34 million Americans, or around 13% of the US population, have diabetes mellitus.¹ T2DM, which is characterized by progressive loss of adequate insulin secretion from the pancreas and peripheral resistance to insulin, accounts for 90-95% of those cases. The hallmark of diabetes is an abnormal elevation in blood glucose, or hyperglycemia. Chronic hyperglycemia puts patients at risk for damage to both small (microvascular) and large (macrovascular) blood vessels, resulting in damage to the eyes, nerves, and kidneys, as well as cardiovascular events and limb ischemia.¹³ Obesity is a major risk factor for developing T2DM with more than 60% of people with diabetes having a body mass index (BMI) in the obese or extreme obesity range.^{1,14} Other risk factors include older age, family history, a history of gestational diabetes, high blood pressure, a sedentary lifestyle, and a low level of HDL.¹⁵

Minorities bear a disproportionate burden of T2DM. Among adults, the prevalence of diagnosed T2DM is highest in American Indian/Alaska Natives (approximately 14%); for Asian Indian, non-Hispanic Black and Hispanic populations, prevalence estimates are around 12% for each group.¹ These populations are also at greater risk of developing diabetes, with the incidence in Black and Hispanic populations 1.7-1.8 times greater than in White populations.^{1,16} For example, Blacks and Hispanics had lower rates of receipt for HbA1c testing than Whites and were less likely to receive annual cholesterol testing and retinal examination, which may be correlated with higher rates of complications in these populations.¹⁷

Complications of diabetes can be severe and life-altering. Diabetes is the leading cause of new blindness, end-stage renal disease, and limb amputations, and it is the seventh leading cause of death in the US.¹ In 2016, diabetes was associated with 16 million emergency department visits, 7.8 million hospitalizations, and 1.7 million hospitalizations for cardiovascular disease.¹ Total direct and indirect costs for diabetes exceeded \$327 billion in 2017, with 72% of the total considered direct costs (e.g., health care services or technologies).² Furthermore, patients with diabetes are more likely to suffer financial hardship from medical bills and its consequences, including cost-related medication nonadherence, delayed or skipped medical care, high financial distress, and food insecurity.¹⁸

Management of T2DM is focused on both shorter-term goals such as controlling hyperglycemia and managing comorbidities, and longer-term goals such as preventing complications of disease. Treatment goals are based in part on measurements of HbA1c, a measure of average blood sugar over three months. Since tight control of blood sugar has been shown to decrease microvascular and macrovascular complications, clinical practice guidelines recommend a HbA1c target of 7.0% or less in most patients, with less stringent control accepted in patients with a higher risk of hypoglycemia, more severe comorbidities, and shorter life expectancy.¹⁹ Achieving near-

normal glycemic control with intensive therapy may have additional benefits in terms of slowing onset or progression of neuropathy, retinopathy, and nephropathy.²⁰ However, such tight control of glucose may come at the expense of more hypoglycemia.

The cornerstones of therapy for T2DM includes lifestyle modifications such as diet, exercise, and weight loss, and management of cardiovascular risk factors such as high cholesterol and high blood pressure.^{19,21} Self-management is a critical component of managing diabetes, and individually tailored, culturally appropriate diabetes self-management education can improve outcomes and reduce costs.^{22,23} For example, medical nutrition therapy delivered by a registered dietician or diabetic educator is associated with a 0.3-2% decrease in HbA1c in patients with T2DM.¹⁹ For some patients, intensive lifestyle changes may be enough to control blood sugars and prevent progression. However, many patients will require pharmacologic therapy during their disease course to achieve adequate glycemic control.

Metformin is recommended as initial pharmacotherapy for patients with T2DM due to its efficacy and favorable safety profile and should be continued as long as it is tolerated and not contraindicated. It is associated with modest weight loss, does not have a significant risk of hypoglycemia, and may improve cardiovascular outcomes, though there are no direct cardiovascular outcomes trials.²⁴ Combination therapy with additional agents can be considered if patients do not meet their HbA1c goal on metformin and lifestyle changes. Older oral medications such as sulfonylureas (SU) and thiazolidinediones (TZD) are inexpensive but have significant side effects such as hypoglycemia (with SU) and heart failure (with TZDs). Newer agents include oral agents such as dipeptidyl peptidase-4 (DPP-4) inhibitors and SGLT-2 inhibitors, and oral and injectable GLP-1 RA. These newer agents are effective at lowering blood glucose without a substantial risk of hypoglycemia, and SGLT-2 inhibitors and GLP-1 RAs are also associated with weight loss, reduction in cardiovascular events, and slowed progression of renal disease. Gastrointestinal side effects are particularly common with GLP-1 RAs and use of SGLT-2 inhibitors are associated with genitourinary infections and ketoacidosis.²⁵⁻²⁸

Since 2008, when the Food and Drug Administration (FDA) issued recommendations for the evaluation of cardiovascular risk for new antihyperglycemic therapies, multiple cardiovascular outcome trials (CVOTs) have been conducted, adding greater certainty in the assessment of the relative risks and benefits of each therapy.²⁹ Thus, for patients with or at high risk of ASCVD, heart failure, or established chronic kidney disease (CKD) recommended second-line therapy includes either a GLP-1 RA or SGLT-2 inhibitor with proven cardiovascular or renal benefit, independent of HbA1c target or baseline metformin use.⁵ For patients without those comorbidities, choice of drug is dependent on comorbidities, cost, side effects, and patient preferences. More details on each drug class can be found in the [Supplement A2](#).

Although many options for therapy for T2DM exist, nearly half of patients may not be at an adequate level of glycemic control.¹ Tirzepatide (Eli Lilly), a novel, once-weekly injectable dual GIP and GLP-1 receptor agonist, has been developed to treat patients with T2DM. Trials are also ongoing to assess tirzepatide for the treatment of obesity and cardiovascular disease. The manufacturer announced the submission of a biologics license application with priority review to the FDA for T2DM on October 27, 2021, with a decision expected in mid-2022.

Table 1.1. Interventions of Interest

Intervention Brand Name (Generic Name)	Mechanism of Action	Delivery Route	Prescribing Information
Tirzepatide	Dual GIP inhibitor/GLP-1 receptor agonist	Subcutaneous injection	5-15 mg once weekly

GIP: glucose-dependent insulinotropic polypeptide, GLP-1: glucagon-like peptide-1, mg: milligrams

2. Patient and Caregiver Perspectives

This evidence report was developed with input from diverse stakeholders, including patients, clinicians, researchers, and manufacturers of the agents of focus in this review.

Patients and patient groups discussed that T2DM has a substantial impact on daily life, including trying to manage diet and blood glucose readings, affording testing supplies and medications, and managing complications of the disease. Patients described the frustrations of trying to follow prescribed diets, particularly when the dietary recommendations conflict (e.g., patients with diabetes are recommended to limit carbohydrates while patients with CKD are recommended to avoid a high protein diet). Glycemic control, particularly HbA1c levels, is one of the most important actionable and short-run outcomes to patients, and patients value potentially getting to near normal glycemic levels.³⁰ Improvements in glucose tracking with new technologies such as continuous glucose monitors (CGM) are empowering to patients to help manage their daily life and achieve their glycemic goal. Weight loss is another critical struggle described by patients, and patients would welcome medications that would produce significant weight loss. Patients also spoke about feelings of guilt and self-blame for their diabetes, describing the stigma they feel about the perception of diabetes as a “lifestyle” disease that they should be able to control with their actions, even though the causes of diabetes are complex, and genetics can play an important role.

Affordability of diabetes medications and access to CGMs were cited as major barriers to managing T2DM, particularly for patients who are taking medications. Medication costs for patients with diabetes can be substantial, even for those who are insured, and this can affect adherence to medication. For example, co-pays and co-insurance for insulin and newer medications like GLP-1 RAs can be very high, particularly for patients with high deductible health plans, even if they successfully navigate a time-consuming prior authorization process. For patients with lower socioeconomic status, the inability to pay can lead to skipping medication doses and cancelling medical appointments, which may be associated with adverse clinical outcomes.³¹ Additionally, patients often have multiple comorbidities such as cardiac or renal disease, which can further increase the financial burden of medications. Patients may turn to foundations or prescription assistance programs from manufacturers to help with medication costs; however, patients with Medicare are usually not eligible for manufacturer prescription assistance programs so such programs have limited reach. Finally, although patients described new technologies such as CGMs as “game changers” in terms of helping them manage their disease, obtaining insurance coverage for CGMs can be very challenging.

Patients discussed that another unmet need is more detailed and comprehensive education about disease management and trajectory, particularly in the time period around diagnosis. In addition, linguistically and culturally sensitive education delivered within the community appears to be a key

to improving patient engagement and self-management of the disease. This is particularly important because the burden of diabetes is substantial in minority populations, with Native Americans/Alaska Natives, Blacks, Hispanics, and Asians having a higher rate of disease compared with White Americans. Thus, patient groups mentioned that a greater focus on efforts to decrease disparities in prevention and treatment of diabetes and accurate modeling of the health effects of diabetes are critical to improving health equity.

Clinicians were enthusiastic about the prospect of therapies that both improved glycemic control and resulted in substantial weight loss. However, they emphasized the importance of cardiovascular outcomes on their assessments of efficacy and were eagerly awaiting the results of the CVOT for tirzepatide. Cost of therapy for patients was an additional issue of great importance to both clinicians and patients and was cited as a major factor in the choice of therapy.

Manufacturers and payers highlighted additional non-glycemic outcomes that may be important to consider when assessing the value of therapy, including improvements in cardiovascular and renal outcomes, satisfaction with the delivery device, and the ability of the therapy to address comorbidities related to obesity that may be impacted by weight loss (e.g., obstructive sleep apnea, nonalcoholic fatty liver disease, etc.).

3. Comparative Clinical Effectiveness

3.1. Methods Overview

Procedures for the systematic literature review assessing the evidence of tirzepatide for T2DM are detailed in [Supplement D1](#).

Scope of Review

We reviewed the clinical effectiveness of tirzepatide added on to background therapy (metformin +/- sulfonylureas or thiazolidinediones) versus background therapy alone, or injectable semaglutide or empagliflozin added on to background therapy in adults with T2DM with inadequate glycemic control despite current treatment with antihyperglycemic agent(s). We sought evidence on patient-important outcomes such as change in HbA1c levels, weight, waist circumference, lipid levels, blood pressure, health-related quality of life measures, and adverse events. Additionally, we reviewed available information on each treatment's effect on micro- and macrovascular outcomes including retinopathy, nephropathy, and neuropathy, all-cause mortality, cardiovascular (CV) mortality, myocardial infarction (MI), stroke, and heart failure requiring hospitalization.

We also looked for data on subpopulations of interest, including T2DM patients with (1) established ASCVD, (2) congestive heart failure, (3) moderate-to-severe renal impairment, (4) requiring a second and/or third antihyperglycemic agent, (5) who are overweight or obese, and (6) identified by race, ethnicity, or socioeconomic status.

The full scope of the review is detailed in [Supplement D1](#).

Evidence Base

The SURPASS clinical trial program contains five Phase 3 global registration studies comparing tirzepatide against placebo (background diet and exercise alone), injectable semaglutide, insulin degludec, insulin glargine, and background insulin glargine ± metformin.³² SURPASS-2 and SURPASS-4 were selected as studies of interest due to a relevant comparator arm or population and are outlined below. We also reviewed two Phase 2 trials to assess the efficacy of tirzepatide added on to background therapy compared with background therapy alone.

Tirzepatide versus Background Therapy

Our search identified two randomized controlled trials (RCT) of tirzepatide that provide evidence on the efficacy and safety of tirzepatide plus background therapy (diet and exercise alone or with stable metformin treatment) versus background therapy alone. These include a Phase 2b study comparing once-weekly injectable tirzepatide (1, 5, 10, or 15 mg) versus once-weekly injectable dulaglutide (1.5 mg) and placebo (Frias 2018)⁶, and a follow-up Phase 2 study evaluating the tolerability of three tirzepatide dose-escalation regimens (12 mg or two versions of a 15 mg arm) versus placebo (Frias 2020).³³ Participants in both trials were largely on stable metformin doses for background therapy. Baseline characteristics of the two trials are outlined in Table 3.1. The two Phase 2 trials were qualitatively assessed for evidence on the efficacy and safety of tirzepatide as an add-on to background therapy.

Table 3.1 Selected Baseline Characteristics of Two Phase 2 Tirzepatide Trials

	Frias 2018 Phase 2				Frias 2020 Phase 2		
Study Arms	PBO	TZP 5 mg	TZP 10 mg	TZP 15 mg	PBO	TZP 15 mg-1*	TZP 15 mg-2*
Background Therapy	MET				MET		
Study Duration, weeks	26 weeks				12 weeks		
Mean Age, years	56.6	57.9	56.5	56	56	55.5	56.6
Sex, Male, %	57	62	59	42	46.2	57.1	82.1
HbA1c, %	8	8.2	8.2	8.1	8.2	8.5	8.4
Weight, kg	91.5	92.8	92.7	89.1	89.6	88.7	89.8
BMI, kg/m ²	32.4	32.9	32.6	32.2	32.5	32	31.1
Race, White, %	80	84	74	81	NR	NR	NR
Metformin, %	92.2	89.1	86.3	96.2	88.5	89.3	82.1

kg: kilogram, m: meter, MET: metformin, mg: milligram, NR: not reported, PBO: placebo, TZP: tirzepatide

*In the Frias 2020 trial, there were two versions of a 15 mg arm. See [Supplement D3](#) for further details.

Additionally, we conducted a network meta-analysis that provided an indirect comparison of tirzepatide added to background therapy versus background therapy alone using five Phase 3 trials. The NMA is described in further detail in the evidence base for the tirzepatide versus empagliflozin scenario and in [Supplement D2](#).

Tirzepatide versus Semaglutide

SURPASS-2 is a Phase 3 head-to-head open-label trial evaluating the clinical efficacy and safety of tirzepatide (5, 10, or 15 mg) versus injectable semaglutide (1 mg) over a 40-week period in patients with T2DM inadequately controlled with baseline metformin therapy. Baseline characteristics can be found in Table 3.2, and additional details of the study design can be found in [Supplemental Table D4.3](#).

Tirzepatide versus Empagliflozin

There are no head-to-head trials comparing the clinical efficacy and safety of tirzepatide versus empagliflozin. Thus, to be able to draw an indirect comparison between the two treatments, we conducted a NMA on the available intermediate outcomes of changes in HbA1c, weight, low-density lipoprotein (LDL) cholesterol, and systolic blood pressure (SBP). These outcomes were presented as between-treatment mean differences of change from baseline at 40 weeks. We included five Phase 3 trials and five drugs into the network: tirzepatide (SURPASS-2), injectable semaglutide (SURPASS-2, SUSTAIN-2), oral semaglutide (PIONEER-2, PIONEER-3), empagliflozin (PIONEER-2), and sitagliptin (SUSTAIN-2, PIONEER-3, HARMONY-3). The study design and baseline characteristics of the five trials are outlined in Table 3.2.

Table 3.2 Selected Study Design and Baseline Characteristics of the Randomized Controlled Trials in the Network Meta Analysis Evaluating Tirzepatide, Semaglutide, Empagliflozin, and Background Therapy

	SURPASS-2		SUSTAIN-2		PIONEER-2		PIONEER-3		HARMONY-3	
Study Arms	TZP 15 mg	SEM 1 mg	SEM 1 mg	SITA 100 mg	OSEM 14 mg	EMPA 25 mg	OSEM 14 mg	SITA 100 mg	PBO	SITA 100 mg
N	470	469	409	407	411	410	465	467	101	302
Background Therapy	MET		MET ± TZD (4.5%)		MET		MET ± SU (47.1%)		MET	
Study Duration	40 weeks		56 weeks		52 weeks		78 weeks		104 weeks	
Mean Age, years	55.9	56.9	56	54.6	57	58	57	58	56.1	54.3
Sex, Male, %	45.5	48	50	51	50.1	51	53.1	51.0	49.5	46
HbA1c, %	8.3	8.3	8	8.2	8.1	8.1	8.3	8.3	8.2	8.1
Weight, kg	93.8	93.7	89.2	89.3	91.9	91.3	91.2	90.9	91.6	90.3
BMI, kg/m ²	34.5	34.2	32.5	32.5	32.9	32.8	32.3	32.5	32.8	32.5
Race, White, %	71.1	71.6	68	69	86.4	86.1	68.2	71.3	63.4	74.5
Metformin, %	100	100	100	100	100	100	100	100	100	100

EMPA: empagliflozin, kg: kilogram, m: meter, MET: metformin, N: number of subjects in full analysis set, OSEM: oral semaglutide, PBO: placebo/background therapy, SEM: injectable semaglutide, SITA: sitagliptin, SU: sulfonylurea, TZD: Thiazolidinediones, TZP: tirzepatide

Cardiovascular Outcomes Trials

Tirzepatide

The SURPASS-CVOT is evaluating the non-inferiority and superiority of once weekly tirzepatide versus dulaglutide (1.5mg) in participants with T2DM and increased cardiovascular risk.³⁴ The trial has an estimated completion date of October 17, 2024; interim data was unavailable at the time of this report.³⁴

In lieu of SURPASS-CVOT data, we sought data on baseline characteristics, efficacy, and adverse event parameters of SURPASS-4, a Phase 3 trial that evaluated three doses of tirzepatide (5, 10, 15 mg) against insulin glargine in adults with T2DM on 1-3 oral antihyperglycemic medications and increased cardiovascular risk.³⁵ Cardiovascular events were recorded as safety events. Additional details of the trial are available in [Supplemental Table D4.2](#).

Due to the unavailability of long-term CVOT data and lack of placebo arm in the above tirzepatide trials, we were unable to include any microvascular (neuropathy, nephropathy, and retinopathy) or macrovascular outcomes (death from cardiovascular causes, nonfatal myocardial infarction, nonfatal stroke, or hospitalization for unstable angina or heart failure) in our NMA.

Injectable Semaglutide

SUSTAIN-6 was a CVOT that assessed the efficacy and safety of semaglutide (0.5 or 1 mg) versus placebo as an add-on therapy to background therapy (73.2% on metformin, 58% on insulin, 42.8% on sulfonylurea) in patients with T2DM and established ASCVD, CKD, or both.⁸ A total of 3297 patients with a mean age of 64.6 years and baseline HbA1c of 8.7% were observed for a median period of 2.1 years.⁸ The pooled semaglutide group was analyzed for non-inferiority and superiority versus the placebo group for the primary composite outcome of MACE-3 (death from cardiovascular causes, nonfatal myocardial infarction, or nonfatal stroke).⁸ Additional details of the trial are available in [Supplemental Table D4.2](#).

Empagliflozin

EMPA-REG OUTCOME was a CVOT that assessed the efficacy and safety of empagliflozin (10 or 25 mg) as an add-on to background therapy (74% on metformin, 48% on insulin, 43% on sulfonylurea) in patients with T2DM and established cardiovascular disease.^{9,36} In this trial, 7020 patients with a mean age of 63.1 years and a baseline HbA1c of 8.1% were treated and observed for a median period of 2.6 and 3.1 years, respectively.⁹ The pooled empagliflozin group was analyzed for non-inferiority and superiority versus the placebo group for the primary composite outcome of MACE-3 (death from cardiovascular causes, nonfatal myocardial infarction, or nonfatal stroke) and the key secondary composite outcome of MACE-3 or hospitalization for unstable angina.⁹ Additional details of the trial are available in [Supplemental Table D4.2](#).

Subgroup Analysis of Patient-Important Outcomes

Our population of interest for this review are adults with T2DM with inadequate glycemic control despite current treatment with background therapy. In addition to our primary analysis, we sought to evaluate all available patient-important outcomes in the context of the following six patient subgroups:

1. Established ASCVD, e.g., a history of acute coronary syndrome (ACS), MI, stable or unstable angina, coronary or other arterial revascularization, stroke, transient ischemic attack, or peripheral arterial disease.³⁷

Currently, there is no available evidence to evaluate the clinical efficacy or safety of tirzepatide for this subgroup. The SURPASS-4 trial included participants with an increased risk of ASCVD as well as participants with established ASCVD; cardiovascular events were recorded as safety events.³⁵ When available, we qualitatively assessed any subgroup analyses done in this population in the semaglutide and empagliflozin CVOTs.

2. Established congestive heart failure (New York Heart Association class II or III)

Currently, there is no available evidence to evaluate the clinical efficacy and safety of tirzepatide for this subgroup. When available, we qualitatively assessed any subgroup analyses done in this population in the semaglutide and empagliflozin CVOTs.

3. Established moderate-to-severe renal impairment (CKD Stage 3 or higher (eGFR <60 mL/min per 1.73 m²))

Academic-in-confidence data on change in HbA1c (%) from baseline at week 40 between the subgroups of eGFR <60 and ≥60 mL/min per 1.73 m² for the SURPASS-2 trial was provided by the manufacturer.³⁸ When available, we qualitatively assessed any subgroup analyses done in this population in the semaglutide and empagliflozin CVOTs.

4. Tirzepatide as a second- or third-line antihyperglycemic agent

We did not find any data assessing the efficacy or safety of tirzepatide specific to these two subpopulations. When available, we qualitatively assessed any subgroup analyses done in this population in the semaglutide and empagliflozin CVOTs.

5. Overweight (BMI 25.0-29.9 kg/m²) or obese (≥30.0 kg/m²)

Academic-in-confidence data on change in HbA1c (%) from baseline at week 40 between the subgroups of BMI <30 kg/m², ≥30 to <35 kg/m², and ≥35 kg/m² was provided by the manufacturer for the SURPASS-2 trial.³⁸ When available, we qualitatively assessed any subgroup analyses done in these populations the semaglutide and empagliflozin CVOTs.

6. Race and ethnicity or socioeconomic status

Academic-in-confidence data on change in HbA1c (%) from baseline at week 40 between the subgroups of Hispanic/Latino and Not Hispanic/Latino was provided by the manufacturer for the SURPASS-2 trial.³⁸ When available, we qualitatively assessed any subgroup analyses done in this population in the semaglutide and empagliflozin CVOTs.

3.2. Results

A subset of trials within our review report treatment effects using two estimands. The treatment-regimen, also called treatment policy, estimand evaluates the treatment effect for all randomized patients regardless of premature trial product discontinuation or use of rescue medication.^{7,39} The efficacy, or trial product, estimand evaluates the treatment effect of all randomized patients who had completed the study without the use of rescue medication. The treatment-regimen estimand and the efficacy estimand provide different perspectives on a drug's efficacy; we feel that the treatment-regimen/treatment policy estimand is more relevant to the patient and clinician experience as it considers the common challenges of medication adherence and need for rescue medication in the diabetes treatment realm.

When possible, we prioritized the reporting of values from the treatment-regimen estimand for several reasons: to ensure consistency with our 2019 ICER Type 2 Diabetes report, the estimand's reflection of the intention-to-treat principle, and its preference by regulatory agencies such as the FDA.⁴⁰ If the treatment-regimen estimand was not available, we reported efficacy estimands and have noted as such via in-text and evidence table references.

We conducted an NMA comparing tirzepatide added to background therapy to background therapy alone, and injectable semaglutide or empagliflozin added to background therapy. Oral semaglutide and sitagliptin were used as linkages in the NMA and therefore were not emphasized in this report. Additionally, due to limited data reported in publications and/or provided by manufacturers, there were only four available outcomes for which we could provide inputs into the NMA: change from baseline in HbA1c, body weight, LDL, and SBP at week 40. ([Supplemental Tables D2.2-2.3](#))

Clinical Benefits

Tirzepatide versus Background Therapy

The efficacy of tirzepatide compared with background therapy was evaluated through two Phase 2 trials (Frias 2018 and Frias 2020) and the NMA. Tirzepatide consistently showed a dose-dependent decrease in HbA1c of -1.6% to -2.4% compared with background therapy (efficacy estimand). Results from the NMA were consistent with these trials, with an estimated treatment difference in HbA1c from placebo of -1.7%. Furthermore, in the Frias 2018 trial, all three tirzepatide doses had a

greater proportion of participants who achieved a HbA1c target of <7.0% (69.1-90.0%) compared to placebo (11.8%) ($p<.0001$ for all three comparisons), and 30.2% of participants in the tirzepatide 15 mg arm achieved HbA1c levels of <5.7%, an indication of normal glycemic control. ([Supplement Table D4.4](#))

In terms of weight, tirzepatide also showed a dose-dependent decrease in weight ([Supplement Table D4.4](#)), with the greatest weight loss seen in the tirzepatide 15 mg group (-11.3 kg from baseline in the Frias 2018 trial and -9.51 kg difference from placebo in the NMA). Additionally, in the Frias 2018 trial, more than one-third of patients in the 10 mg and 15 mg groups achieved body weight reduction of $\geq 10\%$ and almost one-quarter of participants in the 15 mg arm achieved $\geq 15\%$ reduction. Impact on waist circumference was mixed between the two trials. ([Supplement Table D4.4](#))

The impact of tirzepatide on other outcomes was mixed. Tirzepatide had a statistically significant decrease of 7.46 mmHg in systolic blood pressure compared with placebo in the NMA (random effects model) but not the Phase 2 trials. Similarly, tirzepatide had a statistically significant decrease in LDL of 4.33 mg/dL from baseline in the NMA (random effects model) that was not seen in the Phase 2 trials. Tirzepatide also decreased mean total cholesterol and triglycerides concentrations ($p<.05$ for all interactions) in the Frias 2018 trial ([Supplement Table D4.4](#)).

Tirzepatide versus Semaglutide

The primary outcome of the SURPASS-2 trial was the mean change in HbA1c from baseline to 40 weeks between three arms of tirzepatide (5 mg, 10 mg, or 15mg) and injectable semaglutide (1 mg). Tirzepatide reduced HbA1c levels by 2%, 2.2%, and 2.3% in the 5 mg, 10 mg, and 15 mg dose groups, respectively, compared with 1.86% with semaglutide. The estimated treatment differences for all groups compared with semaglutide were statistically significant: tirzepatide 5 mg -0.15% (95% CI -0.28 to -0.03); tirzepatide 10 mg -0.39% (-0.51 to -0.26); tirzepatide 15 mg -0.45% (-0.57 to -0.32).

Additionally, the two larger tirzepatide doses (10 mg, 15 mg) also had a greater proportion of participants who achieved a HbA1c target of <7.0% (86%) compared to semaglutide (79%) ($p<.05$ for both dose groups). Nearly half of participants in the tirzepatide 10 mg and 15 mg arms achieved near-normal glucose levels (HbA1c <5.7%), compared with 19% in the semaglutide arm, $p<.001$).

Treatment with tirzepatide produced a dose-dependent change in mean body weight (kg) from baseline compared with semaglutide. At week 40, participants experienced a weight loss of 7.6 kg (5 mg), 9.3 kg (10 mg), and 11.2 kg (15 mg) compared with 5.7 kg in the semaglutide group ($p<.0001$ for all three comparisons). The percentage of participants achieving body weight reduction of $\geq 10\%$ was 24% (semaglutide), 34% (5 mg), 47% (10 mg), and 57% (15 mg), and 36% of participants in the 15 mg arm achieved a $\geq 15\%$ reduction in body weight. Finally, at week 40, there was a greater

reduction in mean waist circumference (cm) in all three dosage arms of tirzepatide compared with semaglutide ([Supplement Table D4.5](#)).

The 15 mg tirzepatide arm had a greater reduction in systolic and diastolic blood pressure from baseline (–6.5 mmHg and –2.9 mmHg, respectively) as compared with 1 mg semaglutide (–3.6 mmHg and –1.0 mmHg, respectively). Treatment with tirzepatide also resulted in a greater reduction in mean HDL and triglycerides concentrations versus semaglutide across the three dosage arms (efficacy estimand) ([Supplement Table D4.5](#)). Changes in mean total cholesterol and LDL concentrations were not statistically significant among the four treatment groups (efficacy estimand).

Treatment with 15 mg of tirzepatide resulted in better overall quality of life than semaglutide across several quality-of-life measures, including the Diabetes Treatment Satisfaction Questionnaire change version (DTSQc), EQ-5D-5L (index score), EQ-5D-5L visual analogue scale (VAS), Impact of Weight on Quality of Life-Lite Clinical Trials Version (IWQOL-Lite-CT) (psychosocial, physical, and physical functioning score).³⁸ (data on file)

Tirzepatide versus Empagliflozin

As there are no head-to-head trials comparing tirzepatide to empagliflozin, all results for this comparison are derived from the NMA. Compared with empagliflozin, tirzepatide decreased HbA1c by 1.1% (statistically significant change using random effects model) ([Supplemental Table D2.4](#)). A similar advantage is seen in the comparison of weight loss, with an estimated mean difference of –7.2kg between tirzepatide and empagliflozin (statistically significant change using random effects model) ([Supplemental Table D2.4](#)).

Tirzepatide appears to decrease systolic blood pressure by 2.6 mmHg compared with empagliflozin (statistically significant change using random effects model) ([Supplemental Table D2.4](#)). For the outcome of LDL, tirzepatide appears to lower LDL by 7.5 mg/dL compared with empagliflozin (statistically significant change using random effects model) (efficacy estimand)([Supplemental Table D2.4](#)).

Cardiovascular Outcomes

Tirzepatide

Although the CVOT for tirzepatide is ongoing, the manufacturer performed a CV safety meta-analysis within SURPASS-4 and across the five SURPASS trials on the adjudicated composite outcome (MACE-4) of death from cardiovascular or undetermined causes, myocardial infarction, stroke, and hospitalization for unstable angina. In SURPASS-4, there was no evidence of an increase in the occurrence of MACE-4 and there was an overall trend towards benefit (HR: 0.74; 95% CI: 0.51 to 1.08; p=NR).³⁵ Additionally, the tirzepatide 15 mg group showed a decrease in MACE-4 (HR 0.50;

95% CI 0.26-0.95) over a median of 85 weeks of follow-up.³⁵ Likewise, pooled data from the SURPASS program showed that tirzepatide had a trend towards benefit in the occurrence of MACE-4 as compared to pooled comparators (HR: 0.81; 97.85% CI: 0.52 to 1.26; p=NR).⁴¹

Injectable Semaglutide

The primary outcome of the SUSTAIN-6 CVOT was the first occurrence of MACE-3 (death from cardiovascular causes, nonfatal myocardial infarction, or nonfatal stroke). The pooled doses of semaglutide (0.5 mg and 1 mg) were noninferior and superior to placebo in reducing the occurrence of MACE-3 (HR: 0.74; 95% CI: 0.58 to 0.95; p<0.001 for noninferiority; p=0.02 for superiority). When expanding the primary composite outcome to also include the occurrences of revascularization (coronary or peripheral), and hospitalization for unstable angina or heart failure, semaglutide was noninferior to placebo (HR: 0.74; 95% CI: 0.62 to 0.89; p=.002). The reduction of risk in both composite outcomes was largely driven by semaglutide's ability to reduce the occurrence of nonfatal stroke versus placebo (HR: 0.61; 0.38 to 0.99; p<.04); the differences in risk reduction between semaglutide and placebo were not statistically significant for the remaining components of the composite outcomes: cardiovascular or all-cause death, nonfatal MI, hospitalization for unstable angina or heart failure ([Supplemental Table D4.10](#)).

SUSTAIN-6 included several pre-specified subgroup analyses of interest for the primary outcome of MACE-3: established cardiovascular disease, established chronic heart failure, established moderate-to-severe renal impairment, baseline BMI, race, and ethnicity. None of the subgroups had a statistically significant treatment interaction between semaglutide and placebo ([Supplemental Table D4.11](#)).

There was mixed evidence on semaglutide's impact on microvascular outcomes. When compared to placebo, semaglutide reduced the risk of new or worsening nephropathy (HR, 0.64; 95% CI: 0.46 to 0.88; p=.005) but was associated with an increased the risk of diabetic retinopathy complications (HR: 1.76; 95% CI: 1.11 to 2.78; p=.02), although it is not clear whether semaglutide's effect on retinopathy differs between patients with and without baseline retinopathy.⁴² The incidence of diabetic neuropathy was not reported in SUSTAIN-6.

Data from the SUSTAIN-6 CVOT led to semaglutide's approved indication of reducing the risk of cardiovascular death, nonfatal MI, or nonfatal stroke in adults with T2DM and established cardiovascular disease.⁴³

Empagliflozin

The primary outcome of the EMPA-REG OUTCOME CVOT was the first occurrence of MACE-3. The pooled doses of empagliflozin (10 mg and 25 mg) were noninferior and superior to placebo in reducing the occurrence of MACE-3 (HR: 0.86; 95% CI: 0.74 to 0.99; p<0.001 for noninferiority; p=0.04 for superiority). When expanding the primary composite outcome to also include

hospitalization for unstable angina, empagliflozin was noninferior but not superior to placebo (HR: 0.89; 95% CI: 0.78 to 1.01; $p < 0.001$ for noninferiority; $p = 0.08$ for superiority).

The reduction of risk in both composite outcomes was largely driven by empagliflozin's ability to reduce the occurrence of cardiovascular death by 38% (HR: 0.62; 0.49 to 0.77; $p < .001$); the differences in risk reduction between empagliflozin and placebo were not statistically significant for the remaining components of the composite outcomes ([Supplemental Table D4.10](#)). Patients in the empagliflozin group also had lower risk than placebo from hospitalization for heart failure and death from any cause ([Supplemental Table D4.10](#)).

EMPA-REG OUTCOME included several pre-specified subgroup analyses of interest for the outcomes of MACE-3 and cardiovascular death: established moderate-to-severe renal impairment, baseline BMI, race, and ethnicity (CV death only). For MACE-3, none of the subgroups had a statistically significant treatment interaction between empagliflozin and placebo ([Supplemental Table D4.11](#)).

There was mixed evidence on empagliflozin's impact on microvascular outcomes. There was a statistically significant difference between the empagliflozin and placebo group on the reduced risk of new or worsening nephropathy (HR, 0.61; 95% CI: 0.53 to 0.70; $p < .001$) but not the risk of diabetic retinopathy complications (HR: 0.69; 95% CI: 0.43 to 1.12; $p = .134$). The incidence of diabetic neuropathy was not reported in EMPA-REG OUTCOME.

Data from the EMPA-REG OUTCOME CVOT led to empagliflozin's approval for the indication of reducing the risk of cardiovascular death in adults with T2DM and established cardiovascular disease.⁴⁴

Participants treated with empagliflozin in the EMPA-REG OUTCOME CVOT group experienced statistically significantly lower rates of both acute renal failure ($p < .01$) and acute kidney injury ($p < .05$) than placebo, suggesting a renal protective effect. These promising results have led to further study of empagliflozin's effect on kidney disease progression or CV death in adults with established chronic kidney disease both with and without diabetes (EMPA-KIDNEY), scheduled to conclude in 2022.⁴⁵

Table 3.3. CVOT Key Trial Results

Cardiovascular Outcomes		SUSTAIN-6		EMPA-REG OUTCOME	
		SEM 1 mg	PBO 1 mg	PBO	EMPA (10/25 mg)
Composite Outcome	n (%)	108 (6.6)	146 (8.9)	282 (12.1)	490 (10.5)
	HR (95%CI)	0.74 (0.58 to 0.95)		0.86 (0.74 to 0.99)	
	p-value, NON	<0.001	REF	<.001	REF
	p-value, SUP	0.02	REF	0.04	REF
Expanded Composite Outcome	n (%)	199 (12.1)	264 (16.0)	333 (14.3)	599 (12.8)
	HR (95%CI)	0.74 (0.62 to 0.89)		0.89 (0.78 to 1.01)	
	p-value, NON	0.002	REF	<.001	REF
	p-value, SUP	NA	NA	0.08	REF

CI: confidence interval, EMPA: empagliflozin, HR: hazard ratio, n: number, NA: not applicable, NON: noninferiority, PBO: placebo, REF: reference, SEM: injectable semaglutide, SUP: superiority

Harms

Tirzepatide versus Background Therapy

The most frequent adverse events in the Frias 2018 and 2020 trials for tirzepatide compared with background therapy were gastrointestinal-related. Participants in the tirzepatide 15 mg arm experienced nausea (40%), diarrhea (32%), or vomiting (26%) at greater proportion than placebo ([Supplemental Table D4.6](#)). The discontinuation rate in the 15 mg arm was substantial in the Frias 2018 study (24.5% vs. 3.9% placebo).

Other adverse events in the two trials included hypoglycemia (plasma glucose of ≤ 70 mg/dL) (10-18%), decreased appetite (18-28%), headache (9-21%), abdominal pain (5-18%), dizziness (9-11%), and injection site reaction (6-8%). There were no reported episodes of severe hypoglycemia across the two trials.

Tirzepatide versus Semaglutide

The safety profile of tirzepatide in the 40 week SURPASS-2 trial was consistent with Phase 2 results; the most frequent adverse events were gastrointestinal-related ([Supplemental Table D4.6](#)). A greater proportion of participants in the tirzepatide 15 mg arm had a serious adverse event than in the semaglutide arm (5.7% vs. 2.8%). Adverse events resulted in the discontinuation of the active treatment in 8.5% (15 mg tirzepatide) and 4.1% (1 mg semaglutide) of participants.

Injection-site reactions (4.5% vs. 0.2%) and hypoglycemia (1.7% vs. 0.4%) were more frequent in the tirzepatide 15 mg arm compared with semaglutide. Both drugs had similar rates of adjudicated pancreatitis, hypersensitivity, and cholelithiasis ([Supplemental Table D4.6](#)). There were no cases of

diabetic retinopathy among the 15 mg tirzepatide or 1 mg semaglutide arms, with two cases (0.4%) in the 10 mg tirzepatide arm.

Tirzepatide versus Empagliflozin

There were no available direct or indirect comparisons of safety outcomes between tirzepatide and empagliflozin.

Subgroup Analyses and Heterogeneity

Established ASCVD

Currently, there is no available evidence to evaluate the clinical efficacy and safety of tirzepatide for this subgroup. The SUSTAIN CVOT trial is ongoing and expected to be completed in 2024.

Established CHF

Currently, there is no available evidence to evaluate the clinical efficacy and safety of tirzepatide for this subgroup.

Established moderate-to-severe renal impairment

In the 40 week SURPASS-2 trial, participants receiving tirzepatide 15 mg achieved a greater reduction in HbA1c from baseline than semaglutide, irrespective of renal impairment (eGFR <60, ≥60 mL/min per 1.73 m²).³⁸(data on file) Participants with established moderate-to-severe renal impairment (eGFR <60 mL/min per 1.73 m²) experienced a smaller reduction in HbA1c from baseline in both trial arms.³⁸(data on file)

Obesity

In the 40 week SURPASS-2 trial, participants receiving tirzepatide 15 mg achieved a greater reduction in HbA1c from baseline than semaglutide, irrespective of obesity status (BMI categories <30 kg/m², ≥30 to <35 kg/m², ≥35 kg/m²).³⁸(data on file)

Race and ethnicity or socioeconomic status

In the 40 week SURPASS-2 trial, Hispanic/Latino participants experienced a greater reduction in HbA1c from baseline in all trial arms compared with non-Hispanic/Latino participants, with the pattern of relative HbA1c decline between groups consistent with the overall trial results (i.e., greater reduction in HbA1c in the tirzepatide arms compared with semaglutide) ³⁸ (data on file)

Heterogeneity

Table 3.2 outlines the baseline characteristics across the five trials used in our NMA. There were no notable differences between studies in the distribution of baseline characteristics of age, sex, race, ethnicity, HbA1c, weight, BMI, and background use of metformin. We were unable to assess baseline history of microvascular or macrovascular events. Participants in the SUSTAIN-2 and PIONEER-3 trials received additional background therapy via thiazolidinediones and sulfonylurea, respectively.

Uncertainty and Controversies

For the newer antihyperglycemic drugs, particularly GLP-1 Ras and SGLT-2 inhibitors, the glucose-lowering effect of the drug is only one facet of its overall clinical value. The importance of other outcomes such as weight loss, prevention of cardiovascular events, and renal protection have gained prominence, as reflected both by separate indications for such benefits and in clinical practice guidelines favoring GLP-1 Ras and SGLT-2 inhibitors for certain groups of patients with T2DM.⁴⁶ Thus, the value assessment of new T2DM therapies includes examining data beyond glucose lowering and glycemic control.

Tirzepatide shows an impressive impact on glucose-lowering and glycemic control, with an average HbA1c lowering of around 2% compared with background therapy, 0.45% in direct comparison to semaglutide and 1.1% in indirect comparison to empagliflozin. Additionally, a substantial number of patients taking 15 mg of tirzepatide achieved near-normal glycemic levels, which is both important to patients and may have benefits in terms of slowing progression of disease.²⁰

Tirzepatide is also associated with more weight loss than its comparators. Given that the majority of patients with T2DM are overweight or obese and that nearly half of patients with T2DM are inadequately controlled on current therapy, these results are promising. However, gastrointestinal side effects were frequent with tirzepatide, and this may affect real-world acceptance and adherence of the drug. Furthermore, GLP-1 Ras may be associated with an increased risk of pancreatitis, acute kidney injury, and thyroid cancer, and GIP inhibition is a new mechanism of action so long-term side effects of this type of drug are unknown.⁴⁷ Finally, the durability of tirzepatide's effect on glycemic control and other parameters such as weight loss, are not yet known. Since there are already multiple effective options for treatment of T2DM, the unknown harms of tirzepatide may influence treatment decision-making by clinicians and patients.

The data on cardiovascular and renal outcomes for tirzepatide is currently immature, and thus it is not yet clear whether tirzepatide improves cardiovascular and renal outcomes, as has been demonstrated by its comparators. It is reassuring that tirzepatide did not show any cardiovascular harm in a safety meta-analysis across the SURPASS trials and encouraging that it had a greater impact on surrogate cardiovascular outcomes such as blood pressure and lipids in comparison to semaglutide and empagliflozin and a trend towards cardiovascular benefit overall in SURPASS-4.

However, SURPASS-4 was only powered to demonstrate cardiovascular safety, and the formal CVOT to provide more definitive data on cardiovascular outcomes is ongoing. Additionally, although GLP-1 receptor agonism is part of the mechanism for tirzepatide, the drug has an additional mechanism of GIP inhibition, which may have some direct effects in the myocardium. There are mixed data from pre-clinical, clinical, and epidemiological studies on whether GIP inhibition is helpful or harmful with regard to protection from cardiovascular events.⁴⁸⁻⁵¹ Finally, cardiovascular benefit is not uniform across the GLP-1 RA class.⁵² Therefore, we hesitate to fully extrapolate the cardiovascular benefits of GLP-1 Ras like semaglutide to tirzepatide, and eagerly await the results from tirzepatide's CVOT.


The lack of head-to-head comparison between tirzepatide and empagliflozin made our assessment of the net benefit of tirzepatide more challenging. Since tirzepatide and empagliflozin have completely different mechanisms of action, without a direct comparison, it is difficult to judge whether tirzepatide may represent a substantial improvement over empagliflozin, particularly in patients with established or at high risk of ASCVD, CKD, or heart failure; three common co-morbid conditions. However, tirzepatide did show statistically significant and likely clinically significant improvements in HbA1c, weight loss, SBP and lipid parameters, though we have only moderate certainty about the results from the indirect comparison through the NMA.

Finally, T2DM is a disease that disproportionately affects minority populations. However, trials of antihyperglycemic drugs including tirzepatide do not reflect the demographics of the disease in the US. The lack of inclusion of minority populations in the clinical trials poses a serious issue in terms of evaluating whether new treatments may increase or decrease health inequities. Furthermore, as access to newer, potentially more expensive drugs tends to be limited at least during initial launch, missing potential differences in population subgroups may prevent populations who may derive greater benefit from the drug from being able to access it. More diverse representation in clinical trials can also provide more precise data to help match drugs with the patients who have the most favorable benefit/side effect profile, which may in turn improve adherence and clinical outcomes of treatment.

3.3. Summary and Comment

An explanation of the ICER Evidence Rating Matrix (Figure 3.1) is provided in the Supplement (Section D).

Figure 3.1. ICER Evidence Rating Matrix



A = "Superior" - High certainty of a substantial (moderate-large) net health benefit
B = "Incremental" - High certainty of a small net health benefit
C = "Comparable" - High certainty of a comparable net health benefit
D = "Negative" - High certainty of an inferior net health benefit
B+ = "Incremental or Better" - Moderate certainty of a small or substantial net health benefit, with high certainty of at least a small net health benefit
C+ = "Comparable or Incremental" - Moderate certainty of a comparable or small net health benefit, with high certainty of at least a comparable net health benefit
C- = "Comparable or Inferior" - Moderate certainty that the net health benefit is either comparable or inferior with high certainty of at best a comparable net health benefit
C++ = "Comparable or Better" - Moderate certainty of a comparable, small, or substantial net health benefit, with high certainty of at least a comparable net health benefit
P/I = "Promising but Inconclusive" - Moderate certainty of a small or substantial net health benefit, small (but nonzero) likelihood of a negative net health benefit
I = "Insufficient" - Any situation in which the level of certainty in the evidence is low

In this review, we set out to compare the clinical efficacy and safety of tirzepatide to background therapy, injectable semaglutide and empagliflozin.

For the comparison between tirzepatide and background therapy, we evaluated data from two Phase 2 trials on change in glucose levels, weight, lipid profiles, and blood pressure, as well as indirect comparisons on glycemic control, weight loss, LDL, and SBP via results from a five trial

NMA. Compared to background therapy, tirzepatide treatment provided consistent, substantial, and clinically meaningful reductions in HbA1c and weight across the two trials; results for systolic blood pressure and LDL are more modest. The NMA results support the superiority of tirzepatide over background therapy, with statistically significant treatment differences in HbA1c, weight, LDL and SBP. The impact of tirzepatide on cardiovascular outcomes has yet to be determined; however, SURPASS-4 provides a glimpse into the positive direction of tirzepatide's cardiovascular safety and potential benefit (in comparison with insulin). Finally, although there was a relatively high discontinuation rate due to gastrointestinal side effects seen in the Phase 2 trial, dose titration appears to mitigate these effects, and severe adverse events are rare. Therefore, we have high certainty that tirzepatide provides a substantial net benefit compared with background therapy for glycemic control and weight loss. We judge tirzepatide to be incremental or better ("B+") for this comparison.

For the comparison between tirzepatide and injectable semaglutide, we relied on head-to-head evidence from a Phase 3 trial (SURPASS-2) to evaluate change from baseline in the intermediate outcomes of glucose levels, weight, lipid profiles, blood pressure, and quality-of-life. Tirzepatide demonstrated a small estimated net benefit in reducing HbA1c and substantial net benefit in weight loss when compared to semaglutide. The drugs appear to be comparable in terms of effects on lipid profile and blood pressure. However, while semaglutide has been demonstrated to improve cardiovascular outcomes and has an indication for this purpose, these data are currently lacking for tirzepatide. We hesitate to fully extrapolate the favorable cardiovascular outcomes data for GLP-1 Ras to tirzepatide, as cardiovascular benefit is not uniform across the GLP-1 RA class, and because of the dual GLP-1/GIP mechanism of action. In consideration of tirzepatide's superiority over semaglutide in outcomes of glycemic control and weight loss, similar safety profiles but current lack of definitive cardiovascular outcome data and the uncertainty about the impact of the GIP moiety, we judge tirzepatide to be comparable or incremental ("C+") for this comparison, with a high certainty of at least a comparable net benefit.

For the comparison between tirzepatide and empagliflozin, tirzepatide provided a substantial net benefit versus empagliflozin via changes in HbA1c, weight, LDL, and SBP. However, with no direct head-to-head trials, we were limited to making an indirect comparison via NMA and thus there is more uncertainty in the precision of these estimates. Additionally, empagliflozin has established cardiovascular and renal benefits in adults with T2DM; data on such outcomes is less certain for tirzepatide, though early signals for the cardiovascular benefit of the drug are promising. Thus, tirzepatide may provide anywhere from comparable net health benefits (if tirzepatide shows equivalent cardiovascular and renal benefits to empagliflozin) to substantial net health benefits (if the improvements in intermediate outcomes over empagliflozin translate into larger gains in cardiovascular and renal outcomes). Considering the benefits seen for tirzepatide over empagliflozin for intermediate outcomes as well as the current lack of definitive cardiovascular and

renal outcomes for tirzepatide, we judge that tirzepatide is comparable or better (C++) for this comparison.

Table 3.4. Evidence Ratings

Treatment	Comparator	Evidence Rating
Tirzepatide	Background Therapy	B+
Tirzepatide	Injectable Semaglutide	C+
Tirzepatide	Empagliflozin	C++

4. Long-Term Cost Effectiveness

4.1. Methods Overview

We used a patient-level microsimulation relying on the UKPDS-OM2,¹¹ which was an adaptation of a published microsimulation⁵³ and an update of the 2019 ICER report on diabetes therapies.^{40,54} This particular modeling approach was selected in large part due to the complexity of co-occurring comorbidities in people with T2DM. Our [model analysis plan](#) indicated use of the BRAVO risk engine,⁵⁵ however feasibility testing after the publication of the model analysis plan suggested that the information available on BRAVO within the public domain was insufficient to fully implement that engine in a microsimulation model.⁵⁵ We therefore used the same risk engine as described in the 2019 ICER report evaluating oral semaglutide, as we deemed this risk engine as the best publicly available source.^{40,54}

Consistent with the Comparative Clinical Effectiveness assessment, the intervention of interest was tirzepatide plus background therapy versus (1) injectable semaglutide plus background therapy, (2) empagliflozin plus background therapy, and (3) background therapy alone. For consideration of the equal-value of life years (evLYs) gained measure and the purpose of estimating Health Benefit Price Benchmarks, we identified injectable semaglutide plus background therapy as a primary comparator for this version of the report. We chose injectable semaglutide plus background therapy as the primary comparator because of: (1) mechanism of action overlap between tirzepatide and injectable semaglutide, (2) public and clinical expert feedback that suggested injectable semaglutide is an appropriate comparator that would be considered as an alternative to tirzepatide (rather than added on to tirzepatide), (3) availability of a randomized head-to-head trial on intermediate outcomes, and (4) prior research that suggests injectable semaglutide is cost-effective.⁵⁶⁻⁵⁸

The model was informed by clinical trials, the ICER NMA of relevant clinical trials, quality-of-life literature, and validation versus other prior economic models.^{54,59,60} The base-case analysis took a health care sector perspective and thus focused on direct medical care costs only. Costs and outcomes were discounted at 3% per year.¹² Because long-term cardiovascular outcomes trial data are immature for the primary intervention under examination in this review, tirzepatide, health benefits for all modeled therapies were informed by intermediate outcomes: HbA1c, body weight, LDL, and SBP, that are predictors in the UKPDS-OM2 risk engine.¹¹ Because intermediate outcomes alone may not capture a treatment's potential reduction in CVO event risks, as predicted by the UKPDS-OM2 equations, we further multiplied the predicted event risks by the corresponding trial-based CVO hazard ratios and their respective confidence intervals.

The model (Figure 4.1) is an individual patient-level, Monte Carlo-based microsimulation of costs, quality of life, clinical events, and mortality associated with T2DM among US adults with the

disease. Two modeling steps were used: (1) event microsimulation and (2) calculation of mean results from the pool of simulated patients' lifetime outcomes. Patients, with data from multiple NHANES surveys, were separately simulated through the modeling steps for tirzepatide and each comparator (each added to current ongoing background antihyperglycemic therapy), plus background therapy alone. The two model steps are explained below:


- (1) Event microsimulation.** Each NHANES patient was sequentially run through the event microsimulation. Each model cycle was one year in duration. The UKPDS-OM2 risk equations were used to calculate the incidence of any clinical event(s) and/or mortality in each year until the simulated patient died. Effects of each included therapy, such as change in HbA1c after the first cycle, were included depending on data availability from the NMA. All event and/or mortality associated costs and health state utility weights were applied concurrently. The UKPDS-OM2 risk equations accounted for patient history upon entering the model as well as new clinical events that occurred during the microsimulation; for example, a patient who experienced a first MI in a given year of the microsimulation then had the history of MI covariate turned on in each subsequent year.
- (2) Calculation of mean results.** After each simulated patient died, the model recorded the patient's lifetime cost, quality-adjusted life years (QALYs), equal-value of life years (evLYs), life years, and clinical events. Each outcome was then averaged over the entire pool of patients to derive overall model results.

Further details on the economic modeling methods used are available in [Supplement E](#).

Based on public feedback, the following changes were made to the draft report:

1. Patients whose HbA1c levels exceeded 8.5% added insulin to modeled active therapies rather than replacing active therapies to align with the 2019 T2DM report and reflect common clinical practice.
2. The MACE hazard ratio and uncertainty estimates applied to tirzepatide in the base case were derived from SURPASS-4 results, while the former base case (tirzepatide MACE hazard ratio equal to 1.0) was included as a scenario analysis.
3. 2- and 3-year modeled outcomes compared to existing cardiovascular outcome trials for semaglutide and empagliflozin were added for external validation (Supplement Table E6).
4. Additional scenario analyses were added to test model assumptions, including those around weight loss utility and risk factor progression.
5. Additional model outcomes were added to the supplement, including disaggregated and undiscounted outcomes, for increased model transparency.
6. Point of clarification: Risk factor progression was reflected in the model first using trial based NMA results and then using published equations for HbA1c and weight change progression for diabetes patients on therapy. The assumption of risk factor progression was then tested in a scenario analysis.

Figure 4.1. Model Structure



4.2. Key Model Assumptions and Inputs

Below is a list of key model choices:

- Long-term survival and the incidence of diabetes-related clinical events were modeled by applying the UKPDS-OM2 risk equations to a US patient population in an event microsimulation, adjusting the risk equation outputs using available long-term outcome data.¹¹ The UKPDS-OM2 equations were created from a population of T2DM patients in the United Kingdom and may not reflect US T2DM patient demographics, including differences in race and differences in risk of T2DM-related events.
- Major adverse cardiovascular, heart failure, and nephropathy event rates for injectable semaglutide and empagliflozin were adjusted using data (HRs) from their respective cardiovascular outcomes trials, and cardiovascular event rates for tirzepatide were also adjusted using the HR point estimate and its uncertainty based on the results of SURPASS-4 across pooled tirzepatide doses.^{8,9,35} Tirzepatide was not adjusted for heart failure and nephropathy event rates in the base case, reflecting its currently immature CV and renal outcomes and its lack of in-class comparators to use as proxies.
- Quality of life was modeled within QALYs using projected patient survival weighted using regression-based disutilities for each diabetes-related complication experienced in each model cycle.
- The model included all treatment costs associated with each individual drug regimen, including drug acquisition costs, downstream treatments such as insulin, and supportive care costs (e.g., clinician visits and self-monitoring), as well as costs associated with diabetes-related complications experienced in each model cycle.
- All model outcomes were calculated over a lifetime time horizon.¹²
- Life-years, QALYs, and health care cost outcomes were discounted at 3% per year.¹²

Our model included several assumptions stated below.

Table 4.1. Key Model Assumptions

Assumption	Rationale
Patient diabetes outcomes were simulated using the UKPDS OM-2 equations; predicted model outcomes for MACE, HF, and nephropathy were adjusted by hazard ratios to align with existing trial evidence where possible.	Accurately modeling the complexity of diabetes disease progression is notoriously difficult. The UKPDS equations remained the best available tool for this modeling exercise despite the known limitations when applied to a modern US population. These limitations were addressed in multiple ways, including hazard ratio adjustments using available clinical evidence and short-term external validation.
Initial patient clinical characteristics were based on NHANES survey data.	NHANES is a nationally representative and federally funded survey repeated every two years in the United States, uniquely capturing the wide range of demographic and clinical data necessary to understand and predict diabetes disease progression in a modern population.
Modeled risk equation adjustment for CV and renal outcomes for active comparators with CVOT data is maintained while patients remain on treatment.	Active treatment comparators (injectable semaglutide and empagliflozin) have data from CVOTs. We adjusted the event prediction output of the risk equations based on how the model outcomes compare to those trial outcomes in the base-case analysis.
Modeled risk equation adjustment for CV and renal outcomes for tirzepatide is maintained while patients remain on treatment.	Long-term effectiveness of tirzepatide is currently unknown. We evaluated the impact of not applying any adjustment of relative effect to tirzepatide for changes in heart failure and renal outcomes in the base case. Tirzepatide's MACE outcomes were adjusted based on SURPASS-4 findings in the base case.
Ongoing background therapy was assumed the same for all comparators, and all patients in all model arms who discontinue treatment (including those on background therapy alone) transition to insulin.	The goal is to evaluate direct comparisons among the treatments of interest and not multiple possible treatment sequences.
Patients discontinued their add-on treatment at a rate of 9.1% in the second model cycle. The discontinuation rate was derived from the EMPA-REG EXTEND trial, the only trial to present discontinuation data contingent on a successful initial treatment period. Those who were on add-on treatment after the second model cycle were assumed to remain on add-on treatment and received the corresponding treatment benefits, risks, and costs for their remaining life. Patients whose HbA1c reached 8.5% or above during any model cycle after the first were assumed to initiate insulin as add-on treatment.	<p>Equivalent discontinuation was assumed for active treatments after the initial model cycle as this measure was not included in the NMA and we had insufficient evidence to suggest differential long-term discontinuation across all therapies.</p> <p>We evaluated changes to the 8.5% HbA1c threshold for adding insulin therapy in scenario analyses.</p>

The patient population for this analysis was derived from the National Health and Nutrition Examination Survey (NHANES), conducted by the United States Centers for Disease Control and Prevention. Survey participants from years 2013-2014, 2015-2016, and 2017-2018 were included if they had self-reported diabetes, HbA1c levels greater than 7%, and were already taking metformin (with or without sulfonylureas), but not another type of add-on diabetes therapy. This resulted in 387 unique patients, whose characteristics are described in Table 4.2, and whose patient-level baseline clinical inputs were used to power the microsimulation model.

Table 4.2. Base-Case Model Cohort Characteristics

Description	Mean (SD) or Percentage (N)
<u>Patient Characteristics</u>	
Age at Time of survey	60.4 (11.53)
% Female	42.9% (166)
Duration of disease	9.9 (7.91)
Race	26.6% (103)
White, %	22.7% (88)
Black or African American, %	14.5% (56)
Asian, %	11.4% (44)
Hispanic, %	24.8% (96)
Other	21.2% (82)
Weight (kg)	88.8 (24.94)
BMI (kg/m ²)	31.9 (7.55)
HbA1c (%)	8.4 (1.54)
LDL (mmol/L)	2.7 (0.93)
SBP (mmHg)	132.5 (18.88)
% Current smokers	36.7% (142)
% on Metformin	100.0% (387)
% on Sulfonylurea	42.9% (166)
<u>Disease History</u>	
Myocardial infarction	5.9% (23)
Stroke	4.4% (17)
Heart failure	4.1% (16)
Ischemic heart disease	7.8% (30)
Angina	4.1% (16)
Renal disease	15.8% (61)

HbA1c: hemoglobin A1c, LDL: low-density lipids, N: number of subjects in full analysis set, SBP: systolic blood pressure, SD: standard deviation

The efficacy of tirzepatide, injectable semaglutide, and empagliflozin as compared to background therapy alone were measured via intermediate outcomes such as changes in HbA1c, lipid levels, blood pressure, and body weight. Intermediate outcome efficacy estimates for each active comparator against background therapy were derived from the NMA described in Chapter 3 and were applied after the first cycle of the model. Each model cycle subsequently utilized updated patient-level input parameters based on the predictions of the UKPDS-OM2 risk equations and

patient history. UKPDS equations, which are available in the public domain, estimate the probability of 12 diabetes complication events and four mortality events in each cycle.¹¹ To supplement UKPDS-OM2 output, time-varying risk factor values of HbA1c and weight were calculated using additional published equations.⁶¹ To adjust the UKPDS event predictions to existing trial data, we multiplied the probability of events occurring predicted by UKPDS equations by the relevant HR in each model cycle for three event measurements: composite MACE, heart failure, and renal events. Thus, the probabilities informed by the regression equations were adjusted by the HRs. These HRs were abstracted directly from the semaglutide and empagliflozin CVOTs, and from the SURPASS-4 trial in the case of tirzepatide.

Health state utilities and mortality risk equations are further detailed in [Supplement E2](#).

Calculated net drug prices were applied to each patient while they remained on treatment. For each comparator, we obtained net pricing estimates from SSR Health, LLC, which combines data on unit sales with publicly-disclosed US sales figures that are net of discounts, rebates, concessions to wholesalers and distributors, and patient assistance programs, to derive a net price.⁶² We estimated net prices by comparing the most recent four-quarter averages (i.e., third quarter of 2020 through second quarter of 2021) of both net prices and wholesale acquisition cost (WAC) per unit to arrive at a mean discount from WAC for the drug. Finally, we applied this average discount to the most recent available WAC (accessed September 2021) to arrive at an estimated net price per unit. Tirzepatide was given a placeholder price equal to semaglutide, the only suitable proxy drug in the injectable GLP drug class, until its price is publicly available. Drug cost inputs can be found in Table 4.3.

Table 4.3. Drug Cost Inputs

Drug	WAC per 30-Pill Bottle/Pen	Net Price Per 30-Pill Bottle/Pen	Discount From WAC	Net Price per Year‡
Tirzepatide*	(4 weekly doses)	(4 weekly doses)	-	Placeholder \$4,643.50
Semaglutide (Ozempic®) 4 mg/3 mL pen†	\$851.60 (4 weekly doses)	\$355.97 (4 weekly doses)	58.20%	\$4,643.50
Empagliflozin (Jardiance®) 30-tablet bottle§	\$548.54	\$107.51	80.40%	\$1,402.43
Metformin#	\$0.83	-	-	\$10.04
Sulfonylureas‡	\$5.05	-	-	\$61.48

WAC: wholesale acquisition cost

*As a placeholder, we used the net price of Ozempic® (semaglutide), which is a once weekly injectable GLP-1; WAC pricing and discounts reflect the number of pen doses and quantity of pens necessary for Ozempic® use.

†The 4 mg/3 mL Ozempic® pen includes four 1 mg doses; assumes 1 mg weekly dose.

‡One year = 365.25 days or 52 weeks

§Assumes 25 mg daily dose of empagliflozin. Source: Red Book.

#Assumes 1000 mg daily dose of metformin. Source: Red Book.

‡Assumes 20 mg daily dose of glipizide. Source: Red Book.

The average clinical costs of experiencing a diabetes-related event within a cycle, as well as the average annual clinical costs of subsequent years after a health event, were adapted from the literature.^{53,63,64} Patients were able to experience events concurrently within the model. Costs were inflated to 2021 USD. Additional details on the economic model can be found in [Supplement E](#).

4.3. Results

Base-Case Results

Due to the characteristics of the microsimulation, the model's base-case results are presented as the mean and 95% credible range estimates from probabilistic sensitivity analyses, by jointly varying all model parameters and UKPDS-OM2 risk engine equation coefficients over 3,000 simulations per patient, then calculating 95% credible range estimates for each model outcome based on the distributions of those simulations. The base-case discounted results are presented in Table 4.4, with incremental comparisons of tirzepatide to the three comparators presented in Table 4.5. Equal value of life years gained were not reported given tirzepatide average life years were not greater than injectable semaglutide average life years. Disaggregated base-case results, including costs and disease-related events, are available in the [Supplement Table E3.1](#) and undiscounted results are available in [Supplement Table E3.2](#).

Table 4.4. Results for the Base Case for Tirzepatide and Comparators: Injectable Semaglutide, Empagliflozin, and Background Therapy – Mean (95% Credible Range)*

Treatment	Add-On Drug Cost		Total Cost (including background therapy and insulin)		QALYs		Life-years‡	
	Mean	95% Credible Range	Mean	95% Credible Range	Mean	95% Credible Range	Mean	95% Credible Range
Tirzepatide†	\$40,500	(\$38,200 - \$42,900)	\$306,200	(\$275,100 - \$338,600)	4.90	(4.68 – 5.12)	9.36	(8.91 – 9.83)
Injectable Semaglutide	\$41,200	(\$38,800 - \$43,500)	\$309,200	(\$280,000 - \$339,400)	4.85	(4.64 – 5.05)	9.53	(9.08 – 9.97)
Empagliflozin	\$12,000	(\$11,300 - \$12,700)	\$275,700	(\$247,600 - \$304,600)	4.60	(4.40 – 4.79)	9.17	(8.73 – 9.61)
Background Therapy	\$0	NA	\$261,800	(\$234,500 - \$290,800)	4.13	(3.95 – 4.33)	8.34	(7.93 – 8.77)

QALY: quality-adjusted life-year

*All costs and outcomes discounted at 3% annually

†Using a Placeholder Price equal to the net price of semaglutide

‡Equal value of life years gained were not reported given tirzepatide average life years were not greater than injectable semaglutide average life years.

Of the known drug costs, injectable semaglutide had the highest lifetime drug costs at \$41,200 (a placeholder cost was used for tirzepatide). For total lifetime costs, including background treatment costs and costs of cardiovascular and renal complications, injectable semaglutide had the highest costs at approximately \$309,200, followed by tirzepatide (\$306,200), empagliflozin (\$275,700), and background therapy (\$261,800). Tirzepatide was estimated to produce the highest QALYs of all considered therapies, however the QALY 95% credible ranges for active comparators overlapped. All active comparators produced higher QALYs than background therapy alone without overlapping credible ranges.

Table 4.5. Incremental Cost-Effectiveness Ratios for the Base Case – Mean (95% Credible Range)[§]

Treatment	Comparator	Cost per QALY Gained		Cost per Life Year Gained‡	
		Mean	95% Credible Range	Mean	95% Credible Range
Tirzepatide*	Injectable Semaglutide	Less Costly, More Effective†	(-\$1,546,000 - \$1,384,000)	\$17,000	(-\$709,100 - \$633,700)
Tirzepatide*	Empagliflozin	\$101,000	(-\$54,800 - \$331,100)	\$160,000	(-\$950,700 - \$1,265,100)
Tirzepatide*	Background Therapy Alone	\$58,000	(\$10,900 - \$99,100)	\$44,000	(\$10,400 - \$82,500)

QALY: quality-adjusted life-year

*Using a Placeholder Price equal to the net price of semaglutide

†Although the mean ICER point value indicates lower cost and greater health benefit of tirzepatide over semaglutide, neither of the ratio's inputs (savings of \$2,900 for an additional 0.05 QALYs) were statistically different from zero

‡Equal value of life years gained were not reported given tirzepatide average life years were not greater than injectable semaglutide average life years.

§Ratios and credible ranges displayed in table are the mean values across all microsimulations rather than derivations of output in table 4.4


The incremental cost-effectiveness measures reported in Table 4.5 are based on a placeholder price equal to the net value of injectable semaglutide and should be interpreted with caution.

Sensitivity Analyses

One-way sensitivity analyses (OWSA) to identify the impact of parameter uncertainty and key drivers of model outcomes were performed. Details about the sensitivity analysis methods can be found in Supplement E4. Compared to injectable semaglutide, the inputs that were associated with the largest variation in incremental costs were annual costs of semaglutide and tirzepatide, semaglutide and tirzepatide event hazard ratios, and the HbA1c discontinuation threshold (Figure 4.2). The reader should only use these one-way sensitivity analysis findings to gain insights into the general magnitude of impact that unique model inputs have on incremental costs (and health gains). The base-case incremental results remain a more robust average estimate compared with

the central values displayed in the one-way sensitivity figures given the added computer simulation requirements needed for the sensitivity analyses. The tornado diagram for incremental costs was truncated to include inputs with at least a \$500 difference between the high and low value. Compared to injectable semaglutide, the inputs that were associated with the largest variation in incremental QALYs were the tirzepatide and semaglutide event hazard ratios, the HbA1c discontinuation threshold, and semaglutide and tirzepatide effect on body weight (Figure 4.3). The tornado diagram for incremental QALYs was truncated to include inputs with more than a 0.01 QALY difference between the high and low value. Additional tornado diagrams are presented in [Supplement E4](#).


Figure 4.2. Incremental Costs Tornado Diagram (Tirzepatide vs. Injectable Semaglutide)*



CHF: congestive heart failure, HbA1c: Hemoglobin A1c/glycosylated hemoglobin, HR: hazard ratio, MACE: major adverse cardiovascular event, Tx: treatment

*Using a placeholder price for tirzepatide

Figure 4.3. Incremental QALYs Tornado Diagram (Tirzepatide vs. Injectable Semaglutide)



CHF: congestive heart failure, HbA1c: Hemoglobin A1c/glycosylated hemoglobin, HR: hazard ratio, MACE: major adverse cardiovascular event, Tx: treatment

Unlike a typical cohort model, our probabilistic sensitivity analysis for this microsimulation is presented as the base-case results in order to account for patient and risk equation uncertainty. The cost-effectiveness acceptability curves from the base case are presented in [Supplement Figure E5](#).

Scenario Analyses

Multiple scenario analyses can be found in [Supplement E5](#). We sought to produce model estimates using a modified societal perspective and found a paucity of data to inform such calculations for a simulation. However, we present a calculated estimate of cost-effectiveness under a modified societal perspective using an estimate of productivity costs saved based on assumptions that fill in missing data. The results of these calculations are also presented in the Supplement.

Threshold Analyses

The annual drug costs at which tirzepatide plus background therapy would reach cost-effectiveness thresholds ranging from \$50,000 to \$200,000 per QALY gained compared to semaglutide plus background therapy, empagliflozin plus background therapy, and background therapy alone, are presented below in Table 4.6. Equal value of life years gained were not reported given tirzepatide average life years were not greater than injectable semaglutide average life years.

Table 4.6. Cost per Outcome Threshold Analysis Results for Tirzepatide vs. Background Therapy

	Net Price per Unit	Annual Price to Achieve \$50,000 per outcome	Annual Price to Achieve \$100,000 per outcome	Annual Price to Achieve \$150,000 per outcome	Annual Price to Achieve \$200,000 per outcome
Tirzepatide plus Background Therapy vs. Semaglutide plus Background Therapy					
Mean QALYs Gained	To be determined	\$5,200	\$5,500	\$5,700	\$6,000
Tirzepatide plus Background Therapy vs. Empagliflozin plus Background Therapy					
Mean QALYs Gained	To be determined	\$3,000	\$4,600	\$6,200	\$7,700
Tirzepatide plus Background Therapy vs. Background Therapy Alone					
Mean QALYs Gained	To be determined	\$4,000	\$8,000	\$12,000	\$15,900

QALY: quality-adjusted life-year

Net price for tirzepatide has not been publicly stated at the time of this report; Equal value of life years gained were not reported given tirzepatide average life years were not greater than injectable semaglutide average life years.

Model Validation

We used several approaches to validate the microsimulation model's output. First, we provided the preliminary methods to manufacturers, patient groups, and clinical experts. Based on feedback from these groups, we refined the model approach (outlined in Section 4.1). Second, we varied the model input parameters to evaluate the face validity of changes to those inputs on the results in the OWSA. We also performed model verification for model calculations using internal reviewers. As part of ICER's efforts in acknowledging modeling transparency, we also shared the model with the manufacturers for external verification following the publication of the draft report.

For further model validation and calibration, we compared the cardiovascular and renal trial outcomes versus model projections in the revised report, found in supplement section E6. This comparison was informed by microsimulations on modified patient populations and for a shorter time horizon to reflect specific trial conditions more accurately.

Uncertainty and Controversies

This model represents a simplified version of a complex disease and should therefore be interpreted with its limitations in mind. Cost and long-term outcome data for tirzepatide were unavailable at the time of the report and could not be used to validate its clinical results; any economic output for tirzepatide is based on a placeholder price and the results should not be interpreted as factual. Additionally, there are noted limitations to the UKPDS-OM2 risk equations used for this analysis, including that they were developed based on a patient cohort from the United Kingdom in decades past, had a notably different demographic and clinical patient population than generally seen in the current United States, and may not fully capture the impact of weight loss on CV and renal outcomes. Our choice to use UKPDS-OM2 risk equations was made based on lack of feasible alternatives at the time, and adjustments were used in an attempt to ensure the model estimates aligns with clinical evidence. We present shorter time horizon simulations, aligned with the cardiovascular and renal outcomes trials for semaglutide and empagliflozin, in the supplement in order to compare our model event outcomes with those found in the trials to understand the chosen model's impact on our estimates.

Finally, we acknowledge challenges in modeling tirzepatide given the immaturity of the evidence of its impact on micro- and macrovascular outcomes. In the threshold analyses, we compared tirzepatide to injectable semaglutide and estimated the health gains of tirzepatide through changes in intermediate outcomes, assuming the same hazard ratio point estimate as injectable semaglutide for composite MACE events and no adjustment to other outcomes to reflect tirzepatide's cardiovascular safety study. The scenario analysis that includes tirzepatide's changes only through intermediate outcomes, with no adjustment from hazard ratios, provides estimates of the potential relative impact on health outcomes if the trends observed in tirzepatide's cardiovascular safety study are not sustained long-term.

4.4 Summary and Comment

We used a patient-level microsimulation of T2DM patients taking an add-on therapy over a lifetime time horizon to assess the cost-effectiveness of the novel injectable agent tirzepatide in addition to background therapy against injectable semaglutide plus background therapy, empagliflozin plus background therapy, and background therapy alone. Tirzepatide had the highest average lifetime discounted QALYs of all considered therapies, however the QALY 95% credible ranges for active comparators overlapped. Using a placeholder price equal to injectable semaglutide, the incremental costs per QALY gained for tirzepatide were around or under \$100,000 versus all comparators with mean differences in health gains and costs being smallest in comparison to injectable semaglutide. Uncertainty analyses suggested a wide range of plausible cost-effectiveness estimates for tirzepatide. These results paired with tirzepatide's unknown price preclude strong conclusions about its cost-effectiveness at this time.

5. Contextual Considerations and Potential Other Benefits

Our reviews seek to provide information on potential other benefits offered by the intervention to the individual patient, caregivers, the delivery system, other patients, or the public that was not available in the evidence base nor could be adequately estimated within the cost-effectiveness model. These elements are listed in the table below, with related information gathered from patients and other stakeholders. Following the public deliberation on this report the appraisal committee will vote on the degree to which each of these factors should affect overall judgments of long-term value for money of the intervention(s) in this review.

Table 5.1. Contextual Considerations

Contextual Consideration	Relevant Information
Acuity of need for treatment of individual patients based on the severity of the condition being treated	Although there are multiple medication options for the treatment of T2DM, nearly half of patients have not reached adequate glycemic control. ¹ Thus, additional options for treatment are beneficial.
Magnitude of the lifetime impact on individual patients of the condition being treated	Patients with inadequately controlled T2DM are at risk for microvascular and macrovascular complications, which can substantially affect both quality of life and longevity. Additionally, many patients with T2DM are overweight or obese, and weight loss may be of benefit to prevent complications of obesity.
Other (as relevant)	NA

Table 5.2. Potential Other Benefits or Disadvantages

Potential Other Benefit or Disadvantage	Relevant Information
Patients' ability to achieve major life goals related to education, work, or family life	Prevention or delay of microvascular and macrovascular complications may allow for greater work or educational productivity.
Caregivers' quality of life and/or ability to achieve major life goals related to education, work, or family life	Prevention or delay of microvascular and macrovascular complications may lessen caregiving burden over the lifetime.
Patients' ability to manage and sustain treatment given the complexity of regimen	The medication delivery device may be preferred by some patients compared with the delivery device of other GLP-1 RAs.
Health inequities	T2DM disproportionately affects minority populations, and significant disparities exist in prevalence, disease control, and rates of complications. ⁶⁵ ICER calculated the Health Improvement Distribution Index, looking at the relative proportion of any health gains from treatment of T2DM for the following groups with a higher prevalence of T2DM than the general US population (see Supplement A1): American Indian/Alaska Native = 1.4 Hispanic = 1.2 Asian Indian = 1.2 Non-Hispanic Black = 1.1
Other (as relevant)	NA

The main potential other benefit of tirzepatide relates to its impact on obesity and obesity-related diseases and complications. The majority of patients in the US with T2DM are overweight or obese, and at risk for obesity-related complications – e.g., obstructive sleep apnea, nonalcoholic fatty liver disease, high blood pressure, cardiovascular disease, and osteoarthritis – any medication that induces weight loss could have beneficial effects outside of its impact on glycemic control. Tirzepatide appears to induce more substantial weight loss in patients than its comparator drugs, and thus may have an impact on obesity-related diseases as well.

Additionally, a substantial number of patients achieved near-normal glycemic control (defined as a HbA1c <5.7%) in SURPASS-2. Not only is this a boost to patients, who cite glycemic control as one of the more important outcomes for managing their diabetes, but it may also slow progression of disease, particularly if achieved early in the disease course.²⁰ Prevention of progression and potentially of development of the micro- and macrovascular complications of T2DM could both improve the productivity of patients and lessen caregiving burden. For example, prevention of diabetic neuropathy may lead to fewer amputations and thus less disability.

6. Health Benefit Price Benchmarks

Health Benefit Price Benchmarks (HBPBs) for the annual cost of treatment with the tirzepatide are presented in Table 6.1 below. The HBPB for a drug is defined as the price range that would achieve incremental cost-effectiveness ratios between \$100,000 and \$150,000 per QALY or per evLY gained. Based on our model simulations, we arrive at a HBPB for tirzepatide from \$5,500 to \$5,700 per QALY gained compared to injectable semaglutide plus background therapy. Equal value of life years gained were not reported given tirzepatide average life years were not greater than injectable semaglutide average life years. Discounting from WAC to reach the threshold price for tirzepatide is not applicable as it is currently based on a placeholder WAC price and should be updated when WAC pricing is established.

Table 6.1. Annual Cost-Effectiveness Health Benefit Price Benchmarks for Tirzepatide plus Background Therapy vs. Semaglutide plus Background Therapy

Outcome for Annual HBPB Calculation	Annual WAC	Annual Price at \$100,000 Threshold	Annual Price at \$150,000 Threshold	Discount from WAC to Reach Threshold Prices
Tirzepatide plus Background Therapy vs. Semaglutide plus Background Therapy				
QALYs Gained	NA*	\$5,500	\$5,700	NA*

HBPB: health benefit price benchmark, QALY: quality-adjusted life year, WAC: wholesale acquisition cost

*Not applicable (NA) as placeholder prices were used

7. Potential Budget Impact

7.1. Overview of Key Assumptions

Using results from the cost-effectiveness model, we estimated the potential budget impact of adding tirzepatide to current background therapy for patients with T2DM with inadequate glycemic control. We used the tirzepatide placeholder price from the base-case analysis (placeholder price of \$4,643.50 per year) and three threshold prices (at \$50,000, \$100,000, and \$150,000 per QALY) when comparing tirzepatide plus background therapy to injectable semaglutide plus background therapy in order to align with the HBPB comparisons made within the cost-effectiveness analyses. Potential budget impact is defined as the total differential cost of using tirzepatide rather than the relevant existing therapy for the treated population, calculated as intervention costs minus any offsets in these costs from averted health care events. All costs were undiscounted and estimated over a five-year time horizon.

The analysis included the estimated number of individuals in the US who would be eligible for treatment with tirzepatide. To estimate the size of the potential candidate population for treatment, we used inputs for the total US adult population size (~265 million)⁶⁶, overall T2DM prevalence (14.6%)⁶⁷, proportion of patients with diagnosed T2DM (76.7%)⁶⁷, and the proportion of patients having failed background therapy and considering a second-line treatment and thus eligible for tirzepatide (16.2%).⁶⁸ Applying these sources results in estimates of 4,800,000 eligible patients in the US. For the purposes of this analysis, we assumed that 20% of these patients would initiate tirzepatide treatment in each of the five years, or approximately 960,000 patients per year. Tirzepatide drew market share proportionally from each of the model comparators over the five-year time horizon.


Market shares were derived from analyst projections based on primary market research, company reports, and key opinion leader surveys.⁶⁹ We set the initial market shares for injectable semaglutide equal to a calculated 2021 market share for the GLP-1 RA class (15.1%), while the market share for empagliflozin was set to a calculated 2021 market share for the SGLT-2 inhibitor class (12.7%). The remaining market share was attributed to background therapy.

The aim of the potential budgetary impact analysis is to document the percentage of patients who could be treated at select prices within five years without crossing a potential budget impact threshold that is aligned with overall growth in the US economy. For 2021-2022, the five-year annualized potential budget impact [threshold](#) that should trigger policy actions to manage access and affordability is calculated to be approximately \$734 million per year for new drugs.

7.2. Results

Assuming the tirzepatide placeholder price of \$4,643.50 per year, 20.1% of the eligible patients could be treated within five years (assuming 20% uptake each year) without crossing the ICER potential budget impact threshold of \$734 million per year. In contrast, 14.4%, 12.7%, and 11.1% of eligible patients could be treated within five years without crossing the ICER potential budget impact threshold at the annual price to reach \$50,000 per QALY (\$5,200), \$100,000 per QALY (\$5,500), or \$150,000 per QALY (\$5,700) versus injectable semaglutide plus background therapy, respectively. Figure 7.1 depicts the potential budgetary impact of tirzepatide at the placeholder price and the three threshold prices. Due to a large eligible population, the budget impact results were very sensitive to the price of tirzepatide and corresponding cost offsets. For instance, a \$129 increase in tirzepatide annual placeholder price (~3%) led to a 10% relative reduction in the number of T2D patients that could be treated without crossing the ICER potential budget impact threshold.

Figure 7.1. Budgetary Impact of Tirzepatide in Adults with T2DM at Placeholder Price and Threshold Annual Prices



References

1. Centers for Disease Control and Prevention USDoHaHS. *National Diabetes Statistics Report, 2020*. Atlanta, GA: Centers for Disease Control and Prevention, U.S. Dept of Health and Human Services;2020.
2. American Diabetes A. 1. Improving Care and Promoting Health in Populations: Standards of Medical Care in Diabetes-2020. *Diabetes Care*. 2020;43(Suppl 1):S7-S13.
3. American Diabetes A. 6. Glycemic Targets: Standards of Medical Care in Diabetes-2020. *Diabetes Care*. 2020;43(Suppl 1):S66-S76.
4. American Diabetes A. 9. Pharmacologic Approaches to Glycemic Treatment: Standards of Medical Care in Diabetes-2020. *Diabetes Care*. 2020;43(Suppl 1):S98-S110.
5. American Diabetes A. 10. Cardiovascular Disease and Risk Management: Standards of Medical Care in Diabetes-2020. *Diabetes Care*. 2020;43(Suppl 1):S111-S134.
6. Frias JP, Nauck MA, Van J, et al. Efficacy and safety of LY3298176, a novel dual GIP and GLP-1 receptor agonist, in patients with type 2 diabetes: a randomised, placebo-controlled and active comparator-controlled phase 2 trial. *The Lancet (British edition)*. 2018;392(10160):2180-2193.
7. Frias JP, Davies MJ, Rosenstock J, et al. Tirzepatide versus Semaglutide Once Weekly in Patients with Type 2 Diabetes (SURPASS-2). *The New England journal of medicine*. 2021;385(6):503-515.
8. Marso SP, Bain SC, Consoli A, et al. Semaglutide and Cardiovascular Outcomes in Patients with Type 2 Diabetes (SUSTAIN-6). *The New England journal of medicine*. 2016;375(19):1834-1844.
9. Zinman B, Wanner C, Lachin JM, et al. Empagliflozin, Cardiovascular Outcomes, and Mortality in Type 2 Diabetes (EMPA-REG-OUTCOME). *The New England journal of medicine*. 2015;373(22):2117-2128.
10. Wanner C, Inzucchi SE, Lachin JM, et al. Empagliflozin and Progression of Kidney Disease in Type 2 Diabetes. *N Engl J Med*. 2016;375(4):323-334.
11. Hayes AJ, Leal J, Gray A, Holman R, Clarke P. UKPDS outcomes model 2: a new version of a model to simulate lifetime health outcomes of patients with type 2 diabetes mellitus using data from the 30 year United Kingdom Prospective Diabetes Study: UKPDS 82. *Diabetologia*. 2013;56(9):1925-1933.
12. Neumann PJ, Sanders GD, Russell LB, Siegel JE, Ganiats TG. *Cost-effectiveness in health and medicine*. Oxford University Press; 2016.
13. Fowler MJ. Microvascular and Macrovascular Complications of Diabetes. *Clinical Diabetes*. 2008;26(2):77-82.
14. Bae JP, Lage MJ, Mo D, Nelson DR, Hoogwerf BJ. Obesity and glycemic control in patients with diabetes mellitus: Analysis of physician electronic health records in the US from 2009-2011. *J Diabetes Complications*. 2016;30(2):212-220.
15. American Diabetes A. 8. Obesity Management for the Treatment of Type 2 Diabetes: Standards of Medical Care in Diabetes-2020. *Diabetes Care*. 2020;43(Suppl 1):S89-S97.
16. Piccolo RS, Subramanian SV, Pearce N, Florez JC, McKinlay JB. Relative Contributions of Socioeconomic, Local Environmental, Psychosocial, Lifestyle/Behavioral, Biophysiological, and Ancestral Factors to Racial/Ethnic Disparities in Type 2 Diabetes. *Diabetes Care*. 2016;39(7):1208-1217.
17. Meng YY, Diamant A, Jones J, et al. Racial and Ethnic Disparities in Diabetes Care and Impact of Vendor-Based Disease Management Programs. *Diabetes Care*. 2016;39(5):743-749.

18. Caraballo C, Valero-Elizondo J, Khera R, et al. Burden and Consequences of Financial Hardship From Medical Bills Among Nonelderly Adults With Diabetes Mellitus in the United States. *Circ Cardiovasc Qual Outcomes*. 2020;13(2):e006139.
19. American Diabetes A. 6. Glycemic Targets: Standards of Medical Care in Diabetes-2021. *Diabetes Care*. 2021;44(Suppl 1):S73-s84.
20. Ismail-Beigi F, Craven T, Banerji MA, et al. Effect of intensive treatment of hyperglycaemia on microvascular outcomes in type 2 diabetes: an analysis of the ACCORD randomised trial. *Lancet*. 2010;376(9739):419-430.
21. American Diabetes A. Addendum. 10. Cardiovascular Disease and Risk Management: Standards of Medical Care in Diabetes-2020. *Diabetes Care* 2020;43(Suppl. 1):S111-S134. *Diabetes Care*. 2020;43(8):1977-1978.
22. He X, Li J, Wang B, et al. Diabetes self-management education reduces risk of all-cause mortality in type 2 diabetes patients: a systematic review and meta-analysis. *Endocrine*. 2017;55(3):712-731.
23. Strawbridge LM, Lloyd JT, Meadow A, Riley GF, Howell BL. One-Year Outcomes of Diabetes Self-Management Training Among Medicare Beneficiaries Newly Diagnosed With Diabetes. *Med Care*. 2017;55(4):391-397.
24. Han Y, Xie H, Liu Y, Gao P, Yang X, Shen Z. Effect of metformin on all-cause and cardiovascular mortality in patients with coronary artery diseases: a systematic review and an updated meta-analysis. *Cardiovasc Diabetol*. 2019;18(1):96.
25. Bradley C, Lewis KS. Measures of psychological well-being and treatment satisfaction developed from the responses of people with tablet-treated diabetes. *Diabet Med*. 1990;7(5):445-451.
26. Sun F, Chai S, Yu K, et al. Gastrointestinal adverse events of glucagon-like peptide-1 receptor agonists in patients with type 2 diabetes: a systematic review and network meta-analysis. *Diabetes technology & therapeutics*. 2015;17(1):35-42.
27. Chaplin S. SGLT2 inhibitors and risk of genitourinary infections. *Prescriber*. 2016;27(12):26-30.
28. Douros A, Lix LM, Fralick M, et al. Sodium–glucose cotransporter-2 inhibitors and the risk for diabetic ketoacidosis: a multicenter cohort study. *Annals of Internal Medicine*. 2020;173(6):417-425.
29. Administration. UDoHaHSFaD. Guidance for Industry. Diabetes Mellitus - Evaluating Cardiovascular Risk in New Antidiabetic Therapies to Treat Type 2 Diabetes. In: Food and Drug Administration H, ed2008:77724-77725.
30. Gelhorn H, Balantac Z, Shinde S, Thieu VT, Boye KS. The Burden of Type 2 Diabetes and the Value of Achieving Near Normoglycemia from the Patient Perspective. *Diabetes Ther*. 2021;12(7):1821-1837.
31. Kim YY, Lee JS, Kang HJ, Park SM. Effect of medication adherence on long-term all-cause-mortality and hospitalization for cardiovascular disease in 65,067 newly diagnosed type 2 diabetes patients. *Sci Rep*. 2018;8(1):12190.
32. Min T, Bain SC. The Role of Tirzepatide, Dual GIP and GLP-1 Receptor Agonist, in the Management of Type 2 Diabetes: The SURPASS Clinical Trials. *Diabetes therapy*. 2021;12(1):143-157.
33. Frias JP, Nauck MA, Van J, et al. Efficacy and tolerability of tirzepatide, a dual glucose-dependent insulinotropic peptide and glucagon-like peptide-1 receptor agonist in patients with type 2 diabetes: A 12-week, randomized, double-blind, placebo-controlled study to evaluate different dose-escalation regimens. *Diabetes, obesity & metabolism*. 2020;22(6):938-946.
34. Trials C. A Study of Tirzepatide (LY3298176) Compared With Dulaglutide on Major Cardiovascular Events in Participants With Type 2 Diabetes (SURPASS-CVOT). U.S National

- Library of Medicine. <https://clinicaltrials.gov/ct2/show/NCT04255433>. Published 2021. Accessed 2021.
35. Del Prato S, Kahn S. Tirzepatide versus insulin glargine in type 2 diabetes and increased cardiovascular risk (SURPASS-4): a randomised, open-label, parallel-group, multicentre, phase 3 trial. *The Lancet*. 2021.
 36. Zinman B, Inzucchi SE, Lachin JM, et al. Rationale, design, and baseline characteristics of a randomized, placebo-controlled cardiovascular outcome trial of empagliflozin (EMPA-REG OUTCOME™). *Cardiovascular diabetology*. 2014;13(1):1-8.
 37. Association AD. 8. Cardiovascular Disease and Risk Management. *Diabetes Care*. 2016;39(Supplement 1):S60-S71.
 38. Lilly E. Data on File. In: 2021.
 39. Rosenstock J, Allison D, Birkenfeld AL, et al. Effect of Additional Oral Semaglutide vs Sitagliptin on Glycated Hemoglobin in Adults With Type 2 Diabetes Uncontrolled With Metformin Alone or With Sulfonylurea: The PIONEER 3 Randomized Clinical Trial. *JAMA*. 2019;321(15):1466-1480.
 40. Rind DM, Guzaukas G, Fazioli K, et al. *Oral Semaglutide for Type 2 Diabetes: Effectiveness and Value*. Institute for Clinical and Economic Review; 2019.
 41. Lilly's tirzepatide achieves all primary and key secondary study outcomes against insulin glargine in adults with type 2 diabetes and increased cardiovascular risk in SURPASS-4 trial [press release]. Lilly 2021.
 42. Bain SC, Klufas MA, Ho A, Matthews DR. Worsening of diabetic retinopathy with rapid improvement in systemic glucose control: a review. *Diabetes, Obesity and Metabolism*. 2019;21(3):454-466.
 43. FDA. OZEMPIC (semaglutide) Injection Package Insert. In. U.S Food & Drug Administration 2021.
 44. FDA. JARDIANCE (Empagliflozin tablets) Package Insert. In. U.S Food & Drug Administration 2021.
 45. US FDA grants Fast Track designation to Jardiance® for the treatment of chronic kidney disease [press release]. 2021.
 46. American Diabetes A. 11. Microvascular Complications and Foot Care: Standards of Medical Care in Diabetes-2020. *Diabetes Care*. 2020;43(Suppl 1):S135-S151.
 47. Filippatos TD, Panagiotopoulou TV, Elisaf MS. Adverse effects of GLP-1 receptor agonists. *The review of diabetic studies: RDS*. 2014;11(3):202.
 48. Heimbürger SM, Bergmann NC, Augustin R, Gasbjerg LS, Christensen MB, Knop FK. Glucose-dependent insulinotropic polypeptide (GIP) and cardiovascular disease. *Peptides*. 2020;125:170174.
 49. Jujic A, Atabaki-Pasdar N, Nilsson PM, et al. Glucose-dependent insulinotropic peptide and risk of cardiovascular events and mortality: a prospective study. *Diabetologia*. 2020;63(5):1043-1054.
 50. Kahles F, Ruckbeil MV, Arrivas MC, et al. Association of Glucose-Dependent Insulinotropic Polypeptide Levels With Cardiovascular Mortality in Patients With Acute Myocardial Infarction. *J Am Heart Assoc*. 2021;10(13):e019477.
 51. Mori Y, Matsui T, Hirano T, Yamagishi SI. GIP as a Potential Therapeutic Target for Atherosclerotic Cardiovascular Disease-A Systematic Review. *Int J Mol Sci*. 2020;21(4).
 52. Sattar N, Lee MMY, Kristensen SL, et al. Cardiovascular, mortality, and kidney outcomes with GLP-1 receptor agonists in patients with type 2 diabetes: a systematic review and meta-analysis of randomised trials. *Lancet Diabetes Endocrinol*. 2021;9(10):653-662.
 53. Laiteerapong N, Cooper JM, Skandari MR, et al. Individualized Glycemic Control for U.S. Adults With Type 2 Diabetes: A Cost-Effectiveness Analysis. *Ann Intern Med*. 2018;168(3):170-178.

54. Guzauskas GF, Rind DM, Fazioli K, Chapman RH, Pearson SD, Hansen RN. Cost-effectiveness of oral semaglutide added to current antihyperglycemic treatment for type 2 diabetes. *Journal of Managed Care & Specialty Pharmacy*. 2021;27(4):455-468.
55. Shao H, Fonseca V, Stoecker C, Liu S, Shi L. Novel risk engine for diabetes progression and mortality in USA: building, relating, assessing, and validating outcomes (BRAVO). *Pharmacoeconomics*. 2018;36(9):1125-1134.
56. Hunt B, Malkin SJ, Moes RG, Huisman EL, Vandebrouck T, Wolffenbuttel BH. Once-weekly semaglutide for patients with type 2 diabetes: a cost-effectiveness analysis in the Netherlands. *BMJ Open Diabetes Research and Care*. 2019;7(1):e000705.
57. Johansen P, Chubb B, Hunt B, Malkin SJP, Sandberg A, Capehorn M. Evaluating the Long-Term Cost-Effectiveness of Once-Weekly Semaglutide Versus Once-Daily Liraglutide for the Treatment of Type 2 Diabetes in the UK. *Advances in Therapy*. 2020;37(5):2427-2441.
58. Lee M, Lauren BN, Zhan T, et al. The cost-effectiveness of pharmacotherapy and lifestyle intervention in the treatment of obesity. *Obes Sci Pract*. 2019;6(2):162-170.
59. Hong D, Si L, Jiang M, et al. Cost effectiveness of sodium-glucose cotransporter-2 (SGLT2) inhibitors, glucagon-like peptide-1 (GLP-1) receptor agonists, and dipeptidyl peptidase-4 (DPP-4) inhibitors: a systematic review. *Pharmacoeconomics*. 2019;37(6):777-818.
60. Reifsnider OS, Kansal AR, Gandhi PK, et al. Cost-effectiveness of empagliflozin versus canagliflozin, dapagliflozin, or standard of care in patients with type 2 diabetes and established cardiovascular disease. *BMJ Open Diabetes Research and Care*. 2021;9(1):e001313.
61. Willis M, Asseburg C, Nilsson A, Johnsson K, Kartman B. Multivariate Prediction Equations for HbA1c Lowering, Weight Change, and Hypoglycemic Events Associated with Insulin Rescue Medication in Type 2 Diabetes Mellitus: Informing Economic Modeling. *Value in health : the journal of the International Society for Pharmacoeconomics and Outcomes Research*. 2017;20(3):357-371.
62. SSR Health L. <https://www.ssrhealth.com/>. Published 2018. Accessed.
63. Ward A, Alvarez P, Vo L, Martin S. Direct medical costs of complications of diabetes in the United States: estimates for event-year and annual state costs (USD 2012). *J Med Econ*. 2014;17(3):176-183.
64. Yang W, Cintina I, Hoerger T, et al. Estimating costs of diabetes complications in people < 65 years in the US using panel data. *Journal of Diabetes and its Complications*. 2020;34(12):107735.
65. Peek ME, Cargill A, Huang ES. Diabetes health disparities: a systematic review of health care interventions. *Med Care Res Rev*. 2007;64(5 Suppl):101S-156S.
66. US Census Bureau. 2017 National Population Projections Datasets. Projected Population by Single Year of Age, Sex, Race, and Hispanic Origin for the United States: 2016 to 2060. 2017.
67. Wang L, Li X, Wang Z, et al. Trends in Prevalence of Diabetes and Control of Risk Factors in Diabetes Among US Adults, 1999-2018. *JAMA*. 2021;326(8):704-716.
68. Centers for Disease Control and Prevention. National Health and Nutrition Examination Survey. In: Prevention CfDCa, ed2018.
69. Datamonitor. Patient-Based Forecast Model - Diabetes Type 2. 2021.
70. American Diabetes A. 2. Classification and Diagnosis of Diabetes: Standards of Medical Care in Diabetes-2020. *Diabetes Care*. 2020;43(Suppl 1):S14-S31.
71. Group UPDS. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *The lancet*. 1998;352(9131):837-853.
72. Grundy SM, Stone NJ, Bailey AL, et al. 2018
AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APHA/ASPC/NLA/PCNA Guideline on the

- Management of Blood Cholesterol: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. 2019;139(25):e1082-e1143.
73. National Cholesterol Education Program Expert Panel on Detection E, Treatment of High Blood Cholesterol in A. Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation*. 2002;106(25):3143-3421.
 74. Tsai IT, Wang CP, Lu YC, et al. The burden of major adverse cardiac events in patients with coronary artery disease. *BMC Cardiovasc Disord*. 2017;17(1):1.
 75. Brazier JE, Harper R, Jones NM, et al. Validating the SF-36 health survey questionnaire: new outcome measure for primary care. *BMJ*. 1992;305(6846):160-164.
 76. Kolotkin RL, Head S, Brookhart A. Construct validity of the Impact of Weight on Quality of Life Questionnaire. *Obes Res*. 1997;5(5):434-441.
 77. Karter AJ, Parker MM, Moffet HH, Gilliam LK, Dlott R. Association of Real-time Continuous Glucose Monitoring With Glycemic Control and Acute Metabolic Events Among Patients With Insulin-Treated Diabetes. *JAMA*. 2021;325(22):2273-2284.
 78. Martens TW, Beck RW, Bergenstal RM. Continuous Glucose Monitoring and Glycemic Control in Patients With Type 2 Diabetes Treated With Basal Insulin-Reply. *JAMA*. 2021;326(13):1330-1331.
 79. Rosenstock J, Kahn SE, Johansen OE, et al. Effect of Linagliptin vs Glimepiride on Major Adverse Cardiovascular Outcomes in Patients With Type 2 Diabetes: The CAROLINA Randomized Clinical Trial. *JAMA*. 2019;322(12):1155-1166.
 80. Sherifali D, Nerenberg K, Pullenayegum E, Cheng JE, Gerstein HC. The effect of oral antidiabetic agents on A1C levels: a systematic review and meta-analysis. *Diabetes Care*. 2010;33(8):1859-1864.
 81. Singh S, Loke YK, Furberg CD. Long-term risk of cardiovascular events with rosiglitazone: a meta-analysis. *JAMA*. 2007;298(10):1189-1195.
 82. Dormandy JA, Charbonnel B, Eckland DJ, et al. Secondary prevention of macrovascular events in patients with type 2 diabetes in the PROactive Study (PROspective pioglitAzone Clinical Trial In macroVascular Events): a randomised controlled trial. *Lancet*. 2005;366(9493):1279-1289.
 83. Home PD, Pocock SJ, Beck-Nielsen H, et al. Rosiglitazone evaluated for cardiovascular outcomes in oral agent combination therapy for type 2 diabetes (RECORD): a multicentre, randomised, open-label trial. *Lancet*. 2009;373(9681):2125-2135.
 84. Kahn SE, Zinman B, Lachin JM, et al. Rosiglitazone-associated fractures in type 2 diabetes: an Analysis from A Diabetes Outcome Progression Trial (ADOPT). *Diabetes Care*. 2008;31(5):845-851.
 85. Pratley RE, Salsali A. Inhibition of DPP-4: a new therapeutic approach for the treatment of type 2 diabetes. *Curr Med Res Opin*. 2007;23(4):919-931.
 86. Amori RE, Lau J, Pittas AG. Efficacy and safety of incretin therapy in type 2 diabetes: systematic review and meta-analysis. *JAMA*. 2007;298(2):194-206.
 87. McGuire DK, Shih WJ, Cosentino F, et al. Association of SGLT2 Inhibitors With Cardiovascular and Kidney Outcomes in Patients With Type 2 Diabetes: A Meta-analysis. *JAMA Cardiol*. 2021;6(2):148-158.
 88. Ueda P, Svanstrom H, Melbye M, et al. Sodium glucose cotransporter 2 inhibitors and risk of serious adverse events: nationwide register based cohort study. *BMJ*. 2018;363:k4365.
 89. Hinnen D. Glucagon-Like Peptide 1 Receptor Agonists for Type 2 Diabetes. *Diabetes Spectr*. 2017;30(3):202-210.

90. Samms RJ, Coghlan MP, Sloop KW. How May GIP Enhance the Therapeutic Efficacy of GLP-1? *Trends Endocrinol Metab.* 2020;31(6):410-421.
91. Garber AJ, Handelsman Y, Grunberger G, et al. Consensus Statement by the American Association of Clinical Endocrinologists and American College of Endocrinology on the Comprehensive Type 2 Diabetes Management Algorithm - 2020 Executive Summary. *Endocr Pract.* 2020;26(1):107-139.
92. Cosentino F, Grant PJ, Aboyans V, et al. 2019 ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD: the Task Force for diabetes, pre-diabetes, and cardiovascular diseases of the European Society of Cardiology (ESC) and the European Association for the Study of Diabetes (EASD). *European heart journal.* 2020;41(2):255-323.
93. Excellence NfHaC. Type 2 diabetes in adults: management. In:2015.
94. Cook DJ, Mulrow CD, Haynes RB. Systematic reviews: synthesis of best evidence for clinical decisions. *Ann Intern Med.* 1997;126(5):376-380.
95. Higgins J, Thomas, J, Chandler, J, Cumpston, M, Li, T, Page, MJ, Welch, VA. Cochrane Handbook for Systematic Reviews of Interventions version 6.1 (updated September 2020). <https://training.cochrane.org/handbook/current>. Published 2020. Accessed.
96. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS medicine.* 2009;6(7):e1000097.
97. Agency for Healthcare Research and Quality. U.S. Preventive Services Task Force Procedure Manual. Published 2008. Accessed.
98. Ollendorf DA, Pearson SD. An integrated evidence rating to frame comparative effectiveness assessments for decision makers. *Med Care.* 2010;48(6 Suppl):S145-152.
99. Ollendorf D, Pearson, SD. ICER Evidence Rating Matrix: A User's Guide. <https://icer.org/evidence-rating-matrix/>. Published 2020. Updated January 31, 2020. Accessed.
100. Rhodes KM, Turner RM, Higgins JP. Predictive distributions were developed for the extent of heterogeneity in meta-analyses of continuous outcome data. *Journal of clinical epidemiology.* 2015;68(1):52-60.
101. Ahrén BP, Masmiquel LP, Kumar HM, et al. Efficacy and safety of once-weekly semaglutide versus once-daily sitagliptin as an add-on to metformin, thiazolidinediones, or both, in patients with type 2 diabetes (SUSTAIN 2): a 56-week, double-blind, phase 3a, randomised trial. *The lancet Diabetes & endocrinology.* 2017;5(5):341-354.
102. Rodbard HW, Rosenstock J, Canani LH, et al. Oral Semaglutide versus Empagliflozin in Patients with Type 2 Diabetes Uncontrolled on Metformin: The PIONEER 2 Trial. *Diabetes Care.* 2019;dc190883.
103. Ahren B, Johnson SL, Stewart M, et al. HARMONY 3: 104-Week Randomized, Double-Blind, Placebo- and Active-Controlled Trial Assessing the Efficacy and Safety of Albiglutide Compared With Placebo, Sitagliptin, and Glimepiride in Patients With Type 2 Diabetes Taking Metformin. *Diabetes care.* 2014;37(8):2141-2148.
104. Salsali A, Kim G, Woerle HJ, Broedl UC, Hantel S. Cardiovascular safety of empagliflozin in patients with type 2 diabetes: a meta-analysis of data from randomized placebo-controlled trials. *Diabetes, obesity & metabolism.* 2016;18(10):1034-1040.
105. Shi F-H, Li H, Cui M, Zhang Z-L, Gu Z-C, Liu X-Y. Efficacy and Safety of Once-Weekly Semaglutide for the Treatment of Type 2 Diabetes: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Frontiers in pharmacology.* 2018;9:576-576.

106. Sanders GD, Neumann PJ, Basu A, et al. Recommendations for Conduct, Methodological Practices, and Reporting of Cost-effectiveness Analyses: Second Panel on Cost-Effectiveness in Health and Medicine. *Jama*. 2016;316(10):1093-1103.
107. Lundqvist A, Steen Carlsson K, Johansen P, Andersson E, Willis M. Validation of the IHE cohort model of type 2 diabetes and the impact of choice of macrovascular risk equations. *PLoS One*. 2014;9(10):e110235.
108. McEwan P, Foos V, Palmer JL, Lamotte M, Lloyd A, Grant D. Validation of the IMS CORE diabetes model. *Value in Health*. 2014;17(6):714-724.
109. Allison MA, Ho E, Denenberg JO, et al. Ethnic-specific prevalence of peripheral arterial disease in the United States. *American journal of preventive medicine*. 2007;32(4):328-333.
110. Weng L-C, Preis SR, Hulme OL, et al. Genetic predisposition, clinical risk factor burden, and lifetime risk of atrial fibrillation. *Circulation*. 2018;137(10):1027-1038.
111. Aune D, Feng T, Schlesinger S, Janszky I, Norat T, Riboli E. Diabetes mellitus, blood glucose and the risk of atrial fibrillation: a systematic review and meta-analysis of cohort studies. *Journal of diabetes and its complications*. 2018;32(5):501-511.
112. Neuwahl SJ, Zhang P, Chen H, et al. Patient Health Utility Equations for a Type 2 Diabetes Model. *Diabetes Care*. 2021;44(2):381-389.
113. Shao H, Yang S, Fonseca V, Stoecker C, Shi L. Estimating quality of life decrements due to diabetes complications in the United States: the Health Utility Index (HUI) diabetes complication equation. *Pharmacoeconomics*. 2019;37(7):921-929.
114. Boye KS, Matza LS, Walter KN, Van Brunt K, Palsgrove AC, Tynan A. Utilities and disutilities for attributes of injectable treatments for type 2 diabetes. *The European Journal of Health Economics*. 2011;12(3):219-230.
115. Sullivan PW, Ghushchyan VH. EQ-5D scores for diabetes-related comorbidities. *Value in health*. 2016;19(8):1002-1008.
116. Dall TM, Yang W, Gillespie K, et al. The economic burden of elevated blood glucose levels in 2017: diagnosed and undiagnosed diabetes, gestational diabetes mellitus, and prediabetes. *Diabetes care*. 2019;42(9):1661-1668.
117. Nguyen E, Coleman CI, Nair S, Weeda ER. Cost-utility of empagliflozin in patients with type 2 diabetes at high cardiovascular risk. *Journal of Diabetes and its Complications*. 2018;32(2):210-215.
118. Kuo S, Ye W, Duong J, Herman WH. Are the favorable cardiovascular outcomes of empagliflozin treatment explained by its effects on multiple cardiometabolic risk factors? A simulation of the results of the EMPA-REG OUTCOME trial. *Diabetes research and clinical practice*. 2018;141:181-189.
119. Ramos M, Men P, Wang X, Ustyugova A, Lamotte M. Cost-effectiveness of empagliflozin in patients with type 2 diabetes and established cardiovascular disease in China. *Cost Effectiveness and Resource Allocation*. 2021;19(1):1-11.
120. Capehorn M, Hallén N, Baker-Knight J, Glah D, Hunt B. Evaluating the Cost-Effectiveness of Once-Weekly Semaglutide 1 mg Versus Empagliflozin 25 mg for Treatment of Patients with Type 2 Diabetes in the UK Setting. *Diabetes Therapy*. 2021;12(2):537-555.
121. Si L, Willis MS, Asseburg C, et al. Evaluating the ability of economic models of diabetes to simulate new cardiovascular outcomes trials: a report on the ninth mount hood diabetes challenge. *Value in Health*. 2020;23(9):1163-1170.
122. Ehlers LH, Lamotte M, Ramos MC, et al. The cost-effectiveness of oral semaglutide versus empagliflozin in Type 2 diabetes in Denmark. *Journal of comparative effectiveness research*. 2021;11(1):29-37.

123. Malkin SJ, Hunt B, Huisman EL, Grand TS, Chubb B. The long-term cost-effectiveness of oral semaglutide in the Netherlands based on the PIONEER 2, 3 and 4 randomized controlled trials. *Diabetes Research and Clinical Practice*. 2021;175:108759.
124. Ramos M, Ustyugova A, Hau N, Lamotte M. Cost–effectiveness of empagliflozin compared with liraglutide based on cardiovascular outcome trials in Type II diabetes. *Journal of Comparative Effectiveness Research*. 2020;9(11):781-794.
125. Institute for Clinical and Economic Review. 2020-2023 Value Assessment Framework. <https://icer.org/our-approach/methods-process/value-assessment-framework/>. Published 2020. Accessed.
126. Pearson SD. The ICER Value Framework: Integrating Cost Effectiveness and Affordability in the Assessment of Health Care Value. *Value in health : the journal of the International Society for Pharmacoeconomics and Outcomes Research*. 2018;21(3):258-265.

Supplemental Materials

A. Background: Supplemental Information

A1. Definitions

Diagnosis of Type 2 Diabetes Mellitus (T2DM): T2DM is characterized by progressive loss of adequate insulin secretion by pancreatic beta cells, frequently on the background of insulin resistance.⁷⁰ The diagnosis is made if any of the following criteria are met: (a) fasting plasma glucose ≥ 126 mg/dL; (b) 2-hour post-prandial glucose ≥ 200 mg/dL during 75-gram oral glucose tolerance test; (c) HbA1c $\geq 6.5\%$; or (d) random plasma glucose ≥ 200 mg/dL and symptoms of hyperglycemia.

Glycated Hemoglobin (HbA1c): Blood test that measures the amount of glycated red blood cells in the blood. It reflects average blood sugar levels over the past 2-3 months and increases as the amount of glucose increases in the blood. Patients with HbA1c $\geq 6.5\%$ are considered to have diabetes.

Microvascular complications: Persistent exposure to high levels of blood glucose can lead to damage to small blood vessels in the eyes, kidney, and nerves.

- Retinopathy (eye): Most common microvascular complication, causing around 10,000 new cases of blindness each year in the US.¹³ Most cases develop within 7-20 years after diagnosis.⁷¹
- Nephropathy (kidney): Leading cause of renal failure in the US, can be present at diagnosis.^{13,46} Characterized by elevated levels of protein in the urine and/or decreased estimated glomerular filtration rate (eGFR).⁴⁶
- Neuropathy (nerves): Characterized by symptoms or signs of peripheral nerve dysfunction (e.g., burning, tingling, numbness, sensory loss to light touch, vibration or temperature, autonomic dysfunction) in people with diabetes, after excluding other causes.⁴⁶ Can lead to foot ulcers, injury from falls, and ultimately limb amputation.¹³

Macrovascular complications: Diseases characterized by atherosclerosis, including coronary artery disease and cerebrovascular disease. Patients with T2DM have a much higher risk of cardiovascular events such as myocardial infarction or stroke, and cardiovascular death is the most common cause of death in patients with diabetes.¹³

Cardiovascular Outcomes Trial (CVOT): Long-term, prospective trials of diabetes drugs specifically examining cardiovascular safety. CVOTs became required by the FDA in 2008 for the approval of new diabetes drugs due to cardiovascular safety concerns raised by rosiglitazone, a thiazolidinedione.²⁹ Trial results must demonstrate that the upper bound of the 95% confidence interval of the hazard ratio for cardiovascular events is less than 1.8 for a drug to be considered safe from a cardiovascular perspective. CVOTs have been conducted for newer diabetes drugs, including DPP-4 inhibitors, GLP-1 RA, and SGLT-2 inhibitors.

Atherosclerotic Cardiovascular Disease (ASCVD): Disease of the arteries caused by plaque buildup in artery walls. ASCVD includes the clinical conditions of coronary artery disease, acute coronary syndromes, stroke, transient ischemic attack, peripheral vascular disease, coronary or other arterial revascularization, and aortic aneurysm.⁷² Patients with diabetes are at higher than normal risk for ASCVD.⁷³

Major Adverse Cardiovascular Events (MACE): The major causes of morbidity and death in patients with ASCVD, and an often-used endpoint in clinical trials. There is no standard definition of MACE, but in general it can include: fatal and non-fatal myocardial infarction, heart failure, recurrent angina pain, repeat hospitalization for cardiovascular-related illness, repeat percutaneous coronary intervention, coronary bypass surgery, fatal and non-fatal stroke, and all-cause mortality.⁷⁴ For diabetes drugs, 3-point MACE (MACE-3) is often used as an endpoint in CVOT, including non-fatal myocardial infarction (may or may not include silent infarction), non-fatal stroke, and cardiovascular mortality.

Congestive Heart Failure: A chronic condition where the heart does not pump enough blood for the body's needs, leading to inadequate blood flow to vital organs (e.g., kidneys) and buildup of fluid in other organs (e.g., lungs). This can happen when the heart muscle is weakened or is too stiff. The most common symptoms of heart failure are shortness of breath, leg swelling, and fatigue.

Chronic kidney disease (CKD): A chronic condition where kidney function is decreased, resulting in buildup of waste and fluid in the body. It is defined as a reduction in eGFR to 60 mL/min/1.73 m² or less and/or the presence of protein in the urine. Symptoms of CKD often do not occur until the advanced stages, and can include edema, loss of appetite, nausea, fatigue, high blood pressure, anemia, high potassium, and bone disease. Diabetes is the leading cause of CKD in the US.¹³

Short Form (36) Health Survey (SF-36): A 36-item self-reported questionnaire of health-related quality of life. It includes questions about general health, activity limitations, physical and emotional health, social activities, and pain.⁷⁵ It is often used as a quality-of-life measure in clinical trials and is not specific to any health condition.

Diabetes Treatment Satisfaction Questionnaire (DTSQ): An 8-item self-reported questionnaire of psychological well-being (depression, anxiety, and positive well-being) and treatment satisfaction in

patients with T2DM.²⁵ It is widely used in clinical trials to assess the impact of diabetes interventions on quality of life.

Impact of Weight on Quality-of-Life Questionnaire: A 74-item self-reported questionnaire assessing the effects of obesity on health-related quality of life. The scale measures the impact of obesity on physical function, self-esteem, sexual life, public distress, and work.⁷⁶

Continuous Glucose Monitor (CGM): A non-invasive device that measures interstitial blood glucose levels constantly through skin sensors, providing real-time information about blood glucose levels. Use of CGMs can result in better glycemic control compared with regular blood glucose monitoring in patients with T2DM on insulin therapy.^{77,78}

Health Improvement Distribution Index: The Health Improvement Distribution Index identifies a subpopulation that has a higher prevalence of the disease of interest and therefore, creates an opportunity for proportionately more health gains within the subpopulation. This opportunity may be realized by achieving equal access both within and outside the identified subpopulation to an intervention that is known to improve health. The Health Improvement Distribution Index is defined as the disease prevalence in the subpopulation divided by the disease prevalence in the overall population. For example, if the disease prevalence was 10% in poor Americans whereas the disease prevalence across all Americans was 4%, then the Health Improvement Distribution Index would be $10\%/4\% = 2.5$. For interventions known to increase health in this disease and that accomplish equal access across the entire population, poor Americans would receive 2.5 times the health improvements as compared to the same sized group of Americans without regard to economic status. Health Improvement Distribution Indexes above one suggest that more health may be gained on the relative scale in the subpopulation of interest when compared to the population as a whole. This statistic may be helpful in characterizing a treatment's contextual considerations and potential other benefits (Section 5).

For this calculation, we used data from the Centers for Disease Control 2020 National Diabetes Statistics Report.¹ We used the overall US population prevalence of diagnosed diabetes, 10.2%, as the denominator. We performed calculations for the following subgroups:

- American Indian/Alaska Native: $14.7\%/10.2\% = 1.4$
- Asian Indian: $12.6\%/10.2\% = 1.2$
- Hispanic: $12.5\%/10.2\% = 1.2$
- Black: $11.7\%/10.2\% = 1.1$

A2. Drug Classes for the Treatment of Type 2 Diabetes

Metformin: An orally administered biguanide that decreases glucose production and absorption and improves insulin sensitivity. It is recommended as initial pharmacotherapy for patients with T2DM due to its efficacy and favorable safety profile.³ It decreases HbA1c by 1.1% on average without significant risk of hypoglycemia. Additionally, it is associated with modest weight loss and may improve cardiovascular outcomes, though there are no direct cardiovascular outcomes trials.²⁴ Side effects from metformin are mainly gastrointestinal; there is also a rare risk of lactic acidosis, and the drug should be discontinued in patients with severe chronic kidney disease. Metformin is recommended to be continued as long as it is tolerated and not contraindicated.³

Sulfonylureas (SU): Oral hypoglycemic agents such as glyburide and glipizide that increase insulin secretion by stimulating pancreatic beta cells. They lower HbA1c by 1-2%. The most common side effects are hypoglycemia and weight gain. Sulfonylureas may be associated with progressive dysfunction of pancreatic beta cells and worsening diabetes control in the long-term. Sulfonylureas do not appear to have an impact on cardiovascular outcomes.⁷⁹

Thiazolidinediones (TZD): Oral hypoglycemic agents such as pioglitazone that lower blood glucose by decreasing insulin resistance and decreasing glucose production in the liver. TZDs lower HbA1c by around 1-1.25%⁸⁰, and also have favorable effects on lipids and hepatic steatosis. TZDs have mixed cardiovascular data. Rosiglitazone has been associated with an increased risk of myocardial infarction⁸¹, however, pioglitazone has been shown to decrease fatal and non-fatal stroke in patients with previous strokes.⁸² TZDs are also associated with weight gain and fluid retention, and an increased incidence of heart failure and heart failure hospitalizations.^{82,83} There is also an increased risk of bone fractures with long-term use of TZDs.⁸⁴

Dipeptidyl peptidase-4 (DPP-4) inhibitors: Oral hypoglycemic agents such as sitagliptin, saxagliptin, linagliptin, and alogliptin that lower blood glucose by inhibiting the activity of the DPP-4 enzyme in the plasma. Inhibition of DPP-4 stops degradation of incretins such as GLP-1 and GIP, which in turn increases insulin secretion and decreases gastric emptying.⁸⁵ DPP-4 inhibitors lower HbA1c on average by around 0.75%.⁸⁶ Hypoglycemia is rare, and they are considered weight neutral. However, DPP-4 inhibitors do not appear to impact cardiovascular outcomes.⁷⁹ Common side effects include runny nose, headache, and diarrhea. DPP-4 inhibitors have also been associated with more severe side effects such as severe joint pains, pancreatitis, angioedema, and Stevens-Johnson syndrome.


Sodium-glucose co-transporter 2 (SGLT-2) inhibitors: Oral hypoglycemic agents such as empagliflozin, canagliflozin, and dapagliflozin. SGLT-2 is a protein that is involved in the reabsorption of glucose filtered by the kidneys, and in patients with T2DM, SGLT-2 appears to be upregulated, increasing the rate of renal glucose reabsorption. SGLT-2 inhibitors block glucose reabsorption, resulting in loss of glucose in the urine. As add-on therapy to metformin and/or sulfonylureas, SGLT-2 inhibitors lowered HbA1c by up to an additional 1%. Hypoglycemia is rare with SGLT-2 inhibitors and modest weight loss has been observed. SGLT-2 inhibitors have been shown to reduce cardiovascular events, particularly hospitalization for heart failure and kidney disease.⁸⁷ Side effects of SGLT-2 inhibitors include dehydration, urinary tract infections, genital

yeast infections, acute kidney injury and increased LDL-C. More severe side effects such as urosepsis and ketoacidosis have also been noted, as well as an association with increased diabetic foot amputations.⁸⁸

GLP-1 receptor agonists (GLP-1 RA): Oral and injectable hypoglycemic agents such as semaglutide (has both oral and injectable forms), dulaglutide, exenatide, liraglutide. GLP-1 is an incretin that stimulates release of insulin from pancreatic beta cells in response to glucose. It has also been shown to slow gastric emptying, reduce food intake, and inhibit inappropriate post-meal glucagon release (refs). Injectable GLP-1 RA can be administered twice daily, daily or weekly; oral semaglutide is taken daily. GLP-1 RA have been shown to lower HbA1c (0.8-1.6%), weight (1-3 kg), blood pressure and lipids. Additionally, cardiovascular outcomes trials have demonstrated cardiovascular and renal benefits for some GLP-1 RA.⁸ The most prominent side effects are gastrointestinal, and hypoglycemia is rare. There is a risk of developing thyroid C-tumors, though this has not been seen in clinical practice.⁸⁹

Glucose-dependent insulintropic polypeptide/GLP-1 receptor agonist (GIP/GLP-1 RA): Dual agonist targeting two incretins, GIP and GLP-1, both of which are released after meals to facilitate insulin secretion. Tirzepatide, an injectable agent, is currently the only member of this class. The dual receptor agonism leads to a decrease in blood sugar through an increase in insulin secretion and increase in insulin sensitivity in the liver and skeletal muscles. Both GLP-1 and GIP have additional impacts on other tissues, as depicted in Figure A1. For example, both GLP-1 and GIP mediate weight loss through effects on satiety centers in the central nervous system. GLP-1 RA additionally decreases gastric emptying, and GIP has impacts on the subcutaneous white adipose tissue.

Figure A1. Pleiotropic Effects of Dual GIP Inhibition and GLP-1 Receptor Agonism in the Treatment of T2DM.⁹⁰



Insulin: Insulin is produced in pancreatic beta cells and controls the amount of blood glucose in the bloodstream, helps promote storage of glucose in the liver, adipose tissue, and muscles, and regulates metabolism of carbohydrates, fats, and proteins. There are short, intermediate, and long-acting insulins, delivered mainly by injection, which are used alone or in combination by patients with T2DM to help control blood sugar levels. The main side effects of insulin are hypoglycemia and weight gain.

A3. Potential Cost-Saving Measures in Type 2 Diabetes

ICER includes in its reports information on wasteful or lower-value services in the same clinical area that could be reduced or eliminated to create headroom in health care budgets for higher-value innovative services (for more information, see <https://icer.org/our-approach/methods-process/value-assessment-framework/>). These services are ones that would not be directly affected by therapies for T2DM (e.g., reduction in disability), as these services will be captured in the economic model. Rather, we are seeking services used in the current management of T2DM beyond the potential offsets that arise from a new intervention. During stakeholder engagement and public comment periods, ICER encouraged all stakeholders to suggest services (including

treatments and mechanisms of care) currently used for patients with T2DM that could be reduced, eliminated, or made more efficient. No suggestions were received.

B. Patient Perspectives: Supplemental Information

B1. Methods

ICER engaged with individual patients, patient groups, including representatives from diabetes and kidney disease advocacy organizations, and clinical experts to gather information to better understand patient experiences with the T2DM. In total, we spoke with five individual patients and seven advocacy organizations via focus groups and conference calls. We also spoke with three clinical experts throughout the review process. We also reviewed research literature suggested by or provided to ICER by advocacy organizations.

Patients and advocacy groups provided information on the impact of T2DM on patients throughout the disease course, particularly concerning aspects of the disease and caregiving that are not well-reflected in the current literature. These organizations also assisted with literature review to find information that was considered for inputs into the economic model.

C. Clinical Guidelines

Clinical practice guidelines for the treatment of T2DM have been issued by several US and non-US-based organizations. These guidelines are summarized below.

The American Diabetes Association (ADA)^{2,4}

The ADA's Standards of Medical Care in Diabetes guidelines are updated yearly and include recommendations on diagnosis and treatment of T2DM and its complications. The guidelines recommend that at time of diagnosis of T2DM, all patients, except when contraindicated, should be started on metformin in addition to comprehensive lifestyle modifications (e.g., healthy eating patterns, medical nutrition therapy, regular physical activity, weight management, smoking cessation).⁷⁰ A recommended HbA1c target is less than 7.0% for most nonpregnant adults. However, the guidelines suggest accounting for patient-specific factors, including but not limited to risk of hypoglycemia, comorbidities, disease duration, and patient preference, through which a patients' individualized target HbA1c may be higher or lower than 7.0%.³ Dual pharmacologic therapy should be considered at initiation of newly diagnosed T2DM patients if their HbA1c is greater than or equal to 1.5% of the HbA1c target. These guidelines recommend a patient-centered approach to help guide selection of pharmacologic agents with considerations for comorbidities, risk of hypoglycemia, risk of side effects, cost, and impact on patient weight, along with patient preferences.

If the patient does not have chronic kidney disease (CKD), atherosclerotic cardiovascular disease (ASCVD), or concerns regarding weight management, and the HbA1c target is not achieved after three months of therapy, it is recommended to have a combination of metformin and any of six preferred medication classes which include basal insulin, DPP-4 inhibitors, GLP-1 RA, SGLT-2 inhibitors, sulfonylureas, or thiazolidinediones, dependent upon patient factors and drug-specific effects.³ When there is a compelling need to either minimize weight gain or help promote weight loss, use of either GLP-1 RA or SGLT-2 inhibitors are preferred.

For T2DM patients who also have established ASCVD or multiple ASCVD risk factors, and who do not achieve HbA1c target after three months, use of either SGLT-2 inhibitors or GLP-1 receptor agonists are recommended as part of the treatment regimen. Among T2DM patients who have established ASCVD and heart failure or are at high risk of developing heart failure, use of SGLT-2 inhibitors is preferred. For T2DM patients who also have CKD, use of GLP-1 RA or SGLT-2 inhibitors are preferred.⁴

American Association of Clinical Endocrinology (AACE) and American College of Endocrinology (ACE)⁹¹

The AACE and ACE published a Comprehensive Type 2 Diabetes Management Algorithm in 2019. The guidelines recommend that in addition to promoting lifestyle optimization measures, clinicians should individualize both glycemic targets and choice of therapy. Choice of therapy should be patient-centered, consider ASCVD, heart failure, and CKD status, and achieved through shared decision-making. A HbA1c $\leq 6.5\%$ is considered optimal if it can be achieved in a safe and affordable manner and glycemic therapy should be evaluated frequently (e.g., every 3 months) so that glycemic targets should be achieved as soon as possible. Continuous glucose monitoring is highly recommended to assist patients in reaching glycemic targets.

Choice of therapy is based on comorbidities and HbA1c status at initiation of therapy. For patients with established ASCVD or high risk of ASCVD, CKD stage 3, or heart failure with reduced ejection fraction, a long-acting GLP-1 RA or SGLT-2 inhibitor with proven efficacy in these conditions is preferred. For patients with a HbA1c $\geq 7.5\%$ -9.0% or HbA1c $\geq 9.0\%$ without symptoms at initiation of therapy, dual therapy with metformin and another agent is recommended, with the addition of a third agent if glycemic control is not achieved within 3 months. For patient with a HbA1c $> 9.0\%$ with symptoms, insulin with or without other agents is preferred.

The European Society of Cardiology (ESC) and the European Association of the Study of Diabetes (EASD)⁹²

The ESC and EASD guidelines on diabetes, pre-diabetes, and CVD recommend the use of metformin along with lifestyle modifications (e.g., healthy eating patterns, regular physical activity, smoking cessation, weight management) as first-line therapy in patients with T2DM without established ASCVD or at high CV risk.⁹² A recommended HbA1c target is less than 7.0% for most adults, however, target goals should be individualized on a per-patient basis.

For patients with T2DM and CVD or at high CV risk, the use of GLP-1 receptor agonists or SGLT-2 inhibitors are recommended to reduce the risk of CV events. For patients with T2DM and heart failure, SGLT-2 inhibitors are recommended to reduce the risk of hospitalization from heart failure. Saxagliptin is not recommended for use in patients with heart failure. For patients with T2DM and CKD, SGLT-2 inhibitors are recommended to reduce progression of CKD.

National Institute for Health and Care Excellence (NICE)⁹³

NICE published guidelines for Type 2 diabetes in adults in 2015, and updated the guidelines in 2019. Along with evidence-based patient education delivered by trained educators, a personalized diabetes management plan including advice about diet, exercise, and weight loss, the NICE guideline recommends metformin as initial drug treatment. HbA1c targets should be individualized, based on risk of hypoglycemia (HbA1c \leq 6.5% in patients who are not on drugs that cause hypoglycemia and \leq 7.0% if patients are on drugs that cause hypoglycemia). HbA1c targets can be relaxed in patients for whom tight glucose control is not appropriate.

If metformin is not sufficient to achieve the glycemic target, the guidelines recommend dual therapy with the addition of a sulfonylurea, DPP-4 inhibitor, pioglitazone or SGLT-2 inhibitor. Triple therapy with metformin and two of the following: DPP-4 inhibitor, sulfonylurea, and/or pioglitazone, or starting insulin is recommended as a second intensification step. GLP-1 RA are recommended only if metformin + two other oral drugs is not effective, particularly in patients who are obese or have obesity-related complications, or have relative contraindications to insulin (e.g., patients in whom insulin therapy would have significant occupational implications).

D. Comparative Clinical Effectiveness:

Supplemental Information

D1. Detailed Methods

PICOTS

Population

Our intervention of interest for this review was injectable tirzepatide (Eli Lilly) added to background therapy (metformin +/- sulfonylureas or thiazolidinediones).

Comparators

We compared tirzepatide to background therapy and each of the following add-on agents:

- Semaglutide (Ozempic®, Novo Nordisk), an injectable GLP-1 receptor agonist
- Empagliflozin (Jardiance®, Boehringer Ingelheim and Eli Lilly), an oral SGLT-2 inhibitor

Outcomes

The outcomes of interest are described in the list below.

- Glycated hemoglobin (HbA1c) levels
- Fasting plasma glucose
- Body weight
- Waist circumference
- Blood pressure
- Percentage of patients achieving HbA1C targets of <7.0%, ≤6.5%, and/or <5.7%
- Lipid profile (e.g., Total Cholesterol, Low-Density Lipoprotein Cholesterol, High-Density Lipoprotein Cholesterol, and Triglycerides)
- Use of rescue medication (e.g., additional glucose-lowering medication)
- Health-related quality of life and activities of daily living (e.g., Diabetes Treatment Satisfaction Questionnaire (DTSQ), EuroQol 5-Dimensions Health-Related Quality of Life questionnaire (EQ-5D), Impact of Weight on Quality of Life Questionnaire–Lite (IWQOL-Lite), Short Form Health Survey (SF-36))
- Macrovascular outcomes including:
 - All-cause mortality
 - Cardiovascular mortality
 - Stroke

- Myocardial infarction
 - Heart failure requiring hospitalization or an urgent heart failure visit
- Microvascular outcomes including:
 - Retinopathy
 - Nephropathy
 - Neuropathy
- Adverse events including:
 - Hypoglycemia
 - Pancreatitis
 - Urogenital infections
 - Gastrointestinal effects
 - Fractures
 - Discontinuation (all-cause, due to adverse events)
 - Serious adverse events including death

Timing

Evidence on intervention effectiveness and harms was derived from studies of at least 3 months' duration.

Settings

All relevant settings were considered, with a focus on outpatient settings in the United States.

Study Design

Randomized controlled trials and non-randomized controlled trials with any sample size were included.

Table D1.1 PRISMA 2009 Checklist

		Checklist Items
TITLE		
Title	1	Identify the report as a systematic review, meta-analysis, or both.
ABSTRACT		
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.
INTRODUCTION		
Rationale	3	Describe the rationale for the review in the context of what is already known.
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).
METHODS		
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).

Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I ²) for each meta-analysis.
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.
RESULTS		
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).
DISCUSSION		
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., health care providers, users, and policy makers).
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.
FUNDING		
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.

From: Moher D, Liberati A, Tetzlaff J, Altman DG. The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097

Data Sources and Searches

Procedures for the systematic literature review assessing the evidence on new therapies for T2DM followed established best research methods.^{94,95} We conducted the review in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.⁹⁶ The PRISMA guidelines include a checklist of 27 items, which are described further in Table D1.

We searched MEDLINE, EMBASE, Cochrane Database of Systematic Reviews, and Cochrane Central Register of Controlled Trials for relevant studies. Each search was limited to English-language studies of human subjects and excluded articles indexed as guidelines, letters, editorials, narrative reviews, case reports, or news items. We included abstracts from conference proceedings identified from the systematic literature search. All search strategies were generated utilizing the Population, Intervention, Comparator, and Study Design elements described above. The proposed search strategies included a combination of indexing terms (MeSH terms in MEDLINE and Emtree terms in EMBASE), as well as free-text terms.

To supplement the database searches, we performed manual checks of the reference lists of included trials and systematic reviews and invited key stakeholders to share references germane to the scope of this project. We also supplemented our review of published studies with data from conference proceedings, regulatory documents, information submitted by manufacturers, and other grey literature when the evidence met ICER standards (for more information, see <https://icer.org/policy-on-inclusion-of-grey-literature-in-evidence-reviews/>). Where feasible and deemed necessary, we also accepted data submitted by manufacturers “in-confidence,” in accordance with ICER’s published guidelines on acceptance and use of such data (<https://icer.org/guidelines-on-icers-acceptance-and-use-of-in-confidence-data-from-manufacturers-of-pharmaceuticals-devices-and-other-health-interventions/>).


Table D1.2. Ovid MEDLINE(R) Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Ovid MEDLINE(R) Daily, Ovid MEDLINE and Versions(R) 1946 to Present, and EBM Reviews - Cochrane Central Register of Controlled Trials

	Search Terms
1	exp Diabetes Mellitus, Type 2/
2	((adult or ketosis-resistant or matur* or late or "non-insulin depend*" or "noninsulin depend*" or slow or stable or "type 2" or "type II" or lipoatrophic) adj3 diabet*) or T2D* or NIDDM).ti,ab.
3	1 or 2
4	(tirzepatide or "LY3298176" or LY3298176).ti,ab.
5	(semaglutide or "nn 9535" or nn9535 or ozempic).ti,ab.
6	(empagliflozin or "BI 10773" or BI10773 or jardiance).ti,ab.
7	4 or 5 or 6
8	3 and 7
9	(address or autobiography or bibliography or biography or case reports or clinical trial phase i or comment or conference review or congress or consensus development conference or duplicate publication or dictionary or directory or editorial or guideline or interview or lecture or legal case or legislation or letter or meta analysis or news or newspaper article or note or patient education handout or periodical index or personal narrative or portrait or practice guideline or review or systematic review or video-audio media).pt.
10	conference abstract.pt.
11	8 not (9 or 10)
12	(clinical and trial).ti,ab. or exp 'clinical trials as topic'/ or clinical trial.pt. or random*.ti,ab. or exp 'random allocation'/ or tu.xs
13	11 and 12
14	limit 13 to english language
15	(animals not (human and animals)).sh.
16	14 not 15
17	remove duplicates from 16

Table D1.3. Search Strategy of EMBASE SEARCH

	Search Terms
1	'non insulin dependent diabetes mellitus'/exp OR 'non insulin dependent diabetes mellitus'
2	((('adult OR 'ketosis resistant' OR matur* OR late OR 'non-insulin depend*' OR 'noninsulin depend*' OR slow OR stable OR 'type 2' OR 'type ii' OR lipoatrophic) NEAR/3 diabet*):ti,ab) OR t2d*:ti,ab OR niddm:ti,ab
3	#1 or #2
4	'tirzepatide'/exp
5	'tirzepatide':ti,ab OR 'LY3298176':ti,ab OR 'LY3298176':ti,ab
6	'semaglutide'/exp
7	Semaglutide:ti,ab OR 'nn 9535':ti,ab OR 'nn9535':ti,ab OR 'ozempic':ti,ab
8	'empagliflozin'/exp
9	empagliflozin:ti,ab OR bi10773:ti,ab OR 'bi 10773':ti,ab OR 'jardiance':ti,ab
10	#4 OR #5 OR #6 OR #7 OR #8 OR #9
11	#3 AND #10
12	#11 NOT ('animal experiment'/de OR 'animal model'/de OR 'case report'/de OR 'human cell'/de OR 'human tissue'/de OR 'in vitro study'/de OR 'meta analysis'/de OR 'meta analysis (topic)'/de OR 'network meta-analysis'/de OR 'nonhuman'/de OR 'phase 1 clinical trial (topic)'/de OR 'practice guideline'/de OR 'questionnaire'/de OR 'systematic review'/de OR 'systematic review (topic)'/de OR 'chapter'/it OR 'conference review'/it OR 'editorial'/it OR 'letter'/it OR 'note'/it OR 'review'/it OR 'short survey'/it)
13	'clinical':ti,ab AND 'trial':ti,ab OR 'clinical trial'/exp OR random* OR 'drug therapy':lnk OR 'clinical article'/exp OR 'controlled study'/exp OR 'major clinical study'/exp
14	#12 AND #13
15	#14 AND [english]/lim
16	('animal'/exp OR 'nonhuman'/exp OR 'animal experiment'/exp) NOT 'human'/exp
17	#15 NOT #16
18	#17 NOT [medline]/lim

Figure D1.1 PRISMA Flow Chart Showing Results of Literature Search for Tirzepatide for Type 2 Diabetes



Study Selection

We performed screening at both the abstract and full-text level. Two investigators screened all abstracts identified through electronic searches according to the inclusion and exclusion criteria described earlier. We did not exclude any study at abstract-level screening due to insufficient information. For example, an abstract that did not report an outcome of interest would be accepted for further review in full text. We retrieved the citations that were accepted during abstract-level screening for full text appraisal. Two investigators reviewed full papers and provided justification for exclusion of each excluded study. All literature that did not undergo a formal peer review process is described separately.

Tirzepatide

There are four tirzepatide trials, two Phase 2 trials that compare tirzepatide to background therapy and two Phase 3 trials that compare tirzepatide against injectable semaglutide and tirzepatide against insulin glargine.

NMA Linkages

A total of four references relating to four RCTs were used as linkages for the NMA. Two RCTs related to oral semaglutide, one RCT related to injectable semaglutide, and one RCT contained a placebo and sitagliptin arm.

CVOTs

A total of three references relating to two RCTs evaluating cardiovascular outcomes. One RCT reviewed empagliflozin and the second reviewed injectable semaglutide.

Data Extraction and Quality Assessment

We used criteria published by the US Preventive Services Task Force (USPSTF) to assess the quality of RCTs and comparative cohort studies, using the categories “good,” “fair,” or “poor” (see Table D4.1).⁹⁷ Guidance for quality ratings using these criteria is presented below, as is a description of any modifications we made to these ratings specific to the purposes of this review.

Good: *Meets all criteria: Comparable groups are assembled initially and maintained throughout the study; reliable and valid measurement instruments are used and applied equally to the groups; interventions are spelled out clearly; all important outcomes are considered; and appropriate attention is paid to confounders in analysis. In addition, intention to treat analysis is used for RCTs.*

Fair: *Studies were graded "fair" if any or all of the following problems occur, without the fatal flaws noted in the "poor" category below: Generally comparable groups are assembled initially but some question remains whether some (although not major) differences occurred with follow-up; measurement instruments are acceptable (although not the best) and generally applied equally; some but not all important outcomes are considered; and some but not all potential confounders are addressed. Intention to treat analysis is done for RCTs.*

Poor: *Studies were graded "poor" if any of the following fatal flaws exists: Groups assembled initially are not close to being comparable or maintained throughout the study; unreliable or invalid measurement instruments are used or not applied equally among groups (including not masking outcome assessment); and key confounders are given little or no attention. For RCTs, intention to treat analysis is lacking.*

Note that case series are not considered under this rating system – because of the lack of comparator, these are generally considered to be of poor quality.

Assessment of Level of Certainty in Evidence

We used the [ICER Evidence Rating Matrix](#) to evaluate the level of certainty in the available evidence of a net health benefit among each of the interventions of focus.^{98,99}

Assessment of Bias

As part of our quality assessment, we evaluated the evidence base for the presence of potential publication bias. We performed an assessment of publication bias for tirzepatide, semaglutide, and empagliflozin using the clinicaltrials.gov database of trials. We searched for studies which would have met our inclusion criteria and for which no findings have been published and did not find any evidence of publication bias.

Data Synthesis and Statistical Analyses

Data on relevant outcomes were summarized in evidence tables (see [Section D4](#)) and synthesized qualitatively in the body of the review. In addition, we evaluated the comparative efficacy of tirzepatide, injectable semaglutide, empagliflozin, and background therapy by means of network meta-analysis (NMA), where feasible. Based on data availability, our NMA evaluated HbA1c, body weight, LDL, and SBP outcomes at 40 weeks. Network Meta-Analysis Supplemental Information below ([Section D2](#)) contains a detailed description of the NMA methods. Due to inconsistent or limited data reporting, other outcomes were only described narratively in the body of the report or in [Section D4](#) of the Report Supplement.

D2. Network Meta-Analysis: Supplemental Information

NMA Methods

We evaluated the feasibility of conducting a quantitative synthesis by exploring the differences in study populations, study design, analytic methods, and outcome assessment for each outcome of interest. Trials deemed sufficiently similar in terms of population, intervention type, duration, and outcome definitions were included in the NMAs. Based on data availability, we developed quantitative, indirect comparisons of tirzepatide, injectable semaglutide, empagliflozin, and background therapy using a Bayesian NMA for outcomes of change in HbA1c, weight, LDL, and SBP at 40 weeks in adult patients with T2DM. The primary endpoints of the tirzepatide trial, SURPASS-2 was measured at 40 weeks, PIONEER-2 and 3 were measured at 52 weeks, SUSTAIN-2 at 56 weeks, and HARMONY-3 at 104 weeks. We received academic-in-confidence outcomes data at week 40 from manufacturers of four of the five trials in the NMA. We were unable to access week 40 data

from the HARMONY-3 trial. Thus, we used digitized estimates from published figures to calculate change in HbA1c and weight at week 40; we were limited to using change from baseline to week 104 for the LDL and SBP outcomes. For the outcomes of HbA1c and weight, results were reported using the treatment-regimen or equivalent estimand. For the outcome of LDL, results were reported using the efficacy estimand for SURPASS-2 and treatment-regimen estimand equivalent for SUSTAIN-2, PIONEER-2 and 3.

All four outcomes were analyzed as continuous outcomes using a generalized linear model with identity link (Tables D2.1-2.3). Vague priors were used for study-specific baselines and basic parameters in the Bayesian NMA models. However, between-study heterogeneity could not be precisely estimated in the random effect models with noninformative priors due to the small number of studies available. This was reflected by the wide credible intervals of the summary effects (data available upon request). We considered applying fixed-effect models; however, the fixed-effect model does not account for variation in intervention effects across studies. Therefore, we used random-effects models for all four outcomes and made assumptions about the extent of heterogeneity. We used the information provided by Rhodes et al. to construct informative priors for the between-study variance (τ^2).¹⁰⁰ ([see Table D2.1](#)) All NMAs were conducted using the IndiRect NMA platform (CRG-EVERSANA, 2020TM). We initially discarded the first 10,000 iterations as “burn-in” and based inferences on an additional 50,000 iterations using three chains. We evaluated the convergence of chains through visual examination of the Brook–Gelman–Rubin diagnostic and historical plots.

Table D2.1. NMAs Conducted & Presented

Outcome	Model	Number of trials
Change from Baseline in HbA1c (%)	Generalized linear model with identity link; Random effect model with informative prior (prior for between study variance: $\log t[-3.68, 2.78^2, 5]$) ¹⁰⁰	5
Change from Baseline in Body Weight (kg)	Generalized linear model with identity link; Random effect with informative prior (prior for between study variance: $\log t[-3.44, 2.44^2, 5]$) ¹⁰⁰	
Change from Baseline in LDL (mg/dL)	Generalized linear model with identity link; Random effect model with informative prior (prior for between study variance: $\log t[-3.68, 2.78^2, 5]$) ¹⁰⁰	
Change from Baseline in SBP (mmHg)	Generalized linear model with identity link; Random effect with informative prior (prior for between study variance: $\log t[-3.44, 2.44^2, 5]$) ¹⁰⁰	

%; percentage point, HbA1c: hemoglobin A1c, kg: kilogram, LDL: low-density lipoprotein, mg/dL: milligram/deciliter, mmHg: millimeters of mercury, SBP: systolic blood pressure

Table D2.2. Data Inputs for NMA of HbA1c and Body Weight Loss

		HbA1C (%)			Body Weight (kg)		
Trial	Name	N	Mean	Standard Error	N	Mean	Standard Error
SURPASS-2	TZP		-2.3			-11.2	
	SEM		-1.86			-5.7	
SUSTAIN 2	SEM						
	SITA						
PIONEER 2	OSEM						
	EMPA						
PIONEER 3	OSEM						
	SITA						
HARMONY 3*	PBO	101	0	0.13	101	-0.7	1.9
	SITA	302	-0.5	0.07	302	-0.5	1.0

?: percentage point, HbA1c: hemoglobin A1c, EMPA: empagliflozin, kg: kilogram, N: number, OSEM: oral semaglutide, PBO: placebo, SEM: semaglutide, SITA: sitagliptin, TZP: tirzepatide

*Values from HARMONY 3 were digitized by ICER staff

Table D2.3. Data Inputs for NMA of LDL and SBP

		LDL Cholesterol (mg/dL)			Systolic Blood Pressure (mmHg)		
Trial	Name	N	Mean	Standard Error	N	Mean	Standard Error
SURPASS-2	TZP		-4.5			-6.5	
	SEM		-5.6			-3.6	
SUSTAIN 2	SEM						
	SITA						
PIONEER 2	OSEM						
	EMPA						
PIONEER 3	OSEM						
	SITA						
HARMONY 3*	PBO	101	-1.2	2.7	101	2.2	1.39
	SITA	302	-1.9	1.34	302	0.2	0.85

EMPA: empagliflozin, LDL: low-density lipoprotein, mg/dL: milligrams per deciliter, N: number, OSEM: oral semaglutide, PBO: placebo, SBP: systolic blood pressure, SEM: semaglutide, SITA: sitagliptin, TZP: tirzepatide

*Note: For LDL and SBP, mean values of change were calculated from baseline to week 104

Table D2.4. NMA Results for Change in HbA1c, Body Weight, LDL and SBP at Week 40 Between Tirzepatide vs. Background Therapy/Empagliflozin

	TZP vs. BT	TZP vs. EMPA
	Mean Difference (95% Credible Interval)	
Change in HbA1c, %,	-1.72 (-1.95 to -1.49)	-1.12 (-1.39 to -0.85)
Change in Body Weight, kg	-9.51 (-10.3 to -8.73)	-7.24 (-8.08 to -6.41)
Change in LDL, mg/dL	-4.34 (-5.42 to -3.26)	-7.53 (-8.67 to -6.38)
Change in SBP, mmHg	-7.46 (-8.11 to -6.82)	-2.59 (-3.29 to -1.90)

%, percentage point, BT: background therapy, EMPA: empagliflozin, HbA1c: hemoglobin A1c, kg: kilogram, LDL: low-density lipoprotein, mg/dL: milligram per deciliter, mmHg: millimeters of mercury, SBP: systolic blood pressure, TZP: tirzepatide

D3. Additional Clinical Evidence

Trials of Tirzepatide

We identified four relevant trials of tirzepatide for treatment of T2DM.^{6,7,33,35} The key trials are described in detail below and additional details can be found in Evidence Table D4.3-6. Frias 2018, Frias 2020, and SURPASS-2 have been published and the data for these trials are informed by the clinical trial report. We also identified SURPASS-4 which includes an assessment of the cardiovascular safety of tirzepatide compared with insulin glargine.

Phase 2 (Frias 2018 and 2020)

Frias 2018 and 2020 are two phase 2 trials exploring the efficacy and safety of tirzepatide in patients with T2DM. Frias 2018 was a 26-week randomized, double-blind study where participants were randomized 1:1:1:1 to 1 mg (n=52), 5 mg (n=55), 10 mg (n=51), 15 mg (n=53) of tirzepatide, 1.5 mg of dulaglutide (n=52), or placebo (n=51). For the purposes of this review, only the 5, 10, and 15 mg tirzepatide arms were included as Eli Lilly is not seeking FDA approval for the 1 mg dose. Patients were eligible for the study if they were 18-75 years old with T2DM for at least six months, that was inadequately controlled with diet and exercise alone or with stable metformin therapy for at least 3 months before screening, and a BMI of 23-50 kg/m². The primary outcome was change in HbA1c from baseline at 26 weeks. Secondary outcomes include change in HbA1c at week 12, change in mean bodyweight, and waist circumference from baseline to weeks 12 and 26.

Frias 2020 was a phase 2 dose-ranging study, where patients were randomized to either placebo or one of three tirzepatide doses. For the purpose of this review, the 12 mg arm was not reported on, as Eli Lilly will not be seeking FDA approval for this dose. The two 15 mg arms had different dose titration regimens; the 15 mg-1 group was 2.5 mg for two weeks followed by 5 mg for 2 weeks, 10 mg for 4 weeks and then 15 mg for the final 4 weeks. The 15 mg-2 arm was 2.5 mg for 4 weeks,

followed by 7.5 mg for 4 weeks, and then 15 mg for the final 4 weeks. Inclusion criteria and outcomes were similar to the 2018 study. The timepoint of interest was 12 weeks.

Additional baseline characteristics are available in [Evidence Table D4.3](#).

SURPASS-2

SURPASS-2 was a phase 3, head-to-head open-label trial exploring the efficacy and safety of tirzepatide compared to injectable semaglutide. Patients were randomized 1:1:1:1 to tirzepatide 5 mg (n= 470), 10 mg (n= 469), 15 mg (n= 470) or semaglutide 1mg (n= 469) every week for 40 weeks followed by a 4-week safety period. 1,878 patients included in the study were adults with T2DM that were inadequately controlled with at least 1500 mg of metformin per day. Included patients also had HbA1c levels of 7.0 to 10.5% and a BMI of ≥ 25 kg/m² with a stable weight in the past three months. Patients with type 1 diabetes, an eGFR below 45 mL/min/1.73 m², and a history of pancreatitis were excluded from the study. Included patients had a mean age of 56.6 years, with 53% identifying as female, 82.6% white, an average weight of 93.7 kg, and an eGFR of 96.0 mL/min/1.73 m². Additional baseline characteristics are available in Evidence Table D4.3.

The primary endpoint was change in HbA1c from baseline to week 40. Secondary endpoints include change from body weight, and attainment of HbA1c targets of less than 7.0% and less than 5.7%.

SURPASS-4

SURPASS-4 was an open-label phase 3 study exploring the efficacy and cardiovascular safety of tirzepatide (5, 10 and 15 mg) versus insulin glargine in patients with T2DM. Patients were randomized to either a tirzepatide arm (n= 995) or insulin glargine (n= 1000) for 52-weeks for the primary efficacy endpoint, with an additional variable treatment period of up to 52 additional weeks. The median study duration was 85 weeks. Study participants were adults with T2DM inadequately controlled with any of three oral glucose-lowering medications (i.e., metformin, sulfonylurea, or an SGLT-2 inhibitor), a BMI of 25 kg/m² or more and stable weight and increased risk of CV events. Patients with type 1 diabetes or a history of pancreatitis were not included in the study. Included patients had a mean disease duration of 10.5 years, baseline HbA1c of 8.52%, and a baseline weight of 90.3 kg. 87% of patients had a history of cardiovascular disease.

The primary endpoint was change in HbA1c from baseline to 52 weeks. Key secondary endpoints include change in bodyweight at 52 and achievement of HbA1c target of less than 7%. A prespecified cardiovascular risk comparison between tirzepatide and insulin glargine was also conducted and assessed MACE-4. Additional outcomes are available in Evidence Table D4.12.

Trials of NMA Linkage Studies

We identified four trials to provide linkages for the NMA.^{39,101-103} The trials are described below and additional details can be found in [Evidence Tables D4.7-9](#). All trials are published and are informed by the respective clinical trial reports.

SUSTAIN-2

SUSTAIN-2 was a phase 3a, randomized, double blind, multicenter trial, assessing the efficacy and safety of semaglutide versus sitagliptin in patients with T2DM inadequately controlled on metformin, thiazolidinediones, or both. Included patients were adults with T2DM with insufficient glycemic control for 90 days prior to screening and were on stable treatment with metformin, pioglitazone, rosiglitazone, or metformin and rosiglitazone. Patients were excluded if they were on other glucose lowering drugs not described above, had a history of chronic or acute pancreatitis, impaired renal function, or heart failure at any time. 1,231 patients were randomized 2:2:1:1 to 0.5 mg semaglutide (n=409), 1.0 mg semaglutide (n= 409) or two arms of 100 mg sitagliptin that were pooled for analysis (n=407). Included patients had a mean age of 55.1, years, HbA1c of 64.7mmol/mol, 49.5% female, 68.5% white and an eGFR of 97.50 mL/min/1.73 m². Additional baseline characteristics are available in [Evidence Table D4.7](#).

The primary outcome was change in HbA1c from baseline at week 56. Secondary endpoints include change in bodyweight, proportion of patients who achieved an HbA1c of less than 7.0% and HbA1c of 6.5%.

PIONEER 2 and PIONEER 3

PIONEER 2 and 3 have been previously described in the 2019 T2DM ICER report.⁴⁰ A brief description is provided below.

The PIONEER program was comprised of 10 trials (PIONEER 1-10). The PIONEER trials included in this review (PIONEER 2 and 3) were multinational RCTs comparing oral semaglutide to sitagliptin, empagliflozin, liraglutide, and placebo. PIONEER 2 compared oral semaglutide 14 mg to empagliflozin 25 mg added to metformin and PIONEER 3 compared oral semaglutide 3, 7, and 14 mg to sitagliptin 100 mg added to metformin ± sulfonylurea (47%). Key exclusion criteria included: renal impairment (eGFR <60 mL/min/1.73 m²); MI, stroke, hospitalization for unstable angina, or transient ischemic attack within 180 days; stage IV heart failure; and history of pancreatitis. Baseline characteristics are available in [Evidence Table D4.7](#).

HARMONY-3

The HARMONY 3 trial was a phase 3 randomized, double-blind, placebo-controlled study and evaluated the efficacy and safety of albiglutide versus daily sitagliptin, daily glimepiride, and

placebo in patients with type 2 diabetes. For the purpose of this review, we only used the placebo and sitagliptin arms for the NMA. Patients were eligible to be enrolled if they were adults with T2DM and experienced inadequate glycemic control while taking background metformin at least 3 months before screening. Patients also had to have a HbA1c of 7.0% to 10% and a BMI of 20 to 45 kg/m². 1049 patients met the criteria (placebo, n= 101 and sitagliptin, n= 302) with a mean age of 54.3 to 56.1, and a mean duration of diabetes from 5.8 to 6.7 years. Additional baseline characteristics are available in [Evidence Table D4.7](#).

The primary outcome was change in HbA1c from baseline to week 104. The secondary endpoints include change from baseline for HbA1c and weight.

Cardiovascular Outcomes Trials (CVOT)

We identified two relevant trials exploring the efficacy of semaglutide and empagliflozin on cardiovascular outcomes.^{8,9} The trials are described below with additional details on study design available in [Evidence Table D4.2](#).

SUSTAIN-6

The SUSTAIN-6 trial was a phase 3 randomized, double-blind, placebo-controlled trial evaluating the cardiovascular safety of semaglutide in patients with T2DM. Patients were randomized 1:1:1:1 to either 0.5 or 1.0 mg injectable semaglutide or a matched placebo for 104 weeks with a 5-week follow up period. 3,297 adults met the inclusion criteria of T2DM and HbA1c levels of 7%, and no previous treatment with an antihyperglycemic drug, or more than two oral antihyperglycemic agents with or without basal or premixed insulin. Patients were also included if they were 50+ years of age with established CVD, chronic heart failure, or chronic kidney disease of stage three or higher. Included patients had a mean age of 64.6 years, an average disease duration of 13.8 years and an average Hba1c level of 8.7%. Additional baseline characteristics are available in [Evidence Table D4.10](#).

The primary composite outcome was first occurrence of death from CV causes, nonfatal MI, or nonfatal stroke (MACE-3). A key secondary outcome was first occurrence of an expanded composite CV outcome (death from CV causes, nonfatal MI, nonfatal stroke, revascularization and hospitalization for unstable angina or heart failure).

EMPA-REG -OUTCOME

The EMPA-REG OUTCOME trial was a phase 3 trial evaluating the efficacy and safety of empagliflozin added to standard of care on cardiovascular morbidity and mortality in patients with T2DM and high cardiovascular risk. Patients were eligible for the trial if they were 18 or older with T2DM, a BMI of 45 kg/m² or less and eGFR of at least 30 mL/min/1.73m² and established CVD. Included patients were randomized 1:1:1 to either two doses of EMPA (N= 4687) or placebo (N=

2133). The baseline characteristics were well balanced, with a mean age of 63.1, mean weight of 86.3 kg and a HbA1c of 8.07%. Median observation time was 3.1 years. Additional baseline characteristics are available in [Evidence table D4.10](#).

The primary outcome was MACE-3. The key secondary outcome was a composite of the primary outcome plus hospitalization for unstable angina.

D4. Evidence Tables

Table D4.1. USPSTF Study Quality

Trial	Comparable Groups	Non-differential Follow-up	Patient/ Investigator Blinding	Clear Definition of Intervention	Clear Definitions of Outcomes	Selective Outcome Reporting	Measurements Valid	Intention-to-treat Analysis	Approach to Missing Data	USPSTF Rating
Tirzepatide										
Frias 2018	Yes	Yes	Yes	Yes	Yes	No	Yes	mITT	LOCF	Good
Frias 2020	Yes	No	Yes	Yes	Yes	No	Yes	mITT	MMRM	Fair
SURPASS-2	Yes	Yes	Open-label	Yes	Yes	No	Yes	mITT	MMRM	Good
SURPASS-4	Yes	Yes	Open-label	Yes	Yes	No	Yes	mITT	MMRM	Good
Injectable Semaglutide										
SUSTAIN-2	Yes	Yes	Yes	Yes	Yes	No	Yes	mITT	MI	Good
SUSTAIN-6	Yes	Yes	Yes	Yes	Yes	No	Yes	ITT	MAR	Good
Oral Semaglutide*										
PIONEER 2	Yes	Yes	Open-label	Yes	Yes	No	Yes	ITT	MI	Good
PIONEER 3	Yes	Yes	Yes	Yes	Yes	No	Yes	ITT	MI	Good
Empagliflozin										
EMPA-REG -OUTCOME	Yes	Yes	Yes	Yes	Yes	No	Yes	mITT	LOCF	Good
Sitagliptin*										
HARMONY 3	Yes	Yes	Yes	Yes	Yes	No	Yes	ITT	LOCF	Good

Studies with an asterisk (*) were used as linkages for the NMA. ITT: intention-to-treat analysis, LOCF: last observation carried forward, MAR: missing at random, MI: multiple imputation, mITT: modified intention-to-treat analysis, MM: mixed-effects model, MMRM: mixed-effects model repeated measure, NR: not reported, USPSTF: US Preventive Services Task Force

Table D4.2. Study Design

Trial	Interventions	Background Therapy	Inclusion Criteria	Study Length	Primary Key Outcome	Key Secondary Outcomes
Tirzepatide						
Phase 2 Frias 2018 N= 318 18 to 75 years	PBO TZP 1 mg TZP 5 mg TZP 10 mg TZP 15 mg DUA 1.5 mg	+/- MET	Have had T2DM for ≥6 months according to WHO classification Have HbA1c of 7.0% to 10.5% If on MET, have been treated with stable doses of MET for at least 3 months Have BMI ≥23 and <50 kg/m ²	26 weeks	<u>Primary</u> Change in HbA1c from baseline to Week 26 Bayesian Dose Response	<u>Secondary</u> Percentage of participants with 5% or greater body weight loss from baseline Percentage of participants with 10% or greater body weight loss from baseline Percentage of participants reaching the HbA1c target of ≤6.5% Percentage of participants reaching the HbA1c target of <7.0%
Phase 2 Frias 2020 N= 111 18 to 75 years	TZP 12 mg TZP 15 mg-1 TZP 15 mg-2 PBO	+/- MET	Have T2DM for ≥ 6 months according to ADA 2017 Have HbA1c of 7.0% to 10.5% If on MET, have been treated with stable doses of MET for at least 30 months Have a BMI between 23 and 45 kg/m ²	12 weeks	<u>Primary</u> Change from baseline in HbA1c	<u>Secondary</u> Change from baseline in body weight Change from baseline in waist circumference Number of participants with anti-drug antibodies Number of episodes of total hypoglycemia episodes Pharmacokinetics (PK): Average trough concentration

						(Conctrough) of tirzepatide
<p>SURPASS-2 Frias 2021</p> <p>N= 1,879</p> <p>18 and older</p>	<p>TZP 5 mg</p> <p>TZP 10 mg</p> <p>TZP 15 mg</p> <p>SEM 1 mg</p>	MET	<p>Have been diagnosed with T2DM</p> <p>Have HbA1c between $\geq 7.0\%$ and $\leq 10.5\%$</p> <p>Be on stable treatment with unchanged dose of metformin >1500 mg/day for at least 3 months prior to screening</p> <p>Be of stable weight ($\pm 5\%$) for at least 3 months before screening</p>	40 weeks	<p><u>Primary</u></p> <p>Change from Baseline in HbA1c to week 40</p>	<p><u>Secondary</u></p> <p>Mean change from baseline in daily average 7-Point SMBG values</p> <p>Percentage of participants who achieved weight loss $\geq 5\%$</p> <p>Rate of documented symptomatic hypoglycemic episodes</p> <p>Change from baseline in body weight</p> <p>Percentage of participants achieving an HbA1c target value of $<7\%$</p>
<p>SURPASS-4 Del Proto 2021</p> <p>N= 1878</p> <p>18 and older</p>	<p>TZP 5mg</p> <p>TZP 10mg</p> <p>TZP 15mg</p> <p>Insulin glargine SC once daily</p>	OADs (MET, SGLT-2 inhibitors, and/or SU)	<p>Have been diagnosed with T2DM</p> <p>Have HbA1c between $\geq 7.5\%$ and $\leq 10.5\%$</p> <p>Be on stable treatment with unchanged dose of at least 1 and no more than 3 types of OADs, which may only include metformin, SGLT-2 inhibitors, and/or SU for at least 3 months before screening</p> <p>Have increased risk for CV events</p> <p>Be of stable weight ($\pm 5\%$)</p>	Up to 104 weeks	<p><u>Primary</u></p> <p>Change from baseline in HbA1c to week 52</p>	<p><u>Secondary</u></p> <p>Change from baseline in body weight</p> <p>Percentage of participants achieving an HbA1c target value of $<7\%$</p> <p>Change from baseline in fasting serum glucose</p> <p>Change from Baseline in HbA1c (5 mg)</p>

Injectable Semaglutide						
<p>SUSTAIN-2 Ahren 2017</p> <p>N= 1231</p> <p>18 and older</p>	<p>SEM 0.5 mg SEM 1.0 mg SITA 100 mg + PBO 0.5 mg SITA 100 mg + PBO 1.0 mg</p>	<p>MET ± TZD</p>	<p>Japan: Age minimum 20 years</p> <p>Subjects diagnosed with T2DM and on stable treatment in a period of 90 days prior to screening with either MET above or equal to 1500 mg (or maximum tolerated dose), pioglitazone above or equal to 30 mg (or maximum tolerated dose), rosiglitazone above or equal to 4 mg (or maximum tolerated dose) or a combination of either MET/pioglitazone or MET/rosiglitazone (doses as for individual therapies)</p>	<p>56 weeks</p>	<p><u>Primary</u> Change from Baseline in HbA1c</p>	<p><u>Secondary</u> Change in body weight from baseline</p> <p>Change in FPG from baseline</p> <p>Change in PRO questionnaire</p> <p>Change in systolic and diastolic blood pressure from baseline</p> <p>Subjects who achieve HbA1c below or equal to 6.5% (48 mmol/mol) target (yes/no)</p>
<p>SUSTAIN-6 Marso 2016</p> <p>N= 3260</p> <p>50 and older</p>	<p>SEM 0.5 mg SEM 1.0 mg PBO 0.5 mg SC QW PBO 1.0 mg SC QW</p>	<p>Naïve or treated with OADs or insulin</p>	<p>Men and women with T2DM</p> <p>Age above or equal to 50 years at screening and clinical evidence of cardiovascular disease or age above or equal to 60 years at screening and subclinical evidence of cardiovascular disease</p> <p>Anti-diabetic drug naïve, or treated with one or two OADs</p> <p>HbA1c above or equal to 7.0% at screening</p>	<p>148 weeks</p>	<p><u>Primary</u> First occurrence of a MACE</p>	<p><u>Secondary</u> Change from baseline: urinary albumin to creatinine ratio</p> <p>Change from baseline: vital signs</p> <p>Time from randomization to first occurrence of an expanded composite cardiovascular outcome</p> <p>Change from baseline: HbA1c</p> <p>Change from baseline: Lipid profile</p>

						Change from baseline FBG
Oral Semaglutide						
<p>PIONEER 2 NCT02893328 Rodbard et al.</p> <p>N=822</p> <p>18 and older</p>	<p>SEM 14 mg EMPA 25 mg</p>	<p>MET</p>	<p>Male or female, age above or equal to 18 years at the time of signing informed consent</p> <p>Diagnosed with T2DM at least 90 days prior to day of screening</p> <p>HbA1c of 7.0 to 10.5% (53-91 mmol/mol)</p> <p>Stable daily dose of metformin (at least 1500 mg or MTD) at least 90 days prior to the day of screening</p>	<p>52 weeks</p>	<p><u>Primary</u> Change in HbA1c (week 0 to 26)</p>	<p><u>Secondary</u> Change in body weight (kg) (week 0 to 26)</p> <p>Change in HbA1c (week 0 to 52)</p> <p>Change in body weight (kg) (week 0 to 52)</p> <p>Change in FPG</p>
<p>PIONEER 3 NCT026865 Rosenstock et al.</p> <p>N=1864</p> <p>18 and older</p>	<p>SEM 3 mg SEM 7 mg SEM 14 mg SITA 100 mg</p>	<p>MET +/- SU</p>	<p>Male or female, age at least 18 years at the time of signing informed consent</p> <p>Diagnosed with T2DM for at least 90 days prior to day of screening</p> <p>HbA1c 7.0-10.5% (53-91 mmol/mol)</p> <p>Stable daily dose of MET (at least 1500 mg or MTD) within 90 days prior to the day of screening</p>	<p>78 weeks</p>	<p><u>Primary</u> Change in HbA1c (week 0 to 26)</p>	<p><u>Secondary</u> Change in body weight (week 0 to 26)</p> <p>Change in HbA1c (weeks 0 to 52, 78)</p> <p>Change in body weight (kg) (weeks 0 to 52,78)</p> <p>Change in body weight (%)</p> <p>Change in FPG</p> <p>Change in BMI</p> <p>Change in waist circumference</p>
Empagliflozin						

EMPA- REG OUTCOME (CVOT) NCT01131676 Zinman et al. N=7064	EMPA 10 mg EMPA 25 mg PBO	Naïve or pre- treated with any background therapy	<p>Diagnosis of T2DM</p> <p>Male or female patients on diet and exercise regimen who are drug naive or pre-treated with any background therapy</p> <p>Antidiabetic therapy has to be unchanged for 12 weeks prior to randomization</p> <p>HbA1c of $\geq 7.0\%$ and $\leq 10\%$ for patients on background therapy or HbA1c $\geq 7.0\%$ and $\leq 9.0\%$ for drug naive patients</p> <p>Age ≥ 18 years</p> <p>BMI ≤ 45 kg/m² at Visit 1</p>	Up to 4.6 years	<p><u>Primary</u></p> <p>Time to the first occurrence of any of the following adjudicated components of the primary composite endpoint (MACE-3):</p> <p>CV</p> <p>Death (including fatal stroke and fatal MI)</p> <p>Non-fatal MI (excluding silent MI)</p> <p>Non-fatal Stroke</p>	<p><u>Secondary</u></p> <p>Percentage of participants with the composite of all events adjudicated (MACE-4):</p> <p>CV</p> <p>Death (including fatal stroke and fatal MI)</p> <p>Non-fatal MI (excluding silent MI)</p> <p>Non-fatal stroke</p> <p>Hospitalization for Unstable Angina Pectoris</p> <p>Percentage of participants with silent MI</p>
--	---------------------------------	--	---	-----------------------	---	--

BMI: body mass index, CVOT: cardiovascular outcomes trial, DUA: dulaglutide, EMPA: empagliflozin, FPG: fasting plasma glucose, HbA1c: hemoglobin A1c, MACE: major adverse cardiovascular event, MACE-3: 3-point MACE, MACE-4: 4-point MACE, MET: metformin, mg: milligram, MTD: maximum tolerated dose, OAD: oral antidiabetic drug, PBO: placebo, PK: pharmacokinetics, PRO: patient-reported outcomes, QW: once weekly, SC: subcutaneous, SEM: semaglutide, SGLT-2: sodium-dependent glucose co-transporter-2, SIT: sitagliptin, SMBG: self-monitoring of blood glucose, SU: sulfonylurea, TZP: tirzepatide, T2DM: type 2 diabetes mellitus

Table D4.3. Baseline Characteristics – Tirzepatide^{6,7,33,35}

Study		Frias 2018				Frias 2020			SURPASS -2				SURPASS-4				
Arm		PBO	TZP 5 mg	TZP 10 mg	TZP 15 mg	PBO	TZP 15 mg-1	TZP 15 mg-2	TZP 5 mg	TZP 10 mg	TZP 15 mg	SEM 1 mg	TZP 5 mg	TZP 10 mg	TZP 15mg	Insulin G	Overall I
N		51	55	51	53	26	28	28	470	469	470	469	329	328	338	1000	1995
Age, y	Mean (SD)	56.6 (8.9)	57.9 (8.2)	56.5 (9.9)	56.0 (7.6)	56.0 (10.13)	55.5 (8.54)	56.6 (9.21)	56.3 (10.0)	57.2 (10.5)	55.9 (10.4)	56.9 (52)	62.9 (8.6)	63.7 (8.7)	63.7 (8.6)	63.8 (8.5)	63.6 (8.6)
Sex, n (%)	Men	29 (57)	34 (62)	30 (59)	22 (42)	12 (46.2)	16 (57.1)	23 (82.1)	205 (43.6)	238 (50.7)	214 (45.5)	225 (48)	198 (60)	209 (64)	203 (60)	636 (64)	1246 (62)
	Women	22 (43)	21 (38)	21 (41)	31 (59)	NR	NR	NR	265 (56.4)	231 (49.2)	256 (54.5)	244 (52.0)	131 (40)	119 (36)	135 (40)	364 (36)	749 (38)
Diabetes Duration, y	Mean	8.6	8.9	7.9	8.5	8.8	8.2	8.9	9.1	8.4	8.7	8.3	9.8	10.6	10.4	10.7	10.5
	SD	7	5.7	5.8	6.1	6.43	4.87	6.35	7.16	5.9	6.85	5.8	6.2-15.3	6.5-16.2	5.5-15.7	6.3-16.5	6.2-15.9
HbA1c mean (SD)	%	8.0 (0.9)	8.2 (1.0)	8.2 (1.1)	8.1 (1.1)	8.2 (1.2)	8.5 (1.2)	8.4 (1.1)	8.3 (1.1)	8.3 (1.0)	8.3 (1.0)	8.3 (1.0)	8.52 (0.84)	8.59 (0.91)	8.52 (0.98)	8.50 (0.85)	8.52 (0.88)
Weight, kg	Mean	91.5	92.8	92.7	89.1	89.6	88.7	89.6	92.5	94.8	93.8	93.7	90.3	90.6	90	90.2	90.3
	SD	23.1	19	19.5	22.7	23.7	18.21	16.91	21.76	22.71	21.83	21.12	20.32	18.21	16.34	19	18.66
BMI, kg/ m ²	Mean (SD)	32.4 (6.0)	32.9 (5.7)	32.6 (5.8)	32.2 (6.2)	32.5 (5.7)	32.0 (5.56)	31.1 (4.21)	33.8 (6.85)	34.3 (6.60)	34.5 (7.11)	34.2 (7.15)	32.6 (6.06)	32.8 (5.51)	32.5 (5.02)	34.5 (5.55)	32.6 (5.54)
Waist, cm	Mean (SD)	107.7 (2.06)	110.1 (2.0)	109.6 (2.04)	107.6 (2.17)	109.1 (15.38)	107.0 (12.65)	105.1 (12.19)	108.06 (14.81)	110.55 (16.05)	109.55 (15.60)	109.04 (14.90)	NR	NR	NR	NR	NR
eGFR	Mean (SD)	95.3 (15.3)	92.2 (17.2)	93.7 (18.6)	91.8 (17.9)	NR	NR	NR	96.6 (17.51)	95.5 (16.62)	96.3 (16.92)	95.6 (17.25)	80.3 (22.66)	81.4 (20.44)	81.6 (21.22)	81.5 (20.78)	81.3 (21.11)
Metformin use, N (%)	Yes	47 (92.2)	49 (89.1)	44 (86.3)	51 (96.2)	23 (88.5)	25 (89.3)	23 (82.1)	470 (100)	469 (100)	470 (100)	469 (100)	306 (93)	316 (96)	317 (94)	954 (95)	1893 (95)
Ethnicity, N (%)	Hispanic	27 (59)	22 (49)	26 (57)	23 (46)	NR	NR	NR	325 (69.1)	322 (68.7)	334 (71.1)	336 (71.6)	NR	NR	NR	NR	NR
	Non-Hispanic	19 (41)	23 (51)	20 (44)	27 (54)	NR	NR	NR	145 (30.9)	147 (31.3)	136 (28.9)	133 (28.4)	NR	NR	NR	NR	NR
Race, N (%)	American Indian/	NR	NR	NR	NR	NR	NR	NR	53 (11.3)	53 (11.3)	57 (12.1)	45 (9.6)	NR	NR	NR	NR	NR

	Alaska Native																
	Black	2 (4)	6 (11)	7 (14)	6 (11)	NR	NR	NR	28 (6.0)	21 (4.5)	15 (3.2)	15 (3.2)	13 (4)	17 (5)	11 (3)	32 (3)	73 (4)
	Asian	1 (2)	0	1 (2)	1 (2)	NR	NR	NR	6 (1.3)	11 (2.3)	5 (1.1)	3 (0.6)	15 (5)	16 (5)	8 (2)	31 (3)	70 (4)
	White	41 (80)	46 (84)	37 (74)	43 (81)	NR	NR	NR	382 (81.3)	376 (80.2)	334 (71.1)	336 (71.6)	260 (79)	259 (79)	285 (85)	825 (83)	1629 (82)

A dulaglutide arm and 1mg tirzepatide arm in Frias 2018 and 12mg Tirzepatide arm in Frias 2020 are also available. Frias 2020 has two 15-mg arms following two different dosing schedules. N: number, NR: not reported, PBO: placebo, SD: standard deviation, TZP: tirzepatide

Table D4.4. Outcomes– Tirzepatide Phase 2 Trials ^{6,33}

Study		Frias 2018				Frias 2020			
Arm		PBO	TZP 5 mg	TZP 10 mg	TZP 15 mg	PBO	TZP 12 mg	TZP 15 mg-1	TZP 15 mg-2
N		51	55	51	53	26	29	28	28
Timepoint	26 wks					12 wks			
Glycemia endpoints									
HbA1c, %	Mean, Change from BL	0.1	-1.6	-2	-2.4	0.2	-1.7	-2	-1.8
	ETD	NR	NR	NR	NR	REF	-1.9	-2.2	-2
	95% CI (ETD)	REF	-1.88 to -1.46	-2.04 to -1.61	-2.11 to -1.67	REF	-2.5 to -1.4	-2.8 to -1.7	-2.5 to -1.4
	P-Value (ETD)	REF	<0.0001	<0.0001	<0.0001	REF	<0.001	<0.001	<0.001
Lipid Levels									
Triglycerides, mg/dL	Change from BL	26.6	-44.3	-62	-70.9	NR	NR	NR	NR
Total Cholesterol, mg/dL	Change from BL	11.6	-3.9	-11.6	-11.6	NR	NR	NR	NR
HDL Cholesterol, mg/dL	Change from BL	0	0	0	3.9	NR	NR	NR	NR
LDL Cholesterol, mg/dL	Change from BL	7.7	0	0	-3.9	NR	NR	NR	NR
Body Weight Endpoints									
Weight, kg	Change from BL	-0.4	-4.8	-8.7	-11.3	-0.5	-5.3	-5.5	-5.7
	ETD	NR	NR	NR	NR	REF	-4.8	-5	-5.2
	95% CI	NR	NR	NR	NR	REF	-7.1 to -2.6	-7.2 to -2.7	-7.5 to -2.9
	P-Value	NR	NR	NR	NR	REF	<0.001	<0.001	<0.001
Mean Waist Circumference	Change from BL, cm	-1.3	-5.1	-7.4	-10.2	-2.5	-4.8	-4.9	-4.9
	ETD	NR	NR	NR	NR	REF	-2.2	-2.4	-2.4
	95% CI	NR	NR	NR	NR	REF	-4.7 to 0.2	-4.9 to 0.1	-4.9 to 0.2
	P-Value	NR	NR	NR	NR	REF	0.075	0.065	0.065

Blood Pressure and Pulse Rate									
DBP, mmHg	Change from BL	0.8 (1.26)	-0.7 (1.18)	-0.2 (1.22)	-0.7 (1.27)	NR	NR	NR	NR
	ETD	NR	NR	NR	NR	NR	NR	NR	NR
	95% CI	NR	NR	NR	NR	NR	NR	NR	NR
	P-Value	NR	NR	NR	NR	NR	NR	NR	NR
SBP, mmHg	Change from BL	1.7 (1.98)	-2.6 (1.86)	-1.3 (1.92)	-1.0 (2.00)	NR	NR	NR	NR
	ETD	NR	NR	NR	NR	NR	NR	NR	NR
	95% CI	NR	NR	NR	NR	NR	NR	NR	NR
	P-Value	NR	NR	NR	NR	NR	NR	NR	NR
Achieving HbA1C targets									
<7%	N (%)	6 (11.8)	38 (69.1)	45 (90)	41 (77.4)	NR	NR	NR	NR
	OR (95% CI)	REF	20.98 (7.28 to 60.50)	89.44 (24.41 to 327.76)	33.63 (11.02 to 102.62)	NR	NR	NR	NR
	P value	REF	<0.0001	<0.0001	<0.0001	NR	NR	NR	NR
<6.5%	N (%)	1 (2)	35 (63.6)	41 (82)	31 (58.5)	NR	NR	NR	NR
	OR (95% CI)	REF	73.64 (12.92 to 419.77)	200.87 (32.74 to >999)	56.95 (10.01 to 323.97)	NR	NR	NR	NR
	P value	REF	<0.0001	<0.0001	<0.0001	NR	NR	NR	NR
<5.7%	N (%)	1 (2)	2 (3.6)	9 (18)	16 (30.2)	NR	NR	NR	NR
	OR (95% CI)	REF	1.60 (0.21 to 12.26)	7.75 (1.33 to 44.97)	14.42 (2.61 to 79.75)	NR	NR	NR	NR
	P value	REF	0.6506	0.0225	0.0022	NR	NR	NR	NR
Achieving bodyweight targets									
≥5%	N (%)	0 (0)	26 (47.3)	36 (70.6)	33 (62.3)	NR	NR	NR	NR
	OR (95% CI)	REF	90.91 (5.45 to >999)	236.84 (13.99 to >999)	161.12 (9.62 to >999)	NR	NR	NR	NR
	P value	REF	0.0017	0.0002	0.0004	NR	NR	NR	NR
≥10 %	N (%)	0 (0)	9 (16.4)	20 (39.2)	20 (37.7)	NR	NR	NR	NR

	OR (95% CI)	REF	20.82 (1.22 to 355.42)	67.57 (4.07 to >999)	66.20 (3.99 to >999)	NR	NR	NR	NR
	P value	REF	0.036	0.0033	0.0034	NR	NR	NR	NR
≥15 %	N (%)	0 (0)	3 (5.5)	11 (21.6)	13 (24.5)	NR	NR	NR	NR
	OR (95% CI)	REF	NR	NR	NR	NR	NR	NR	NR
	P value	REF	NR	NR	NR	NR	NR	NR	NR

A dulaglutide arm in Frias 2018 and 12 mg Tirzepatide arm in Frias 2020 are also available. Frias 2020 has two 15-mg arms following two different dosing schedules. BL: baseline, CI: confidence interval, DBP: diastolic blood pressure, ETD: estimated treatment difference, N: number, NR: not reported, OR: odds ratio, PBO: placebo, REF: reference, SBP: systolic blood pressure, SD: standard deviation, TZP: tirzepatide

Table D4.5. Outcomes– Tirzepatide Phase 3 Trials^{7,35}

Study		SURPASS-2				SURPASS-4			
Arm		TZP 5 mg	TZP 10 mg	TZP 15 mg	SEM 1 mg	TZP 5 mg	TZP 10 mg	TZP 15 mg	Insulin Glargine
N		470	469	470	469	328	326	337	998
Timepoint		40 wks				52 wks			
Glycemia endpoints									
HbA1c, %	Mean, Change from BL	-2.01	-2.24	-2.3	-1.86	-2.24	-2.43	-2.58	-1.44
	ETD	-0.15	-0.39	-0.45	REF	-0.80	-0.99	-1.14	REF
	95% CI (ETD)	-0.28 to -0.03	-0.51 to -0.26	-0.57 to -0.32	REF	-0.92 to -0.68	-1.11 to -0.87	-1.26 to -1.02	REF
	P-Value (ETD)	0.02	<0.001	<0.001	REF	<0.0001	<0.0001	<0.0001	REF
Lipid levels, mg/dL									
Triglycerides	Change from BL	-31.4*	-40*	-41.1*	-19.1*	-16.3	-20.1	-22.5	-6.4
Total Cholesterol	Change from BL	-9.4*	-10.2*	-10.7*	-8.2*	-5.2	-5.5	-5.6	0
HDL Cholesterol	Change from BL	2.9*	3.4*	3*	1.9*	6.7	9.7	10.8	2.9
LDL Cholesterol	Change from BL	-6.7*	-4.9*	-4.5*	-5.6*	-6.8	-8.3	-7.9	1.4

Study		SURPASS-2				SURPASS-4			
Arm		TZP 5 mg	TZP 10 mg	TZP 15 mg	SEM 1 mg	TZP 5 mg	TZP 10 mg	TZP 15 mg	Insulin Glargine
Body Weight Endpoints									
Weight, kg	Change from BL	-7.6	-9.3	-11.2	-5.7	-7.1	-9.5	-11.7	1.9
	ETD	-1.9	-3.6	-5.5	REF	-9.0	-11.4	-13.5	REF
	95% CI	-2.8 to -1.0	-4.5 to -2.7	-6.4 to -4.6	REF	-9.8 to -8.3	-12.1 to -10.6	-12.1 to -10.6	REF
	P-Value	<0.001	<0.001	<0.001	REF	<0.0001	<0.0001	<0.0001	REF
Mean Waist Circumference, cm	Change from BL	-6.9*	-9.6*	-9.9*	-5.6*	-8.41†	-8.27†	-9.46†	1.4†
	ETD	NR	NR	NR	NR	NR	NR	NR	NR
	95% CI	NR	NR	NR	NR	NR	NR	NR	NR
	P-Value	NR	NR	NR	NR	NR	NR	NR	NR
Blood pressure and pulse rate									
DBP, mmHg	Change from BL	-4.5	-5.3	-6.5	-3.6	1†	0.94†	0.8†	-0.72†
	ETD	NR	NR	NR	NR	NR	NR	NR	NR
	95% CI	NR	NR	NR	NR	NR	NR	NR	NR
	P-Value	NR	NR	NR	NR	NR	NR	NR	NR
SBP, mmHg	Change from BL	NR	NR	NR	NR	-1.65†	-4.33†	-4.59†	-1.2†
	ETD	NR	NR	NR	NR	NR	NR	NR	NR
	95% CI	NR	NR	NR	NR	NR	NR	NR	NR
	P-Value	NR	NR	NR	NR	NR	NR	NR	NR
Achieving HbA1c targets									
<7%	N (%)	394 (85.5) *	408 (88.9) *	428 (92.2) *	374 (81.1) *	264 (88)	283 (88)	303 (91)	496 (51)
	OR (95% CI)	1.54 (1.06 to 2.23)	2.14 (1.44 to 3.17)	3.03 (1.97 to 4.66)	REF	4.78 (3.47 to 6.58)	9.23 (6.31 to 13.49)	11.87 (7.88 to 17.89)	REF
	P value	NR	NR	NR	NR	<0.0001	<0.0001	<0.0001	REF
<6.5%	N (%)	341 (74) *	377 (82.1) *	404 (87.1) *	305 (66.2) *	215 (66)	244 (76)	271 (81)	310 (32)
	OR (95% CI)	1.63 (1.2 to 2.21)	2.75 (1.98 to 3.82)	3.95 (2.78 to 5.61)	REF	4.86 (3.66 to 6.45)	8.93 (6.53 to 12.21)	11.84 (8.52 to 16.45)	REF

Study		SURPASS-2				SURPASS-4			
Arm		TZP 5 mg	TZP 10 mg	TZP 15 mg	SEM 1 mg	TZP 5 mg	TZP 10 mg	TZP 15 mg	Insulin Glargine
	P value	NR	NR	NR	NR	<0.0001	<0.0001	<0.0001	REF
<5.7%	N (%)	135 (29.3) *	205 (44.7) *	236 (50.9) *	91 (19.7) *	75 (23)	105 (33)	144 (43)	33 (3)
	OR (95% CI)	1.86 (1.35 to 2.57)	3.94 (2.88 to 5.39)	5.1 (3.73 to 6.97)	REF	9.57 (6.16 to 14.86)	17.11 (11.12 to 26.35)	26.53 (17.35 to 40.56)	REF
	P value	NR	NR	NR	NR	<0.0001	<0.0001	<0.0001	REF
Achieving bodyweight targets									
≥5%	N (%)	316 (68.6)*	378 (82.4) *	400 (86.2) *	270 (58.4) *	205 (63)	249 (78)	285 (85)	78 (8)
	OR (95% CI)	1.58 (1.2 to 2.08)	3.49 (2.57, 4.75)	4.6 (3.32, 6.38)	REF	21.42 (15.53 to 29.89)	46.14 (32.05 to 66.42)	76.93 (51.76 to 114.35)	REF
	P value	NR	NR	NR	NR	<0.0001	<0.0001	<0.0001	REF
≥10 %	N (%)	165 (35.8)*	243 (52.9)*	301 (64.9)*	117 (25.3)*	117 (36)	170 (53)	219 (66)	15 (2)
	OR (95% CI)	1.68 (1.26, 2.25)	3.58 (2.69, 4.77)	5.85 (4.37, 7.82)	REF	20.6 to 61.5)	76.79 (44.2 to 132.7)	127.5 (73.5 to 221.1)	REF
	P value	NR	NR	NR	NR	<0.0001	<0.0001	<0.0001	REF
≥15 %	N (%)	70 (15.2)*	127 (27.7)*	185 (39.9)*	40 (8.7)*	45 (14)	77 (24)	122 (37)	5 (<1)
	OR (95% CI)	1.92 (1.27, 2.90)	4.27 (2.9, 6.29)	7.44 (5.09, 10.87)	NR	28.58 (11.88 to 68.75)	59.14 (25 to 139.86)	105.74 (45.1 to 247.87)	REF
	P value	NR	NR	NR	NR	<0.0001	<0.0001	<0.0001	REF

Frias 2020 has two 15-mg arms following two different dosing schedules. BL: baseline, CI: confidence interval, DBP: diastolic blood pressure, ETD: estimated treatment difference, N: number, NR: not reported, OR: odds ratio, PBO: placebo, REF: reference, SBP: systolic blood pressure, SD: standard deviation , TZP: tirzepatide OR: odds ratio, CI: confidence interval, BL: baseline, DBP: diastolic blood pressure, SBP: systolic blood pressure

*Efficacy estimand

†Digitized and calculated by ICER staff

Table D4.6. Safety – Tirzepatide^{6,7,33,35}

Study	Frias 2018					Frias 2020				SURPASS-2		SURPASS-4			
Arm	PBO	TZP 5 mg	TZP 10 mg	TZP 15 mg	DUA 1.5 mg	PBO	TZP 12 mg	TZP 15 mg-1	TZP 15 mg-2	TZP 15 mg	SEM 1 mg	TZP 5 mg	TZP 10 mg	TZP 15 mg	Insulin Glargine
N	51	55	51	53	54	26	29	28	28	470	469	328	326	337	998
Timepoint	26 wks					12 wks				40 wks		104 wks			
Any AE	NR	NR	NR	NR	NR	NR	NR	NR	NR	324 (68.9)	301 (64.2)	NR	NR	NR	NR
Any TEAE, n (%)	27 (52.9)	40 (72.7)	40 (78.4)	45 (84.9)	40 (74.1)	13 (50)	23 (79.3)	19 (67.9)	24 (85.7)	NR	NR	232 (71)	241 (74)	259 (77)	679 (68)
D/C Due to AE, n (%)	1 (2.0)	0	1 (2.0)	2 (3.8)	2 (3.7)	0	0	0	0	40 (8.5) †**	19 (4.1) †**	37 (11)	28 (9)	36 (11)	54 (5)
D/C from Study Drug	2 (3.9)	5 (9.1)	3 (5.9)	13 (24.5)	6 (11.1)	1 (3.8)	1 (3.4)	1 (3.6)	0	NR	NR	8%	7%	9%	3%
Serious AE	2 (3.9)	1 (1.8)	3 (5.9)	2 (3.8)	3 (5.6)	0	1 (3.4)	0	0	27 (5.7)	13 (2.8)	48 (15)	54 (17)	41 (12)	193 (19)
Death	1 (2.0)*	0	0	0	0	0	0	0	0	4 (0.9)	1 (0.2)	15 (5)	2 (<1)	8 (2)	25 (4)
Diarrhea	2 (3.9)	13 (23.6)	12 (23.5)	17 (32.1)	9 (16.7)	2 (7.7)	9 (31.0)	10 (35.7)	9 (32.1)	65 (13.8)	54 (11.5)	12.2	19.5	20.4	3.2
Nausea	3 (5.9)	11 (20)	11 (21.6)	21 (39.6)	16 (29.6)	2 (7.7)	7 (24.1)	11 (39.3)	10 (35.7)	104 (22.1)	84 (17.9)	11.9	15.9	22.2	1.6
Decreased Appetite	1 (2.0)	11 (20)	13 (25.5)	10 (18.9)	3 (5.6)	0	4 (13.8)	6 (21.4)	8 (28.6)	42 (8.9)	25 (5.3)	29 (9)	36 (11)	35 (10)	5 (<1)
Vomiting	1 (2)	4 (7.3)	8 (15.7)	14 (26.4)	5 (9.3)	1 (3.8)	5 (17.2)	5 (17.9)	5 (17.9)	46 (9.8)	39 (8.3)	4.9	8.2	8.3	1.1
Headache	2 (3.9)	2 (3.6)	1 (2)	5 (9.4)	1 (1.9)	2 (7.7)	2 (6.9)	6 (21.4)	5 (17.9)	NR	NR	NR	NR	NR	NR
Dyspepsia	0	1 (1.8)	6 (11.8)	2 (3.8)	2 (3.7)	0	5 (17.2)	3 (10.7)	3 (10.7)	43 (9.1)	31 (6.6)	18 (6)	27 (8)	26 (8)	13 (1)
Abdominal Pain	1 (2.0)	1 (1.8)	0	3 (5.7)	1 (1.9)	1 (3.8)	1 (3.4)	5 (17.9)	1 (3.6)	24 (5.1)	24 (5.1)	NR	NR	NR	NR
Dizziness	2 (3.9)	2 (3.6)	2 (3.9)	5 (9.4)	1 (1.9)	2 (7.7)	0	1 (3.6)	3 (10.7)	NR	NR	NR	NR	NR	NR
Hypoglycemia (≤70 mg/dL)	2 (3.9)	4 (7.3)	5 (9.8)	4 (7.5)	2 (3.7)	0	2 (6.9)	5 (17.9)	5 (17.9)	NR	NR	NR	NR	NR	NR
Hypoglycemia (≤54 mg/dL)	NR	NR	NR	NR	NR	0	0	0	0*	8 (1.7)	2 (0.4)	6.7	5.5	6.5	15.0
Severe hypoglycemia	0	0	0	0	0	0	0	0	0	1 (0.2)	0	NR	NR	NR	NR

Cholecystitis	0	0	1 (2.0)	0	1 (1.9)	0	0	0	0	4 (0.9)	2 (0.4)	3 (<1)	1 (<1)	1 (<1)	4 (<1)
Acute pancreatitis	0	2 (3.6)	0	0	0	0	0	0	0	2 (0.4)	3 (0.6)	NR	NR	NR	NR
Injection site reaction	2 (3.9)	3 (5.5)	4 (7.8)	1 (1.9)	6 (11.1)	0	2 (6.9)	2 (7.1)	0	21 (4.5)	1 (0.2)	1 (<1)	2 (<1)	1 (<1)	4 (<1)
Hypersensitivity	5 (9.8)	2 (3.6)	2 (3.9)	2 (3.8)	0	1 (3.8)	0	0	0	8 (1.7)	11 (2.3)	NR	NR	NR	NR
Diabetic Retinopathy	0	0	0	0	0	NR	NR	NR	NR	0	0	5 (2) ‡†	5 (2) ‡†	4(1) ‡†	15 (2) ‡†

Frias 2020 has two 15-mg arms following two different dosing schedules. AE: adverse event, BL: baseline, CI: confidence interval, D/C: discontinuation, ETD: estimated treatment difference, N: number, NR: not reported, OR: odds ratio, PBO: placebo, REF: reference, SD: standard deviation, TEAE: treatment emergent adverse event, TZP: tirzepatide

*Fatal TEAE

†** Discontinuation from active treatment due to AE

‡†Diabetic retinopathy complications

Safety Outcomes for Injectable Semaglutide and Empagliflozin

Injectable Semaglutide

In the SUSTAIN-6 CVOT, there was a greater incidence of AEs that led to study discontinuation in the pooled semaglutide versus pooled placebo group (13% vs. 6.7%, respectively; $p=NR$)(Table D4.10). The rates of gastrointestinal AEs (nausea, vomiting, diarrhea) were greater in the semaglutide arm than placebo, in line with expectations of the GLP-1 RA class. Semaglutide carries an FDA warning label for pancreatitis; less than one percent of participants in either pooled arm experienced adjudicated acute pancreatitis ($p=NR$). Semaglutide is contraindicated in patients with medullary thyroid carcinoma; neither treatment arm experienced this adverse event. There was little difference between arms in occurrence of malignant neoplasms.

Empagliflozin

In the EMPA-REG OUTCOME CVOT, there was a significantly lower incidence of AEs that led to study discontinuation in the pooled empagliflozin versus placebo group (17.3% vs. 19.4%, respectively; $p<.01$)(Table D4.10). The most frequent adverse events in the trial were urinary tract infection and genital infection (Table D4.10). Among female study participants, those treated with empagliflozin experienced fewer rates of urinary tract infection than placebo (36.4% vs. 40.6%, respectively; $p<.05$). Empagliflozin significantly increased the rate of genital infection versus placebo (6.4% vs. 1.8%). Within the empagliflozin treatment arm, female participants had twice the rate of genital infection versus male participants (10.0% vs. 5.0%).

Table D4.7. Baseline Characteristics– NMA Linkage Studies^{39,101-103}

Study		SUSTAIN-2		PIONEER 2			PIONEER 3		HARMONY 3	
Arm		SEM 1.0 mg	SITA 100 mg	OSEM 14 mg	EMPA 25 mg	Total	SEM 14 mg	SITA 100 mg	PBO	SITA 100mg
N		409	407	411	410	821	465	467	101	302
Age, y	Mean (SD)	56.0 (9.4)	54.6 (10.4)	57 (10)	58 (10)	58 (10)	57 (10)	58 (10)	56.1 (10.0)	54.3 (9.8)
Sex, n (%)	Men	205 (50)	208 (51)	206 (50.1%)	209 (51.0%)	415 (50.6%)	247 (53.1%)	238 (51.0%)	50 (49.5)	139 (46.0)
	Women	204 (50)	199 (49)	205 (49.9)	201 (49.0)	406 (49.5)	218 (46.9)	229 (49.0)	NR	NR
Diabetes Duration, y	Mean	6.7	6.6	7.2	7.7	7.4	8.7	8.8	6.7	5.8
	SD	5.6	5.1	5.8	6.3	6.1	6.1	6	6.6	4.8
HbA1c, mean (SD)	%	8.0 (0.9)	8.2 (0.9)	8.1 (0.9)	8.1 (0.9)	8.1 (0.9)	8.3 (0.9)	8.3 (0.9)	8.2 (0.9)	8.1 (0.8)
Weight, kg	Mean	89.2	89.3	91.9	91.3	91.6	91.2	90.9	91.6	90.3
	SD	20.7	19.7	20.5	20.1	20.3	21.7	21	19.3	19.1
BMI, kg/ m ²	mean (SD)	32.5 (6.6)	32.5 (5.8)	32.9 (6.3)	32.8 (5.9)	32.8 (6.1)	32.3 (6.3)	32.5 (6.2)	32.8 (5.4)	32.5 (5.4)
Waist Circumference, cm	Mean (SD)	NR	NR	NR	NR	NR	NR	NR	NR	NR
eGFR	Mean (SD)	97 (55-171)	98 (53-194)	96 (15)	95 (15)	95 (15)	95 (16)	96 (15)	NR	NR
Metformin use, N (%)	Yes	407 (100)	405 (100)	411 (100)	410 (100)	821 (100)	465 (100)	467 (100)	NR	NR
Ethnicity, N (%)	Hispanic	67 (16)	73 (18)	91 (22.1)	108 (26.3)	199 (24.2)	75 (16.1)	93 (19.9)	32 (31.7)	111 (36.8)
	Non-Hispanic	342 (84)	334 (82)	320 (77.9)	302 (73.7)	622 (75.8)	390 (83.9)	374 (80.1)	NR	NR
Race, N (%)										
American Indian/Alaska Native		NR	NR	NR	NR	NR	5 (1.1)	6 (1.3)	NR	NR
Black		24 (6)	17 (4)	26 (6.3)	33 (8.0)	59 (7.2)	45 (9.7)	39 (8.4)	23 (22.8)	35 (11.6)

Asian	99 (24)	102 (25)	28 (6.8)	21 (5.1)	49 (6.0)	61 (13.1)	59 (12.6)	5 (5.0)	20 (6.6)
White	279 (68)	281 (69)	355 (86.4)	353 (86.1)	708 (86.2)	317 (68.2)	333 (71.3)	64 (63.4)	225 (74.5)

EMPA: empagliflozin, N: number, NR: not reported, OSEM: oral semaglutide, REF: reference, SD: standard deviation, SEM: semaglutide, y: years

Table D4.8. Outcomes– NMA Linkage Studies^{39,101-103}

Study		SUSTAIN-2			PIONEER-2		PIONEER-3		HARMONY 3	
Arm		SEM 0.5 mg	SEM 1.0 mg	SITA 100 mg	OSEM 14 mg	EMPA 25 mg	OSEM 14 mg	SITA 100 mg	PBO	SITA 100 mg
N		409	409	407	411	410	465	467	101	302
Timepoint		56 wks			52 wks				104 wks	
Glycaemia Endpoints										
HbA1c, %	Mean, Change from BL	-1.3	-1.6*	-0.5*	-1.3	-0.9	-1.2	-0.7	NR	NR
	ETD	-0.77	-1.06	NA	-0.4	REF	-0.5	REF	NR	NR
	95% CI (ETD)	-0.92 to -0.62	-1.21 to -0.91	NA	-0.5 to -0.3	REF	-0.6 to -0.3	REF	NR	NR
	P-Value (ETD)	<0.0001	<0.0001	REF	<0.0001	REF	<0.001	REF	NR	NR
Lipid Levels										
Triglycerides , mg/dL	Change from BL	NR	NR	NR	-16.8	-15.9	-10.9	-1.6	-3.5*	-13.2*
Total Cholesterol, mg/dL	Change from BL	NR	NR	NR	-5.4	3.9	-1.7	1.7	-1.9*	-3.5*
HDL Cholesterol, mg/dL	Change from BL	NR	NR	NR	0.4	2.7	0.4	0	1.2*	1.2*
LDL Cholesterol, mg/dL	Change from BL	NR	NR	NR	-3.9	2.7	0	2.8	-1.2*	-1.9*
Body Weight Endpoints										
Weight, kg	Change from BL	-4.3	-6.1	-1.9	-3.8	-3.6	-3.4	-0.8	-1.0*	-0.86*

	ETD	-2.35	-4.20	NA	-0.2	REF	-2.7	REF	NR	NR
	95% CI	-3.06 to -1.63	-4.91 to -3.49	NA	-0.9 to 0.5	REF	-3.3 to -2.1	REF	NR	NR
	P-Value	<0.0001	<0.0001	REF	0.6231	REF	<0.001	REF	NR	NR
Mean Waist Circumference, cm	Change from BL	-4.3	-5.9	-2.2	-3.5	-2.9	-2.6	-0.4	NR	NR
	ETD	-2.10	-3.67	NA	-0.6	REF	-2.2	REF	NR	NR
	95% CI	-2.91 to -1.29	-4.48 to -2.87	NA	-1.4 to 0.2	REF	-3.0 to -1.4	REF	NR	NR
	P-Value	<0.0001	<0.0001	REF	0.1488	REF	<0.001	REF	NR	NR
Blood Pressure and Pulse Rate										
DBP, mmHg	Change from BL	-2.0	-1.9	-1.1	-2	-2	-2	-1	0*	0.2*
	ETD	-0.90	-0.80	NA	0	REF	-1	REF	NR	NR
	95% CI	-2.10 to 0.30	-2.00 to 0.40	NA	-1 to 2	REF	-2 to 0	REF	NR	NR
	P-Value	0.14	0.19	REF	0.6551	REF	0.28	REF	NR	NR
SBP, mmHg	Change from BL	-5.1	-5.6	-2.3	-4	-4	-3	-1	2.2*	0.2*
	ETD	-2.78	-3.32	NA	0	REF	-2	REF	NR	NR
	95% CI	-4.59 to -0.97	-5.13 to -1.52	NA	-2 to 2	REF	-4 to -1	REF	NR	NR
	P-Value	0.0026	0.0003	REF	0.9371	REF	0.01	REF	NR	NR
Achieving HbA1C Targets										
<7%	N (%)	282 (69) *	321 (78) *	148 (36) *	214 (55.7)	149 (39)	53	31	15.5*	31.6*
	OR (95% CI)	4.16 (3.02 to 5.74)	7.92 (5.59 to 11.22)	REF	2.03 (1.50 to 2.74)	REF	22 (16 to 28)	REF	NR	NR
	P value	<0.0001	<0.0001	REF	<0.0001	REF	<0.001	REF	NR	NR
<6.5%	N (%)	215 (53) *	270 (66) *	83 (20) *	182 (47.4)	83 (21.7)	32	14	7.2*	15.2 *

	OR (95% CI)	4.39 (3.15 to 6.12)	8.99 (6.36 to 12.72)	REF	3.36 (2.43 to 4.66)	REF	18 (13 to 24)	REF	NR	NR
	P value	<0.0001	<0.0001	REF	<0.0001	REF	<0.001	REF	NR	NR
<5.7%	N (%)	NR	NR	NR	NR	NR	NR	NR	NR	NR
	OR (95% CI)	NR	NR	NR	NR	NR	NR	NR	NR	NR
	P value	NR	NR	NR	NR	NR	NR	NR	NR	NR
Achieving bodyweight reduction Targets										
≥5%	N (%)	188 (46)	254 (62)	75 (18)	NR	NR	34	12	NR	NR
	OR (95% CI)	3.76 (2.72 to 5.19)	7.47 (5.38 to 10.37)	REF	NR	NR	22 (16 to 27)	REF	NR	NR
	P value	<0.0001	<0.0001	REF	NR	NR	<0.001	REF	NR	NR
≥10 %	N (%)	52 (13)	97 (24)	14 (3)	58 (15)	30 (7.8)	11	3	NR	NR
	OR (95% CI)	4.09 (2.26 to 7.40)	8.85 (5.01 to 15.61)	REF	2.05 (1.28 to 3.28)	REF	8 (5 to 12)	REF	NR	NR
	P value	<0.0001	<0.0001	REF	0.0028	REF	<0.001	REF	NR	NR
≥15 %	N (%)	NR	NR	NR	NR	NR	NR	NR	NR	NR
	OR (95% CI)	NR	NR	NR	NR	NR	NR	NR	NR	NR
	P value	NR	NR	NR	NR	NR	NR	NR	NR	NR

BL: baseline, CI: confidence interval, DBP: diastolic blood pressure, EMPA: empagliflozin, ETD: estimated treatment difference, N: number, NA: not applicable, NR: not reported, OR: odds ratio, OSEM: oral semaglutide, REF: reference, SBP: systolic blood pressure, SD: standard deviation, SEM: semaglutide, y: years, Lipid levels are NR, HbA1c target of <5.7% is NR, bodyweight target of >15% is NR.

*Efficacy estimand

Table D4.9. Safety– NMA Linkage Studies^{39,101-103}

Study	SUSTAIN-2		PIONEER 2		PIONEER 3		HARMONY 3	
Arm	SEM 1.0 mg	SITA 100 mg	OSEM 14 mg	EMPA 25 mg	OSEM 14 mg	SITA 100 mg	PBO	SITA 100 mg
N	409	407	411	410	465	467	101	302
Timepoint	40 wks		52 wks		56 wks		104 wks	
Any AE	NR	NR	289 (70.5)	283 (69.2)	370 (79.6)	388 (83.3)	79.2	79.1
Any TEAE, n (%)	292 (71)	292 (72)	NR	NR	NR	NR	20.8	17.9
D/C Due to AE, n (%)	39 (10)	12 (3)	44 (10.7)	18 (4.4)	54 (11.6)	24 (52.)	5	3.6
D/C from Study Drug	NR	NR	NR	NR	NR	NR	NR	NR
Serious AE	30 (7)	29 (7)	27 (6.6)	37 (9.0)	44 (9.5)	58 (12.4)	12.9	17.9
Death	1 (1)	3 (1)	0	1 (0.2)	1 (0.2)	3 (0.6)	NR	NR
Diarrhea	53 (13)	29 (7)	38 (9.3)	13 (3.2)	57 (12.3)	37 (7.9)	10.9	8.6
Nausea	72 (18)	30 (7)	81 (19.8)	10 (2.4)	70 (15.1)	32 (6.9)	10.9	6.6
Decreased Appetite	27 (7)	11 (3)	21 (5.1)	2 (0.5)	32 (6.9)	14 (3.0)	NR	NR
Vomiting	41 (10)	11 (3)	30 (7.3)	7 (1.7)	42 (9.0)	19 (4.1)	NR	NR
Headache	29 (7)	17 (4)	NR	NR	37 (8.0)	36 (7.7)	NR	NR
Dyspepsia	20 (5)	9 (2)	NR	NR	0	0	NR	NR
Abdominal Pain	NR	NR	5 (1.2)	0	2 (0.4)	2 (0.4)	NR	NR
Dizziness	NR	NR	NR	NR	1 (0.2)	0	NR	NR
Hypoglycemia (plasma glucose ≤70 mg/dL)	NR	NR	45	39	131 (28.2)	112 (24.0)	NR	NR
Hypoglycemia (plasma glucose ≤54 mg/dL)	NR	NR	7 (1.7)	8 (2.0)	36 (7.7)	39 (8.4)	4 (4.0)	5 (1.7)
Severe hypoglycemia	NR	NR	1 (0.2)	1 (0.2)	1 (0.2)	4 (0.9)	0	0
Cholecystitis	1 (1)	0 (0)	NR	NR	NR	NR	NR	NR
Acute pancreatitis	--	--	1 (0.2)	1 (0.2)	3 (0.6)	1 (0.2)	NR	NR
Injection site reaction	NR	NR	NA	NA	NA	NA	2	1.7
Hypersensitivity	NR	NR	NR	NR	NR	NR	NR	NR
Diabetic Retinopathy	0	3 (1)	13 (3.2)	4 (1.0)	17 (3.7)	29 (6.2)	NR	NR

AE: adverse event, CI: confidence interval, D/C: discontinuation, EMPA: empagliflozin, N: number, NA: not applicable, NR: not reported, OR: odds ratio, OSEM: oral semaglutide, REF: reference, SD: standard deviation, SEM: semaglutide, TEAE: treatment emergent adverse event, y: years

Table D4.10. CVOT Studies^{8,9}

Study		SUSTAIN-6		EMPA-REG -OUTCOME	
	Arms	SEM 1mg	PBO 1mg	PBO	EMPA (10/25mg)
N		822	825	2333	4687
	Timepoint	109 wks		156 wks	
Baseline Characteristics					
Age, yr		64.7 (7.1)	64.4 (7.5)	63.2 (8.8)	63.1 (8.6)
Male sex, N (%)		518 (63)	507 (61.5)	1680 (72.0)	3336 (71.2)
Weight, kg (SD)		92.9 (21.1)	91.9 (20.8)	86.6 (19.1)	86.2 (18.9)
BMI, kg/m ² (SD)		32.9 (6.2)	32.7 (6)	30.7 (5.2)	30.6 (5.3)
Type 2 Diabetes					
Diabetes Duration, yr (SD)		14.1 (8.2)	13.2 (7.4)	NR	NR
Hemoglobin, % (SD)		8.7 (1.5)	8.7 (1.5)	8.08 (0.84)	8.07 (0.85)
Race					
White, n (%)		691 (84.1)	676 (81.9)	1678 (71.9)	3403 (72.6)
Asian, n (%)		58 (7.1)	72 (8.7)	511 (21.9)	1006 (21.5)
Black, n (%)		54 (6.6)	59 (7.2)	120 (5.1)	237 (5.1)
Ethnicity					
Hispanic or Latino, n (%)		124 (15.1)	137 (16.6)	418 (17.9)	847 (18.1)
Antihyperglycemic Medication at Baseline					
Metformin, n (%)		594 (72.3)	617 (74.8)	1734 (74.3)	3459 (73.8)
Insulin, n (%)		477 (58.0)	479 (58.1)	1135 (48.6)	2252 (48.0)
Sulfonylurea, n (%)		349 (42.5)	349 (42.3)	992 (42.5)	2014 (43.0)
Cardiovascular Risk Factors					
Systolic Blood Pressure (mmHg)		135.8 (17)	134.8 (17.5)	135.8 (17.2)	135.3 (16.9)
Diastolic Blood Pressure (mmHg)		76.9 (10.2)	76.7 (10.2)	76.8 (10.1)	76.6 (9.7)
LDL Cholesterol		83.3 (41.2)	83.6 (45.9)	84.9 (35.3)	85.9 (36.0)
Never Smoked, n (%)		364 (44.3)	348 (42.2)	957 (41.0)	1925 (41.1)
History of cardiovascular disease, N (%)					

Ischemic Heart Disease		495 (60.2)	496 (60.1)	1763 (75.6)	3545 (75.6)
Myocardial Infarction		264 (32.1)	275 (33.3)	1083 (46.4)	2190 (46.7)
Heart Failure		180 (21.9)	206 (25)	244 (10.5)	462 (9.9)
Ischemic Stroke		89 (10.8)	109 (13.2)	553 (23.7)	1084 (23.1)
Hemorrhagic Stroke		24 (2.9)	29 (3.5)		
Hypertension		771 (93.8)	760 (92.1)	NR	NR
Renal Function (Estimated glomerular filtration rate)					
≥90 mL/min/1.73m ² (normal)		246 (29.9)	252 (30.5)	488 (20.9)	1050 (22.4)
60 to <90 mL/min/1.73m ² (mild)		357 (43.4)	346 (41.9)	1238 (53.1)	2423 (51.7)
<60 mL/min/1.73m ² (moderate to end stage)		219 (26.6)	227 (27.5)	607 (26.0)	1212 (25.9)
Outcomes					
Composite Outcome (CV Causes, Non-fatal MI, or Non-fatal stroke)	n (%)	108 (6.6)	146 (8.9)	282 (12.1)	490 (10.5)
	HR (95%CI)	0.74 (0.58 to 0.95)		0.86 (0.74 to 0.99)	
	p-value, NON	<0.001	REF	<.001	REF
	p-value, SUP	0.02	REF	0.04	REF
Expanded Composite Outcome (Death from CV Causes, Non-fatal MI, Non-fatal Stroke, Revascularization, Hospitalization, HF)	n (%)	199 (12.1)	264 (16.0)	333 (14.3)	599 (12.8)
	HR (95%CI)	0.74 (0.62 to 0.89)		0.89 (0.78 to 1.01)	
	p-value, NON	0.002	REF	<.001	REF
	p-value, SUP	NA	NA	0.08	REF
All Cause Death, Non-fatal MI, or Non-fatal Stroke	n (%)	122 (7.4)	158 (9.6)	NR	NR
	HR (95%CI)	0.77 (0.61 to 0.97)		NR	NR
	p-value	0.03	REF	NR	NR
From Any Cause	n (%)	62 (3.8)	60 (3.6)	194 (8.3)	269 (5.7)
	HR (95%CI)	1.05 (0.74 to 1.50)		0.68 (0.57 to 0.82)	
	p-value	0.79	REF	<.001	REF
From Cardiovascular Cause	n (%)	44 (2.7)	46 (2.8)	137 (5.9)	172 (3.7)
	HR (95%CI)	0.98 (0.65 to 1.48)		0.62 (0.49 to 0.77)	
	p-value	0.92	REF	<.001	REF
Non-fatal MI		n (%)	47 (2.9)	64 (3.9)	121 (5.2)* 213 (4.5)

	HR (95%CI)	0.74 (0.51 to 1.08)		0.87 (0.70-1.09)	
	p-value	0.12	REF	0.22	REF
Non-fatal Stroke	n (%)	27 (1.6)	44 (2.7)	60 (2.6)	150 (3.2)
	HR (95%CI)	0.61 (0.38 to 0.99)		1.24 (0.92 to 1.67)	
	p-value	0.04	REF	0.16	REF
Hospitalization for Unstable Angina	n (%)	22 (1.3)	27 (1.6)	66 (2.8)	133 (2.8)
	HR (95%CI)	0.82 (0.47 to 1.44)		0.99 (0.74 to 1.34)	
	p-value	0.49	REF	0.97	REF
Revascularization	n (%)	83 (5.0)	126 (7.6)	186 (8.0)	329 (7.0)
	HR (95%CI)	0.65 (0.50 to 0.86)		0.86 (0.72 to 1.04)	
	p-value	0.003	REF	0.11	REF
Hospitalization for Heart Failure	n (%)	59 (3.6)	54 (3.3)	95 (4.1)	126 (2.7)
	HR (95%CI)	1.11 (0.77 to 1.61)		0.65 (0.50 to 0.85)	
	p-value	0.57	REF	0.002	REF
Retinopathy Complications	n (%)	50 (3.0)	29 (1.8)	29 (1.2)	41 (0.9)
	HR (95%CI)	1.76 (1.11 to 2.78)		0.69 (0.43 to 1.12)	
	p-value	0.02	REF	0.134	REF
New or worsening nephropathy	n (%)	62 (3.8)	100 (6.1)	388 (18.8)	525 (12.7)
	HR (95%CI)	0.64 (0.46 to 0.88)		0.61 (0.53 to 0.70)	
	p-value	0.005	REF	<0.001	REF
Adverse Events, n (%)					
Adverse Events		732 (89.1)	736 (89.2)	2139 (91.7)	4230 (90.2)
Serious Adverse Event		276 (33.6)	298 (36.1)	592 (25.4)	1100 (23.5)
Severe Adverse Events		207 (25.2)	194 (23.5)	988 (42.3)	1789 (38.2)
Leading to D/C		119 (14.5)	63 (7.6)	453 (19.4)	813 (17.3)
Gastrointestinal Disorder		430 (52.3)	290 (35.2)	NR	NR
Nausea		151 (18.4)	87 (10.5)	NR	NR
Vomiting		122 (14.8)	34 (4.1)	NR	NR
Diarrhea		151 (18.4)	87 (10.5)	NR	NR

Gallbladder Disorder		26 (3.2)	23 (2.8)	NR	NR
Cholelithiasis		17 (2.1)	12 (1.5)	NR	NR
Acute Cholecystitis		0	2 (0.2)	NR	NR
Neoplasm		89 (10.8)	69 (8.4)	NR	NR
Benign		54 (6.6)	34 (4.1)	NR	NR
Malignancy, any		40 (4.9)	35 (4.2)	NR	NR
Malignancy, pancreatic		1 (0.1)	2 (0.2)	NR	NR
Other					
Severe or symptomatic hypoglycemic event		178 (21.7)	173 (21.0)	NR	NR
Acute renal failure		23 (2.8)	35 (4.2)	155 (6.6)	256 (5.2)
Allergic reaction		49 (6.0)	57 (6.9)	NR	NR
Injection Site Reaction		9 (1.1)	12 (1.5)	NR	NR
Cardiac disorder		150 (18.2)	173 (21.0)	NR	NR
Atrial Fibrillation		23 (2.8)	26 (3.2)	NR	NR
Acute Pancreatitis		3 (0.4)	9 (1.1)	NR	NR
Event Consistent with Urinary Tract Infection	Total	NR	NR	423 (18.1)	842 (18)
	Male Patients	NR	NR	158 (9.4)	350 (10.5)
	Female Patients	NR	NR	265 (40.6)	492 (36.4)
Complicated Urinary Tract Infection		NR	NR	41 (1.8)	82 (1.7)
Event Consistent with Genital Infection	Total	NR	NR	42 (1.8)	301 (6.4)
	Male Patients	NR	NR	25 (1.5)	166 (5.0)
	Female Patients	NR	NR	17 (2.6)	135 (10.0)
Acute Kidney Failure		23 (2.8)	35 (4.2)	155 (6.6)	246 (5.2)
Acute Kidney Injury		NR	NR	37 (1.6)	45 (1.0)
Diabetic Ketoacidosis		NR	NR	1 (<0.1)	4 (0.1)
Intermediate Outcomes at Week 104					
HbA1c Change from baseline, %		-1.4	-0.4	-0.1	-0.6†
Body Weight CFB, kg		-4.9	-0.5	-0.8	-3†
SBP CFB, mmHg		-5.37	-2.78	-0.8	-4.1

DBP CFB, mmHg	-1.57	-1.71	-1.5‡	-2.2‡
Total Cholesterol, Ratio to Baseline	0.97	0.99	NR	NR
LDL Cholesterol, Ratio to Baseline	0.97	0.99	1.03	1.05
HDL Cholesterol, Ratio to Baseline	1.01	0.97	1.01	1.05
Triglycerides, Ratio to Baseline	0.92	0.98	NR	NR

CI: confidence interval, CFB: change from baseline, CV: cardiovascular, DBP: diastolic blood pressure, HF: heart failure, HR: hazard ration, N: number, NR: not reported, REF: reference, SBP: systolic blood pressure, SD: standard deviation, y: years

*Nonfatal myocardial infarction excluding silent myocardial infarction

†Outcomes for EMPA 25 mg

‡Digitized and calculated by ICER staff

Table D4.11. CVOT Subgroups^{8,9}

CVOT Subgroups					
Study		SUSTAIN-6		EMPA-REG -OUTCOME	
Arm		SEM 1mg	PBO 1mg	PBO	EMPA
N		822	825	2333	4687
Timepoint		109 wks		156 wks	
Composite Outcome					
BMI (kg/m²)	<30	6.2	8.3	7.57	10.13
	HR: 95% CI; p-value	0.58 (0.39 to 0.87); 0.16		0.74 (0.60 to 0.90)); 0.06	
	>30	6.6	7.7	9.05	9.08
	HR: 95% CI; p-value	0.84 (0.61 to 1.16); 0.16		0.98 (0.80 to 1.21); 0.06	
Race	Asian	6.6	11.2	0.11	0.07
	HR: 95% CI; p-value	0.58 (0.25 to 1.34); 0.88		0.66 (0.48 to 0.95); 0.09	
	Black/AA	4.6	6.2	0.12	0.16
	HR: 95% CI; p-value	0.72 (0.23 to 2.28); 0.88		1.48 (0.80 to 2.72)); 0.09	
	White	6.7	8.7	0.12	0.10
	HR: 95% CI; p-value	0.76 (0.58 to 1.00); 0.88		0.88 (0.74 to 1.04); 0.09	
Ethnicity	Hispanic	5.1	7.5	8.03	12.1
	HR: 95% CI; p-value	0.67 (0.33 to 1.36); 0.80		0.63 (0.44 to 0.90); 0.07	
	Non	6.8	8.7	8.31	9.13
	HR: 95% CI; p-value	0.74 (0.57 to 0.96); 0.80		0.91 (0.77 to 1.07); 0.07	
Chronic HF	No	5.4	8.2	NR	NR
	HR: 95% CI; p-value	0.64 (0.48 to 0.86); 0.09		NR	NR
	Yes	12.3	11.8	NR	NR
	HR: 95% CI; p-value	1.03 (0.64 to 1.66); 0.09		NR	NR
CVD Status	Established	5.4	9.9	NR	NR
	HR: 95% CI; p-value	0.72 (0.55 to 0.93); 0.49		NR	NR
	Risk	3.4	3.4	NR	NR
	HR: 95% CI; p-value	1.00 (0.41 to 2.46); 0.49		NR	NR

eGFR <60 mL/min/1.73m ²	No	5.3	7.9	NR	NR
	HR: 95% CI; p-value	0.67 (0.48 to 0.92)); 0.98		NR	NR
	Yes	9.6	11.3	6.13	6.89
	HR: 95% CI; p-value	0.84 (0.57 to 1.25)); 0.37		0.88 (0.69 to 1.13); 0.20	
CVD Death					
BMI (kg/m ²)	<30	NR	NR	14.36	28.49
	HR: 95% CI; p-value	NR	NR	0.50 (0.37 to 0.68)*; 0.05	
	>30	NR	NR	20.6	26.17
	HR: 95% CI; p-value	NR	NR	0.78 (0.56 to 1.08)*; 0.05	
Ethnicity	Hispanic	NR	NR	14.93	22.32
	HR: 95% CI; p-value	NR	NR	0.53 (0.32 to 0.88)); 0.49	
	Non	NR	NR	17.54	27.2
	HR: 95% CI; p-value	NR	NR	0.64 (0.50 to 0.83); 0.49	
eGFR <60 mL/min/1.73m ²	No	NR	NR	NR	NR
	HR: 95% CI; p-value	NR	NR	NR	NR
	Yes	NR	NR	12.64	16.16
	HR: 95% CI; p-value	NR	NR	0.78 (0.54 to 1.12)2); 0.15	

AA: African American, BL: baseline, CI: confidence interval, DBP: diastolic blood pressure, EMPA: empagliflozin, ETD: estimated treatment difference, N: number, NR: not reported, OR: odds ratio, PBO: placebo, REF: reference, SBP: systolic blood pressure, SD: standard deviation, SEM: semaglutide

*p=0.05 for interaction. All other tests for interaction were non-significant.

†Efficacy estimand

Table D4.12. SURPASS-4 CV and Death Outcomes³⁵

Study	SURPASS-4 CV Outcomes				
Arms	TZP 5 mg	TZP 10 mg	TZP 15 mg	All TZP	Insulin Glargine
N	329	328	338	995	1000
Timepoint	104 weeks				
MACE-4	19 (6)	17 (5)	11 (3)	47 (5)	62 (6)
HR; 95% CI	--	--	--	0.74 (0.51 to 1.08)*	
CV Death	10 (3)	1 (<1)	5 (2)	16 (2)	21 (2)
MI	7 (2)	9 (3)	3 (<1)	19 (2)	21 (2)
Hospitalization for unstable angina	0	2 (<1)	2 (<1)	4 (<1)	8 (<1)
Stroke	5 (2)	5 (2)	1 (<1)	11 (<1)	13 (<1)
Other MACE					
Coronary Interventions	10 (3)	11 (3)	8 (2)	29 (3)	37 (4)
Transient Ischemic Attack	0	2 (<1)	1 (<1)	3 (<1)	0
Hospitalization for HF	1 (<1)	1 (<1)	2 (<1)	4 (<1)	0
HR; 95% CI	--	--	--	--	--
Death	15 (5)	2 (<1)	8 (2)	25 (3)	35 (4)
HR; 95% CI	--	--	--	0.70 (0.42 to 1.17)*	
CV Death	4 (1)	0	2 (<1)	6 (<1)	9 (<1)
Undetermined	6 (2)	1 (<1)	3 (<1)	10 (1)	12 (1)
Non-CV	5 (2)	1 (<1)	3 (<1)	9 (<1)	14 (1)

CI: confidence interval, CV: cardiovascular, HF: heart failure, HR: hazard ration, MI: myocardial infarction, TZP: tirzepatide

*Point estimate and 95%CI of hazard ratio comparing pooled TZP versus insulin

D5. Ongoing Studies

Figure D5.1. Ongoing Studies

Title / Trial Sponsor	Study Design	Comparators	Patient Population	Primary Outcomes	Estimated Completion Dates
A Study of Tirzepatide (LY3298176) Compared with Dulaglutide on Major Cardiovascular Events in Participants with Type 2 Diabetes (SURPASS-CVOT)	Phase III, double-blind, randomized N= 12,500	Arm 1: tirzepatide (SC) Arm 2: Dulaglutide (SC)	<u>Inclusion:</u> Have a diagnosis of T2DM Have confirmed atherosclerotic cardiovascular disease HbA1c $\geq 7.0\%$ to $\leq 10.5\%$ BMI ≥ 25 kg/m ² <u>Exclusion:</u> Type 1 diabetes MACE in the last 60 days History of severe hypoglycemia History of pancreatitis	Time to first occurrence of death from CV causes, MI, or stroke (MACE-3)	October 2024
A Study of Tirzepatide (LY3298176) in Participants with Type 2 Diabetes on Metformin with or Without Sulfonylurea (SURPASS-AP-Combo)	Phase III, open-label N= 917	Arm 1: Tirzepatide 5 mg SC Arm 2: Tirzepatide 10 mg SC Arm 3: Tirzepatide 15 mg SC Arm 4: Insulin glargine (SC)	<u>Inclusion:</u> Have a diagnosis of T2DM Type 2 diabetes mellitus Stable metformin with or without a sulfonylurea for at least 2 months Are insulin-naïve (except for the use of insulin for treatment of gestational diabetes or short-term use [≤ 14 consecutive days] for acute conditions)	Mean change from baseline in HbA1c (10 mg and 15 mg)	November 2021

Title / Trial Sponsor	Study Design	Comparators	Patient Population	Primary Outcomes	Estimated Completion Dates
			<p>HbA1c $\geq 7.5\%$ to $\leq 11.0\%$ at screening</p> <p>Stable weight ($\pm 5\%$) ≥ 3 months</p> <p>BMI ≥ 23 kg/m²</p> <p><u>Exclusion:</u></p> <p>Type 1 diabetes mellitus</p> <p>Have history of chronic or acute pancreatitis</p> <p>Have history of proliferative diabetic retinopathy; or diabetic maculopathy; or non-proliferative diabetic retinopathy that requires acute treatment</p> <p>Have a history of severe hypoglycemia and/or hypoglycemia unawareness within the three months</p>		
A Study of Tirzepatide (LY3298176) Versus Insulin Lispro (U100) in Participants with Type 2 Diabetes Inadequately Controlled on Insulin Glargine (U100) With or Without Metformin (SURPASS-6)	<p>Phase III, randomized, open-label</p> <p>N= 1182</p>	<p>Arm 1: Tirzepatide 5 mg SC + insulin (U100)</p> <p>Arm 2: Tirzepatide 10 mg SC + insulin (U100)</p> <p>Arm 3: Tirzepatide 15 mg SC + insulin (U100)</p> <p>Arm 4: Insulin glargine (SC) + insulin (U100)</p>	<p><u>Inclusion:</u></p> <p>Have been diagnosed with T2DM</p> <p>Have HbA1c between $\geq 7.5\%$ and $\leq 11\%$</p> <p>Have been treated for at least 90 days prior to day of screening with once or twice daily basal insulin with or without stable dose of metformin ≥ 1500 mg/day and up to maximum approved dose per country specific approved label, sulfonylureas or dipeptidyl peptidase 4 inhibitors</p>	Change from baseline in HBA1C (pooled doses) at week 52	November 2022

Title / Trial Sponsor	Study Design	Comparators	Patient Population	Primary Outcomes	Estimated Completion Dates
			<p>Be of stable weight ($\pm 5\%$) for at least 90 days</p> <p>BMI ≥ 23 (kg/m²) and ≤ 45 kg/m² at screening</p> <p><u>Exclusion:</u> Type 1 diabetes mellitus</p> <p>Chronic or acute pancreatitis any time prior to study entry</p> <p>Proliferative diabetic retinopathy or diabetic macular edema or non-proliferative diabetic retinopathy requiring immediate or urgent treatment</p> <p>Disorders associated with slowed emptying of the stomach, have had any stomach surgeries for the purpose of weight loss, or are chronically taking drugs that directly affect gastrointestinal motility</p> <p>Heart attack, stroke, or hospitalization for congestive heart failure in the past two months</p>		
A Study of Tirzepatide (LY3298176) in Participants with Type 2 Diabetes Who Have Obesity or Are Overweight (SURMOUNT-2)	<p>Phase III randomized, double-blind, placebo controlled</p> <p>N= 900</p>	<p>Arm 1: Tirzepatide 10 mg SC + insulin (Uu100)</p> <p>Arm 2: Tirzepatide 15 mg SC</p> <p>Arm 3: placebo</p>	<p><u>Inclusion</u> Have Type 2 Diabetes (T2DM) with HbA1c $\geq 7\%$ to $\leq 10\%$ at screening, on stable therapy for the last 3 months prior to screening</p> <p>Have a BMI of ≥ 27 kg/m²</p>	Percent change from randomization in body weight and percentage of patients who achieve $>5\%$	June 19, 2023

Title / Trial Sponsor	Study Design	Comparators	Patient Population	Primary Outcomes	Estimated Completion Dates
			<p>Are overweight or have obesity</p> <p>Have a history of at least one self-reported unsuccessful dietary effort to lose body weight</p> <p>Are at least 18 years of age and age of majority per local laws and regulations</p> <p><u>Exclusion</u></p> <p>Have Type 1 diabetes mellitus, history of ketoacidosis or hyperosmolar state/coma or any other types of diabetes except T2DM</p> <p>Have at least 2 confirmed fasting self-monitoring blood glucose (SMBG) values >270 mg/dL (on two nonconsecutive days) prior to Visit 3</p> <p>Have proliferative diabetic retinopathy OR diabetic macular edema OR non-proliferative diabetic retinopathy that requires acute treatment</p> <p>Have self-reported change in body weight >5 kg within 3 months prior to screening</p> <p>Have had a history of chronic or acute pancreatitis</p>	body weight reduction from randomization.	

BMI: body mass index, CV: cardiovascular, HbA1c: hemoglobin A1c, kg: kilogram, MACE-3: 3-point MACE, mg: milligram, MI: myocardial infarction, N: number, SC: subcutaneous, T2DM: type 2 diabetes mellitus

Source: www.ClinicalTrials.gov (NOTE: studies listed on site include both clinical trials and observational studies)

D6. Previous Systematic Reviews and Technology Assessments

We identified one ongoing health technology assessments conducted by the National Institute for Health and Care Excellence (NICE) on tirzepatide summarized below. We also identified two meta-analyses that include empagliflozin and subcutaneous injectable semaglutide in its analyses. All reports are summarized below.

NICE Technology Assessments

[Tirzepatide for Treating Type 2 Diabetes](#)

NICE is currently conducting an appraisal of the clinical and cost- effectiveness of tirzepatide for treating type 2 diabetes. The expected publication date is to be confirmed.

Salsali, A. (2016). "Cardiovascular safety of empagliflozin in patients with type 2 diabetes: a meta-analysis of data from randomized placebo-controlled trials"¹⁰⁴

This meta-analysis assessed the effect on empagliflozin on cardiovascular risk in patients with type 2 diabetes across eight placebo-controlled trials. Using data from all available empagliflozin (EMPA) trials greater than 12 weeks duration, the primary endpoint of this analysis was a composite of cardiovascular (CV) death, non-fatal myocardial infarction (MI), non-fatal stroke, and hospitalization for unstable angina (MACE-4). The secondary endpoint was a composite of CV death, non-fatal MI and non-fatal stroke (MACE-3). Across all trials, 3835 patients were assigned to placebo, 3629 to EMPA 10 mg and 3828 to EMPA 25 mg with an average age of 61 and 65% of patients identifying as male. Baseline characteristics were generally similar across all arms. For pooled EMPA, patients had a reduced risk in MACE-4 versus placebo (HR 0.86; CI: 0.76 to 0.98) and MACE-3 (HR: 0.84; CI: 0.73 to 0.96). Similar results were seen for EMPA 10mg alone and 25mg alone.

Shi, F. (2018) Efficacy and Safety of Once-Weekly Semaglutide for the Treatment of Type 2 Diabetes: A Systematic Review and Meta-Analysis of Randomized Controlled Trials¹⁰⁵

A systematic review and meta-analysis were conducted to assess the efficacy and safety of once weekly injectable semaglutide in adult patients with T2DM. Of the 457 initial studies, nine studies met the inclusion criteria, evaluating a total of 9,773 patients. The outcomes of interest in this evaluation included glycemic control, weight control, blood pressure, pulse rate, and safety. Patients assigned to semaglutide had a significant decrease in HbA1c levels compared to other therapies (weight mean difference [WMD]: -0.93; CI: -1.24 to -0.62; $p < 0.001$) although there was significant heterogeneity (I^2 : 92.6%). Semaglutide also significantly reduced body weight (WMD: -3.47; CI: -3.96 to -2.98; $p < 0.001$; I^2 : 17.6) and systolic blood pressure (WMD: -0.29 mmHg; CI: -0.65 to 0.07; $p = 0.0113$). Semaglutide significantly increased pulse rate compared to other therapies (WMD: 2.21 bpm; CI: 1.54 to 2.88; $p < 0.0001$; I^2 : 67.6). The safety profile was generally well

tolerated with no increased safety risk for semaglutide for adverse events (RR:1.04), serious AEs (RR: 0.93), or fatal AEs (RR: 0.90). There was a significant increase in premature treatment discontinuation associated with semaglutide (RR: 2.07; CI: 1.58 to 2.73; $p<0.001$). The most common AEs were gastrointestinal events (42.9%) followed by nausea (18.1%) and diarrhea (14.0%).

E. Long-Term Cost-Effectiveness: Supplemental Information

E1. Detailed Methods

Table E.1. Impact Inventory

Sector	Type of Impact (Add additional domains, as relevant)	Included in This Analysis from [...] Perspective?		Notes on Sources (if quantified), Likely Magnitude & Impact (if not)
		Health Care Sector	Societal	
Formal Health Care Sector				
Health Outcomes	Longevity effects	X	X	
	Health-related quality of life effects	X	X	
	Adverse events	X	X	
Medical Costs	Paid by third-party payers	X	X	
	Paid by patients out-of-pocket	<input type="checkbox"/>	<input type="checkbox"/>	
	Future related medical costs	<input type="checkbox"/>	<input type="checkbox"/>	
	Future unrelated medical costs	<input type="checkbox"/>	<input type="checkbox"/>	
Informal Health Care Sector				
Health- Related Costs	Patient time costs	NA	<input type="checkbox"/>	
	Unpaid caregiver-time costs	NA	<input type="checkbox"/>	
	Transportation costs	NA	<input type="checkbox"/>	
Non-Health Care Sector				
Productivity	Labor market earnings lost	NA	X	
	Cost of unpaid lost productivity due to illness	NA	X	
	Cost of uncompensated household production	NA	<input type="checkbox"/>	
Consumption	Future consumption unrelated to health	NA	<input type="checkbox"/>	
Social services	Cost of social services as part of intervention	NA	<input type="checkbox"/>	
Legal/Criminal Justice	Number of crimes related to intervention	NA	<input type="checkbox"/>	
	Cost of crimes related to intervention	NA	<input type="checkbox"/>	
Education	Impact of intervention on educational achievement of population	NA	<input type="checkbox"/>	
Housing	Cost of home improvements, remediation	NA	<input type="checkbox"/>	
Environment	Production of toxic waste pollution by intervention	NA	<input type="checkbox"/>	
Other	Other impacts (if relevant)	NA	<input type="checkbox"/>	

NA: not applicable

Adapted from Sanders et al.¹⁰⁶

Target Population

The target population of the model was patients with diabetes that were not controlled with first-line anti-hypoglycemic medication who were eligible for an add-on therapy such as tirzepatide, injectable semaglutide, or empagliflozin. Description of the included patient population, derived from the NHANES is included in the main report. We assume our patient population is eligible for add-on therapy to help control their diabetes. The three add-on (to background therapy) therapies we consider are: tirzepatide, 15 milligrams once weekly via subcutaneous injection; semaglutide (Ozempic®), one milligram once weekly via subcutaneous injection, and empagliflozin (Jardiance®), 25 milligrams daily, orally. We additionally consider background therapy alone, consisting of metformin (with or without sulfonylureas). Greater emphasis is placed on the common comparator of background therapy alone in scenario analyses, as some of these scenarios focus on comparisons across modeled results and trial outcomes versus background therapy alone.

E2. Model Inputs and Assumptions

Key model inputs and assumptions are listed in the main text in [section 4.2](#).

Model Inputs

Clinical Inputs

Baseline clinical inputs came from the individual patients in the target population, derived from NHANES dataset.

We used the point estimates and 95% credible ranges from the random effects model output of the network meta-analysis described above as the treatment effects in the model for the following outcomes: decrease in HbA1c level, decrease in weight, decrease in systolic blood pressure (SBP), and decrease in low-density lipids (LDL). These values can be found in the section reporting NMA output.

Response to Treatment

Clinical inputs regarding the efficacy of tirzepatide, injectable semaglutide, and empagliflozin as compared to placebo on intermediate outcomes such as changes in HbA1c, lipid levels, blood pressure, and body weight were derived from the NMA described above. The treatment effect was assumed to take effect after the first model cycle only, at which point it would be maintained while the patient was on treatment. Ongoing changes in weight and HbA1c were modeled using published equations for those measures in treated diabetic patients.⁶¹

We applied hazard ratios to estimated outcomes, including major adverse cardiac events, congestive heart failure, and nephropathy (Table E.2). Hazard ratios were derived from clinical outcome trials SUSTAIN-6 for injectable semaglutide and from EMPA-REG -OUTCOME for empagliflozin.^{8,36} They were applied by multiplying the UKPDS predicted probability of the event occurring by the HR in each model cycle. No adjustment was made to tirzepatide in the base-case for congestive heart failure and nephropathy, reflected by setting the hazard ratios equal to one. Tirzepatide's hazard ratio for major adverse cardiac events was set equal to the risk reduction observed in SURPASS-4, which is also an equivalent point estimate to semaglutide's SUSTAIN-6. The decision to only modify one group of the risk equation estimates for tirzepatide in the base-case was based primarily on two factors: a lack of available cardiovascular and renal outcome data to reference and the lack of reliable proxy data due to tirzepatide's novel dual mechanism of action. We also tested a scenario where all hazard ratios for tirzepatide were neutralized (set equal to 1).

Table E.2. Hazard Ratios Applied in Base-Case Analysis

Regimen	Hazard Ratio	Source
Composite MACE HR		
Tirzepatide HR vs. Background Tx	0.74	SURPASS-4 ³⁵
Semaglutide HR vs. Background Tx	0.74	SUSTAIN-6. ⁸
Empagliflozin HR vs. Background Tx	0.86	EMPA-REG -OUTCOME ³⁶
Congestive Heart Failure HR		
Tirzepatide HR vs. Background Tx	1.00	Assumption
Semaglutide HR vs. Background Tx	1.11	SUSTAIN-6. ⁸
Empagliflozin HR vs. Background Tx	0.65	EMPA-REG -OUTCOME ³⁶
Nephropathy HR		
Tirzepatide HR vs. Background Tx	1.00	Assumption
Semaglutide HR vs. Background Tx	0.64	SUSTAIN-6. ⁸
Empagliflozin HR vs. Background Tx	0.61	EMPA-REG -OUTCOME ³⁶

HR: hazard ratio, MACE: major adverse cardiovascular event, Tx: treatment

Treatment Discontinuation

The only therapy with extension trial data beyond initial clinical trials was empagliflozin (EMPA-REG Extend), with discontinuation data for patients for 52 weeks assuming they continued the therapy for at least the initial 24 weeks. We derived the probability of discontinuation during the first year after the initial trial period (9.1% per year) from patients who discontinued for any reason; the rationale for choosing discontinuation for any reason was to accurately reflect the number of patients no longer benefiting from the drug regardless of reason. This rate was applied to all included active therapies pending long term trial data availability. Discontinuation of background therapy was derived from the placebo arms of trials for the three active comparators: SURPASS 1, EMPA-REG Extend, and SUSTAIN 2.

Addition of Insulin Therapy

Patients whose HbA1c levels reach 8.5% or above (with a range from 8.0 to 9.0% used in scenario analyses) were assumed to add insulin therapy while continuing on active therapy. Patients were assumed to continue both insulin and their active treatment for the remainder of the model time horizon. Patients being modeled on background therapy alone were assumed to discontinue their background therapy when insulin therapy started. The choice for insulin to be added on top of active therapy was made for several reasons: it reflects the same treatments for GLP-1 and SGLT-2 therapies in the 2019 ICER report; it reflects what we believe would occur in clinical practice; and it allows us to be able to evaluate all comparators head-to-head as opposed to evaluating differences in different medication treatment pathways. Insulin treatment costs were modeled using mean doses from a literature review, applied to unit costs similar to the model comparators.⁶¹ Clinical characteristics for patients on insulin were modeled using the UKPDS-OM2 and Willis equations.^{11,61}

Diabetes-related Complications

We modeled diabetes-related complications and mortality based on risk equations from the UKPDS-OM2 risk engine.¹¹ The UKPDS-OM2 risk equations are widely used in diabetes simulation models, and have been shown to accurately predict results for the population in which it was developed as well as other diabetes populations.^{11,107,108} The UKPDS-OM2 complications (13 risk equations) include CHF, ischemic heart disease (IHD), first MI for females, first MI for males, subsequent MI, first stroke, subsequent stroke, blindness, foot ulcer, first amputation without prior ulcer, first amputation with prior ulcer, subsequent amputation, and ESRD.¹¹ In the microsimulation, patients were able to experience multiple and concurrent complications during each modeled year. The UKPDS-OM2 mortality risk equations predict that previous T2DM-related complications (except foot ulcer and blindness) increase the probability of death. The four mutually exclusive mortality risk equations were death without history of complication(s), death in the year of a clinical event, death in subsequent year of prior event(s), and death with history of clinical event(s).¹¹

The effect of included add-on therapies on intermediate outcomes, including several of the time-varying risk factors mentioned above, were applied to the patients and the treatment effect was assumed to persist until patient death or therapy discontinuation. We estimated the effect of included add-on therapies on cardiovascular and renal outcomes in the base case via the reductions in HbA1c, weight, SBP, and LDL, allowing changes in intermediate outcomes to be associated with longer-term outcomes. We also applied hazard ratios from the placebo-controlled cardiovascular outcomes trials for the add-on therapies.

Treatment Discontinuation and Insulin Uptake

We applied a single discontinuation rate due to any cause from the empagliflozin extension trial EMPA-REG EXTEND to all therapies, as long-term discontinuation data was not available for

injectable semaglutide or tirzepatide, at the end of cycle 1. Patients discontinuing their primary modeled treatment were assumed to transition to insulin therapy. This choice was made to facilitate head-to-head comparator evaluations as opposed to evaluating differences in multiple potential treatment pathways. All patients who discontinued used insulin in addition to background treatment for the remainder of the model time horizon. After cycle 1, we also assumed that patients initiated insulin (in addition to their add-on treatment) if their HbA1c reached 8.5 or above. Insulin treatment costs were based on a multivariate prediction model for estimating long-term HbA1c change, weight change, and hypoglycemic events associated with insulin rescue medication.⁶¹ After cycle 1, clinical characteristics for patients pre- and post-insulin were modeled using the equations for HbA1c and weight change,⁶¹ which then influenced the UKPDS-OM2 complication risk equations for those patients. The hypoglycemia equations from the Willis et al. prediction model were not used due to their substantial uncertainty.

Hypoglycemia

Mild, moderate, and severe hypoglycemia were modeled in cycles 2+ based on the previous UKPDS-OM2 adaptation from Laiteerapong et al.⁵³ Patients not yet receiving insulin were assumed to have a 5% probability for a severe hypoglycemic event and a 33% probability for a mild or moderate event each year. Patients receiving insulin were assumed to have a 21% probability of a severe hypoglycemic event and a 52% probability of a mild or moderate hypoglycemic event each year. Patients were assumed to have no more than one mild or moderate hypoglycemic event and one severe hypoglycemic event per year but could have multiple hypoglycemic events during their lifetime.

Atrial Fibrillation and Peripheral Artery Disease

The UKPDS-OM2 equations have coefficients for atrial fibrillation and peripheral vascular (artery) disease but the NHANES patient dataset did not provide this information. Therefore, we utilized age-based cumulative incidence estimates from the US population^{109,110} and (for atrial fibrillation) relative risk estimates based on patients' HbA1c,¹¹¹ to simulate these patient characteristics prior to each microsimulation. Peripheral vascular disease and atrial fibrillation prevalence were modeled independent of existing patient characteristics.

Mortality

The UKPDS-OM2 risk equations predict mortality using four mutually exclusive equations that are stratified based on the person's prior T2DM event history and whether the mortality event is related to a cardiovascular event.¹¹

Utilities

We used consistent health state utility values across treatments evaluated in the model. Separate utilities were used for the year in which a complication occurred and for patient history of each complication, where applicable. Health state utilities were derived from publicly available literature and/or manufacturer-submitted data and applied to the modeled events. We used regression-based estimates for T2DM complications primarily from Shao et al.^{112,113} In Shao et al., the Health Utilities Index Mark 3 (HUI-3) was used to measure health utility in a sample of 8,713 patients from the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial of high cardiovascular disease risk T2DM patients.¹¹³ Lastly, we additively modeled an annual disutility for daily injection of insulin (for patients who progressed) based on Boye et al., who used standard gamble interviews of T2DM patients in Scotland to estimate the utility values for injection-related attributes.¹¹⁴

Table E.3. Utility Values Used in Model

Health event or history	(Dis)Utility	Source
Baseline T2DM Utility	0.800	Shao et al. ¹¹³
Macrovascular complications		
Congestive heart failure event	-0.089	Shao et al. ¹¹³
Congestive heart failure history	-0.041	Shao et al. ¹¹³
Ischemic heart disease history	-0.016	Shao et al. ¹¹³
Myocardial infarction event	-0.042	Shao et al. ¹¹³
Myocardial infarction history	-0.011	Shao et al. ¹¹³
Stroke event	-0.204	Shao et al. ¹¹³
Stroke history	-0.101	Shao et al. ¹¹³
Microvascular complications		
Blindness history	-0.057	Shao et al. ¹¹³
Foot ulcer event	-0.024	Sullivan et al. ¹¹⁵
Amputation event	-0.051	Sullivan et al. ¹¹⁵
Renal disease history	-0.024	Shao et al. ¹¹³
Hypoglycemia		
Hypoglycemia event	-0.036	Shao et al. ¹¹³
Hypoglycemia history	-0.033	Shao et al. ¹¹³

Demographic characteristics		
Age at diagnosis (per year ≥52)	-0.002	Shao et al. ¹¹³
Female	-0.043	Shao et al. ¹¹³
Race (ref = Black)		
Hispanic	-0.045	Shao et al. ¹¹³
Others	-0.010	Shao et al. ¹¹³
White	-0.019	Shao et al. ¹¹³
Current smoker	-0.054	Shao et al. ¹¹³
BMI (per unit ≥32)	-0.007	Shao et al. ¹¹³
Diabetes duration (per year)	-0.005	Shao et al. ¹¹³
<u>Injection-related disutility</u>		
Annual Disutility for Tx Injection	-0.054	Boye et al. ¹¹⁴

BMI: body mass index, Tx: treatment

Economic Inputs

Drug Acquisition Costs

Because tirzepatide is not approved by the FDA, the drug price is not yet available. Based on the [ICER Reference Case](#), we investigated calculation of a placeholder price based on the average of all available once-weekly injectable GLP-1s. However, we uncovered that discounted pricing for once-weekly injectable Bydureon BCise® (exenatide extended-release) is approximately three times that of Ozempic® (semaglutide), suggesting it may be an unsuitable proxy to inform tirzepatide placeholder pricing. Given this, we used the price of Ozempic® (semaglutide) as a placeholder price for tirzepatide and calculated the threshold prices at the standard cost-effectiveness thresholds: \$50,000 through \$200,000 per QALY; we did not calculate cost-effectiveness thresholds in terms of equal value of life years (evLYs) gained given no average increased survival when comparing tirzepatide with injectable semaglutide. We applied the semaglutide drug discount rate to obtain net pricing estimates for tirzepatide.

Health Care Utilization Costs

Table E.4. Cost per T2DM-Related Complication and per Hypoglycemic Event

Incremental Cost in the Year of Event/Diagnosis (per event)	Estimate (2021 USD)*	Source
Year of Event (per event)		Yang ⁶⁴
Congestive Heart Failure	\$34,898	Yang ⁶⁴
Ischemic Heart Disease	\$9,962	Yang ⁶⁴

Myocardial Infarction	\$50,612	Yang ⁶⁴
Stroke	\$26,597	Yang ⁶⁴
Foot Ulcer	\$2,691	Ward ⁶³
Amputation	\$11,330	Ward ⁶³
Hypoglycemia		
Requiring Hospitalization	\$8,563	Yang ⁶⁴
Requiring ER Visit	\$1,643	Yang ⁶⁴
Requiring Glucagon Injection	\$221	Yang ⁶⁴
History of Complication (per year)		
Congestive Heart Failure	\$7,899	Yang ⁶⁴
Ischemic Heart Disease	\$2,386	Ward ⁶³
Myocardial Infarction	\$9,587	Yang ⁶⁴
Stroke	\$5,289	Yang ⁶⁴
Blindness	\$14,534	Yang ⁶⁴
Renal Disease	\$105,394	Yang ⁶⁴
Health Care Use Costs		
Outpatient Visit: Non-insulin	\$584	Laiteerapong ⁵³
Outpatient Visit: Insulin	\$639	Laiteerapong ⁵³

ER: emergency room, ESRD: end-stage renal disease, USD: United States dollars

*Costs inflated to USD 2021 using US government Bureau of Labor Statistics Consumer Price Index

Adverse Event Costs and Disutilities

No serious treatment-related adverse events occurring in greater than 5% of the patient population were observed. Adverse events occurring under that rate generally have minimal effect on the model output and therefore were not included in the model. Overall TRAEs and other reasons for discontinuing treatment were addressed in a global fashion through cardiac events, treatment discontinuation, etc. We also included an annual disutility due to daily insulin injections in the utility section.

Productivity Costs

We sought to include a modified societal perspective in this analysis. However, we found a paucity of data to inform such modeling. Instead, we performed calculations utilizing the model's incremental health gains (QALYs gained) and a measure of the productivity impact of T2DM to approximate the modified societal perspective results. We adjusted Dall et al.'s estimate of the productivity costs of T2DM from 2017 to 2021 US Dollars using the health care component of the Consumer Price Index, resulting in annual productivity costs of \$5842.¹¹⁶ That annual estimate was assumed to be the potential savings associated with each incremental QALY for an intervention versus a comparator. We multiplied the annual productivity estimate times the incremental QALYs to estimate the assumed societal cost savings and combined these assumed productivity-based

savings with the total incremental payer perspective costs in order to calculate cost-effectiveness ratios for the modified societal perspective.

E3. Results

Disaggregated and Undiscounted Base-Case Results

To estimate the overarching lifetime costs and benefits of each active therapy and background treatment alone, the model calculated the average disaggregated values, including the percent of patients experiencing disease-related events, add-on agent costs, background treatment costs, and event-related health care costs. The averages of each of those values, along with 95% credible ranges, are presented in Table E3.1 for each active therapy and background treatment alone. We also calculated cost and outcome variables without discounting, presented in Table E3.2. EvLYs and evLYs gained were not reported given no average increased survival when comparing tirzepatide with injectable semaglutide.

Table E3.1 Disaggregated Results of the Base-Case Model

	Tirzepatide		Semaglutide		Empagliflozin		Background Tx	
	Mean	95% CR	Mean	95% CR	Mean	95% CR	Mean	95% CR
Costs								
Total Cost	\$306,211	(\$275,083 - \$338,644)	\$309,150	(\$280,007 - \$339,362)	\$275,748	(\$247,587 - \$304,617)	\$261,792	(\$234,473 - \$290,804)
Add-on Agent	\$40,480	(\$38,160 - \$42,885)	\$41,222	(\$38,813 - \$43,527)	\$11,967	(\$11,280 - \$12,672)	\$0	(\$0 - \$0)
Background Tx	\$4,155	(\$3,951 - \$4,364)	\$4,229	(\$4,027 - \$4,429)	\$4,070	(\$3,875 - \$4,272)	\$3,703	(\$3,516 - \$3,897)
Insulin	\$8,189	(\$7,132 - \$9,265)	\$10,448	(\$9,294 - \$11,625)	\$13,285	(\$12,071 - \$14,572)	\$14,544	(\$13,287 - \$15,842)
Health Care	\$5,627	(\$5,342 - \$5,916)	\$5,760	(\$5,481 - \$6,042)	\$5,600	(\$5,326 - \$5,874)	\$5,134	(\$4,871 - \$5,412)
CHF	\$18,305	(\$15,537 - \$21,410)	\$21,527	(\$18,593 - \$24,776)	\$16,389	(\$13,701 - \$19,236)	\$18,085	(\$15,245 - \$21,086)
IHD	\$3,263	(\$2,629 - \$3,957)	\$3,444	(\$2,774 - \$4,125)	\$3,280	(\$2,618 - \$3,978)	\$2,905	(\$2,332 - \$3,554)
MI	\$22,413	(\$18,947 - \$26,038)	\$24,950	(\$21,188 - \$28,950)	\$27,207	(\$23,489 - \$31,279)	\$25,950	(\$22,319 - \$29,797)
Stroke	\$10,133	(\$8,147 - \$12,358)	\$11,872	(\$9,645 - \$14,433)	\$13,038	(\$10,742 - \$15,569)	\$12,480	(\$10,322 - \$14,848)
Blindness	\$4,566	(\$2,591 - \$6,951)	\$5,392	(\$3,258 - \$7,778)	\$5,448	(\$3,294 - \$7,875)	\$5,133	(\$3,107 - \$7,546)
Foot Ulcer	\$307	(\$195 - \$453)	\$384	(\$253 - \$542)	\$424	(\$280 - \$599)	\$408	(\$267 - \$577)
Amputation	\$2,486	(\$1,847 - \$3,286)	\$3,034	(\$2,272 - \$3,953)	\$3,276	(\$2,465 - \$4,242)	\$2,995	(\$2,250 - \$3,867)
Renal Disease	\$174,835	(\$147,013 - \$204,294)	\$164,363	(\$139,417 - \$191,177)	\$158,121	(\$133,761 - \$183,606)	\$156,922	(\$132,560 - \$183,590)

Hypoglycemia	\$11,452	(\$10,345 - \$12,573)	\$12,527	(\$11,438 - \$13,723)	\$13,644	(\$12,471 - \$14,897)	\$13,534	(\$12,360 - \$14,781)
Health Outcomes								
QALYs	4.90	(4.68 - 5.12)	4.85	(4.64 - 5.05)	4.60	(4.40 - 4.79)	4.13	(3.95 - 4.33)
Life Years	9.36	(8.91 - 9.83)	9.53	(9.08 - 9.97)	9.17	(8.73 - 9.61)	8.34	(7.93 - 8.77)
Complications								
CHF	29.9%	(25.6% - 34.4%)	34.8%	(30.5% - 39.3%)	27.2%	(23.0% - 31.5%)	30.1%	(25.6% - 34.6%)
IHD	14.5%	(11.4% - 18.1%)	15.3%	(11.9% - 18.9%)	14.6%	(11.1% - 18.1%)	13.3%	(10.1% - 16.5%)
1st MI	27.3%	(23.5% - 31.3%)	29.2%	(25.1% - 33.1%)	32.4%	(28.4% - 36.7%)	34.0%	(29.7% - 38.2%)
Subs. MI	3.7%	(2.1% - 5.9%)	4.4%	(2.3% - 6.7%)	5.4%	(3.1% - 8.0%)	4.8%	(2.8% - 7.2%)
1st Stroke	23.3%	(19.4% - 27.4%)	25.7%	(21.4% - 29.7%)	28.8%	(24.5% - 33.3%)	30.0%	(25.8% - 34.4%)
Subs. Stroke	9.6%	(4.7% - 17.3%)	12.7%	(6.2% - 22.5%)	13.7%	(7.2% - 23.3%)	11.3%	(5.9% - 19.1%)
Blindness	8.6%	(5.9% - 11.6%)	10.0%	(7.0% - 12.9%)	10.2%	(7.2% - 13.2%)	9.9%	(7.0% - 12.9%)
Foot Ulcer	18.2%	(11.1% - 28.4%)	23.0%	(14.5% - 34.6%)	24.7%	(15.8% - 36.7%)	22.6%	(14.5% - 33.3%)
1st Amp, No Ulcer	16.1%	(12.7% - 19.9%)	18.1%	(14.5% - 22.0%)	19.0%	(15.2% - 22.7%)	17.8%	(14.2% - 21.7%)
1st Amp, Ulcer	3.8%	(2.1% - 5.7%)	4.4%	(2.6% - 6.5%)	4.6%	(2.6% - 6.7%)	4.3%	(2.3% - 6.5%)
Subs. Amp	17.2%	(8.5% - 30.2%)	23.0%	(12.1% - 39.0%)	24.2%	(12.9% - 40.3%)	19.1%	(10.1% - 31.8%)
Renal Disease	15.9%	(12.7% - 19.1%)	12.8%	(9.8% - 15.8%)	12.5%	(9.3% - 15.8%)	14.5%	(11.4% - 17.8%)

CHF: congestive heart failure, IHD: ischemic heart disease, MI: myocardial infarction, QALY: quality-adjusted life year, Tx: treatment

Table E3.2 Undiscounted Results from Base Case

	Tirzepatide		Semaglutide		Empagliflozin		Background Tx	
	Mean	95% CR	Mean	95% CR	Mean	95% CR	Mean	95% CR
Undiscounted Costs								
Total Cost	\$407,979	(\$362,068 - \$455,506)	\$413,490	(\$369,729 - \$458,206)	\$366,247	(\$323,939 - \$407,232)	\$340,555	(\$300,422 - \$382,769)
Add-on Agent	\$52,635	(\$49,107 - \$56,364)	\$53,698	(\$50,057 - \$57,276)	\$15,447	(\$14,397 - \$16,522)	\$0	(\$0 - \$0)
Background Tx	\$5,417	(\$5,108 - \$5,750)	\$5,524	(\$5,214 - \$5,847)	\$5,269	(\$4,962 - \$5,583)	\$4,699	(\$4,414 - \$4,998)
Insulin	\$12,642	(\$10,885 - \$14,446)	\$15,923	(\$13,978 - \$17,957)	\$19,342	(\$17,333 - \$21,440)	\$20,180	(\$18,228 - \$22,265)
Health Care	\$7,370	(\$6,928 - \$7,835)	\$7,561	(\$7,104 - \$8,005)	\$7,283	(\$6,853 - \$7,719)	\$6,541	(\$6,146 - \$6,961)
CHF	\$26,019	(\$21,888 - \$30,505)	\$30,513	(\$26,149 - \$35,279)	\$23,140	(\$19,231 - \$27,224)	\$24,576	(\$20,608 - \$28,775)
IHD	\$4,461	(\$3,521 - \$5,483)	\$4,750	(\$3,744 - \$5,793)	\$4,467	(\$3,500 - \$5,535)	\$3,843	(\$3,029 - \$4,767)

MI	\$30,910	(\$25,854 - \$36,098)	\$34,816	(\$29,182 - \$40,836)	\$37,275	(\$31,745 - \$43,155)	\$34,341	(\$29,226 - \$39,737)
Stroke	\$14,443	(\$11,489 - \$17,976)	\$17,158	(\$13,695 - \$21,383)	\$18,509	(\$15,025 - \$22,631)	\$16,993	(\$13,950 - \$20,538)
Blindness	\$7,545	(\$4,280 - \$11,525)	\$9,017	(\$5,415 - \$13,139)	\$8,926	(\$5,336 - \$13,095)	\$8,068	(\$4,824 - \$11,941)
Foot Ulcer	\$490	(\$297 - \$764)	\$618	(\$390 - \$934)	\$666	(\$423 - \$989)	\$608	(\$386 - \$903)
Amputation	\$4,195	(\$3,017 - \$5,889)	\$5,160	(\$3,699 - \$7,127)	\$5,412	(\$3,928 - \$7,389)	\$4,668	(\$3,404 - \$6,264)
Renal Disease	\$225,970	(\$185,935 - \$268,935)	\$211,343	(\$174,801 - \$250,455)	\$201,883	(\$166,385 - \$238,699)	\$198,108	(\$163,818 - \$236,359)
Hypoglycemia	\$15,881	(\$14,212 - \$17,601)	\$17,410	(\$15,719 - \$19,203)	\$18,629	(\$16,864 - \$20,456)	\$17,930	(\$16,227 - \$19,759)
Survival Time-related Outcomes								
Undiscounted Time on Tx (Years)	11.33	(10.60 - 12.13)	11.56	(10.78 - 12.33)	11.02	(10.28 - 11.77)	NA	NA
Undiscounted Life Years	12.20	(11.52 - 12.95)	12.45	(11.74 - 13.15)	11.87	(11.19 - 12.57)	10.59	(9.96 - 11.25)


CHF: congestive heart failure, IHD: ischemic heart disease, MI: myocardial infarction, QALY: quality-adjusted life year, Tx: treatment

E4. Sensitivity Analyses

Additionally, we ran one-way sensitivity analyses to identify the key drivers of model outcomes, using available measures of parameter uncertainty (i.e., standard errors) or reasonable ranges for each input. In order to efficiently operationalize the one-way sensitivity analysis within the framework of the patient-level microsimulation, we fixed the parameter values for all non-patient-level inputs and then performed 10 UKPDS equation simulations for each of 387 NHANES patients for each parameter's low and high value, for each treatment, in order to produce an estimate of uncertainty for each high and low value of each parameter. Therefore, each one-way sensitivity analysis output value represents the average impact over the 10 simulations of the 387 individual patient simulations. Individual one-way sensitivity analyses were performed for incremental costs and incremental QALYs in order to isolate the impact of the inputs on those individual outcomes. The results of these one-way sensitivity analyses are presented as tornado diagrams.

Here we present comparisons between tirzepatide and empagliflozin, and tirzepatide and background therapy alone. The tornado diagram for incremental costs was truncated to include inputs with at least a \$500 difference between the high and low value. The tornado diagram for incremental QALYs was truncated to include inputs with more than a 0.01 QALY difference between the high and low value.


Figure E.1. Incremental Costs Tornado Diagram (Tirzepatide vs. Empagliflozin)*



CHF: congestive heart failure, HbA1c: Hemoglobin A1c/glycosylated hemoglobin, HR: hazard ratio, MACE: major adverse cardiovascular event, Tx: treatment


*Using a placeholder price for tirzepatide

Figure E.2. Incremental QALYs Tornado Diagram (Tirzepatide vs. Empagliflozin)



CHF: congestive heart failure, HbA1c: Hemoglobin A1c/glycosylated hemoglobin, HR: hazard ratio, MACE: major adverse cardiovascular event, Tx: treatment


Figure E.3. Incremental Costs Tornado Diagram (Tirzepatide vs. Background Therapy Alone)*



CHF: congestive heart failure, HR: hazard ratio, MACE: major adverse cardiovascular event, Tx: treatment

*Using a placeholder price for tirzepatide


Figure E.4. Incremental QALYs Tornado Diagram (Tirzepatide vs. Background Therapy Alone)



HR: hazard ratio, Tx: treatment

*Using a placeholder price for tirzepatide

Figure E.5. Base-Case Cost-Effectiveness Acceptability Curves for Tirzepatide vs. Semaglutide, Empagliflozin, and Background Therapy Alone



E5. Scenario Analyses

We ran the following separate scenario analyses to understand the impact of some of our assumptions within the model when comparing tirzepatide plus background therapy to background therapy alone: we shortened the time horizon to 10 years; we set tirzepatide's CV and renal event adjustment equal to that of semaglutide; we gave no adjustment to tirzepatide's CV and renal event estimations; we set the insulin initiation threshold due to advanced HbA1c levels to both 8% and to 9% as opposed to the base case 8.5%; we ran the model without applying risk factor progression for HbA1c and weight; and we removed utility benefit for BMI change. For each of these scenarios, the results presented represent the mean and 95% credible range from probabilistic sensitivity analyses comprised of 500 simulations of each of the 387 people from NHANES with T2DM.

Table E.5. Scenario analyses incremental results for tirzepatide added to background therapy vs. background therapy alone, mean (95% credible range)*

Scenario Description	Incremental Cost per QALY Gained	Incremental Cost per Life Year Gained
Base-Case Scenario	\$58,000	\$44,000
	(\$10,900 to \$99,100)	(\$10,400 to \$82,500)
10-year Time Horizon	\$49,000	\$59,000
	(-\$10,300 to \$101,500)	(-\$54,800 to \$223,500)
Tirzepatide CV and Renal Benefit with No Hazard Ratio Adjustment	\$58,000	\$57,000
	(-\$13,000 to \$115,000)	(-\$20,000 to \$278,500)
Insulin When HbA1c=8%	\$55,000	\$42,000
	(\$12,000 to \$94,300)	(\$11,100 to \$77,200)
Insulin When HbA1c=9%	\$64,000	\$46,000
	(\$19,400 to \$111,700)	(\$15,400 to \$87,400)
No Willis Equations (no risk factor progression assumptions)	\$60,000	\$47,000
	(\$3,500 to \$106,000)	(\$3,900 to \$106,200)
No disutility due to BMI	\$62,000	\$42,000
	(\$13,200 to \$109,100)	(\$10,100 to \$82,900)

BMI: body mass index, CV: cardiovascular, HbA1c: hemoglobin levels, QALY: quality-adjusted life-year

*Using a placeholder price for tirzepatide

Modified Societal Perspective Scenario Estimation, using Missing Data Assumptions

We used a placeholder price for tirzepatide and assumed that the incremental QALYs gained by tirzepatide were associated with avoiding productivity losses associated with T2DM. Our calculation of the estimated value of tirzepatide compared to background therapy alone under this modified societal perspective using missing data assumptions, resulted in incremental cost-effectiveness ratios of \$52,600 per QALY gained and \$39,200 per life year gained.

E6. Model Validation

Model validation followed standard practices in the field. We tested all mathematical functions in the model to ensure they were consistent with the report (and in supplemental materials). We also conducted sensitivity analyses with null input values to ensure the model was producing findings consistent with expectations. Further, independent modelers tested the mathematical functions in the model as well as the specific inputs and corresponding outputs.

We also performed external validation by running the model for 2- and 3-year outcomes without hazard ratio adjustments and presented the results next to available cardiovascular outcome trial data (Table E6). We shortened the time horizon to more closely resemble the conditions in the

trials and to provide insight into clinical event rates and risk factor progression against the base-case lifetime model.

Table E6. 2-year and 3-year Time Horizon Model Event Estimates Compared to Cardiovascular Outcome Trial Output

		Semaglutide		Placebo/Background Only		Empagliflozin		Placebo/Background Only	
Trial Outcomes	Model Outcome Used	SUSTAIN-6	2-year Model	SUSTAIN-6	2-year Model	EMPA-REG - OUTCOME	3-year Model	EMPA-REG - OUTCOME	3-year Model
Composite Outcome	Summed MACE w/o double counting CV death	6.6%	8.1%	8.9%	12.4%	10.5%	14.9%	12.1%	18.0%
All-Cause Mortality	Ratio of Undiscounted LYs to Time Horizon	3.8%	3.0%	3.6%	3.5%	5.7%	6.0%	8.3%	6.7%
CV Mortality	CV Death (UKPDS Eq)	2.7%	1.5%	2.8%	2.9%	3.7%	3.1%	5.9%	4.0%

Prior Economic Models

In our review of the literature, we found no cost-effectiveness model that compared tirzepatide to other T2DM treatment strategies. Our focus in this section is therefore to review and contrast the methodologies used in the modeling of T2DM treatment strategies. The numerous available strategies and pathways available for the treatment of T2DM have led to the development and publication of several cost-effectiveness analyses and modeling exercises in the past few decades, many of which highlight the comparators used in this analysis.^{55,56,108,117,118} Such analyses include both cohort and microsimulation models. For the purposes of this report, we have limited the comparison to other published microsimulation models, specifically the following: 1) the UKPDS OM2 (a model predicting health outcomes in T2DM)¹¹, 2) a microsimulation cost utility model by Laiteerapong et al.⁵³, 3) the model published in ICER's 2019 report of oral semaglutide on T2DM outcomes, of which this model is an adaptation,^{40,54} and 4) models published since the previous ICER T2DM report using the IQVIA Core Diabetes Model assessing our included comparators.^{119,120}

The UKPDS OM2 is an update of the original UKPDS Outcomes Model 1 (OM1), also a patient simulation model that predicts health outcomes of patients with T2DM. The UKPDS OM2 re-estimated the original seven risk equations in the UKPDS OM1 over a longer time-horizon plus additional risk equations for other complications such as diabetic ulcer. Additionally, it also included new risk equations for all-cause mortality in T2DM patients. Our model applied the updated UKPDS OM2 risk equations (developed for the UK population) to a US-specific population that was derived from 2013-2014, 2015-2016, and 2017-2018 NHANES survey data

on 387 patients that fit the baseline characteristics of patients with uncontrolled T2DM taking metformin but not another type of add-on diabetes therapy. Our model also assumed risk factor progression by applying published risk factor progression equations for weight change and HbA1c to the simulation.⁶¹

A key comparison of our model is to the one by Laiteerapong et al.⁵³ Risks of different levels of hypoglycemia in our model are based on the hypoglycemia risk module developed by Laiteerapong et al. in their patient-level, Monte Carlo-based Markov model. Both models use the baseline UKPDS OM2 risk equations in modeling health outcomes, but for T2DM patients in the U.S. Considering differences between our model and the one by Laiteerapong et al., a key difference is unlike their or any other model, we specifically estimate risk for MACE and renal events using HRs derived from an NMA of key trials in our population. This HR is applied to the UKPDS OM2-derived risk equations for specific treatment strategies included in our model. As in ICER's 2019 T2DM review, we believe this approach better accounts for treatment-specific effects on critical outcomes such as MACE and renal outcomes in T2DM patients than the approach used by Laiteerapong et al. or others, who used the unmodified risk equations from the UKPDS OM2. Other differences between the two models include the NHANES population; we used a more recent population compared to theirs which results in slightly different patient characteristics. As mentioned earlier we modeled a new, previously unmodeled treatment (tirzepatide), our treatment costs were different, we limited patients to only receive insulin after discontinuation or HbA1c increase, and we used an adapted approach to applying utility values when individuals had a history of an event.

Another key comparison of our model is to ICER's 2019 report reviewing oral semaglutide.⁴⁰ Other T2DM risk models were considered when preparing the 2022 report on tirzepatide, however the UK-PDS OM2 equations remained the most feasible option and were deemed effective as compared to other available diabetes models when used with CVOT calibration.¹²¹ Updates to the 2019 review included a different set of active treatment comparators and an updated network meta-analysis of treatment effect to reflect tirzepatide clinical trials, additional NHANES patient cohorts, and the addition of change in LDL as an intermediate outcome in the first cycle.

Finally, there have been several additional models published since the publication of ICER's 2019 T2DM report which assess one or more of our comparator therapies, mostly set in countries other than the United States.^{119,120,122-124} As an example, Ramos and colleagues published a cost-effectiveness model of empagliflozin compared to sitagliptin and liraglutide in 2021 from a Chinese health care system perspective using the IQVIA Core Diabetes Model (CDM).¹¹⁹ Like our model, they calibrated their equations to reflect available CVOT outcomes and used indirect treatment comparisons due to lack of head-to-head trial data. They differed from our

approach in assuming no lifetime treatment but a switch to insulin as rescue therapy at an 8.5% HbA1c threshold. Results were not comparable due to the therapies included and the baseline Chinese population. Another study by Capehorn and colleagues likewise used the IQVIA CDM to assess injectable semaglutide versus empagliflozin in a 2021 paper from a United Kingdom health care perspective, also using indirect comparisons via network meta-analysis.¹²⁰ Their analysis, like ours, assumed treatment effects in the first year of analysis, then used risk factor progression equations based on equations published by Willis et al.,⁶¹ and had patients initiate insulin therapy when their HbA1c reached a designated threshold. They concluded that semaglutide conferred a lifetime improvement of 0.23 QALYs over empagliflozin, which was comparable to our estimate of 0.25 incremental QALYs for those two therapies.

F. Potential Budget Impact: Supplemental Information

F1. Methods

We used results from the same model employed for the cost-effectiveness analyses to estimate total potential budget impact. These results from the cost-effectiveness analyses include the costs of add-on antidiabetic agent (i.e., tirzepatide, injectable semaglutide, empagliflozin), health care resource utilization offsets, and averted health care event offsets. In patients who subsequently progressed to insulin, those insulin costs were attributed to the previously discontinued therapy (e.g., tirzepatide [plus background therapy]). Potential budget impact was defined as the total differential cost of using each new therapy rather than relevant existing therapy for the treated population, calculated as differential health care costs (including drug costs) minus any offsets in these costs (for instance, due to offsets in major adverse cardiovascular events). All costs were undiscounted and estimated over one- and five-year time horizons. The five-year timeframe was of primary interest, given the potential for cost offsets to accrue over time and to allow a more realistic impact on the number of patients treated with the new therapy.

To estimate the size of the potential candidate population for treatment, we used inputs for the total US adult population size (~265 million)⁶⁶, overall T2DM prevalence (14.6%)⁶⁷, proportion of patients with diagnosed T2DM (76.7%)⁶⁷, and the proportion of patients having failed background therapy and considering a second-line treatment and thus eligible for tirzepatide (16.2%).⁶⁸ Applying these sources results in estimates of 4,800,000 eligible patients in the US. For the purposes of this analysis, we assumed that 20% of these patients would initiate tirzepatide treatment in each of the five years, or approximately 960,000 patients per year.

Comparators in the budget impact model included injectable semaglutide, empagliflozin, each added to background therapy, as well as background therapy alone. Conventional therapy consisted of metformin and/or sulfonylureas. Market shares were derived from analyst projections based on primary market research, company reports, and key opinion leader surveys.⁶⁹ We set the initial market shares for injectable semaglutide equal to a calculated market share for the GLP-1 RA class (15.1%), while the market share for empagliflozin was set to a calculated market share for the SGLT-2 inhibitor class (12.7%). The remaining market share (72.2%) was attributed background therapy. Market share was captured in proportion to the comparators' initial market shares at a rate of 20% per year, with all patients having switched to tirzepatide by the end of five-year of the time horizon.

Additionally, we used an estimate of net price (using injectable semaglutide's price as a placeholder), and the three threshold prices (at \$50,000, \$100,000, and \$150,000 per QALY versus injectable semaglutide) in our estimates of tirzepatide's potential budget impact. Injectable semaglutide was chosen as the comparator for threshold price determination in order to align with the HBPBs reported from the cost-effectiveness analyses. ICER's methods for estimating potential budget impact are described in detail elsewhere and have recently been updated.^{125,126} The intent of our revised approach to budgetary impact is to document the percentage of patients that could be treated at selected prices without crossing a budget impact threshold that is aligned with overall growth in the US economy.


Using this approach to estimate potential budget impact, we then compared our estimates to an updated budget impact threshold that represents a potential trigger for policy mechanisms to improve affordability, such as changes to pricing, payment, or patient eligibility. As described in [ICER's methods presentation](#), this threshold is based on an underlying assumption that health care costs should not grow much faster than growth in the overall national economy. From this foundational assumption, our potential budget impact threshold is derived using an estimate of growth in US gross domestic product (GDP) +1%, the average number of new drug approvals by the FDA over the most recent five-year period for which data were available, and the contribution of spending on retail and facility-based drugs to total health care spending over the most recent five-year period for which data were available.

For 2021-2022, the five-year annualized potential budget impact [threshold](#) that should trigger policy actions to manage access and affordability is calculated to total approximately \$734 million per year for new drugs.

F2. Results

Figure F2.1 illustrates the cumulative per-patient budget impact calculations for tirzepatide added to background therapy compared to a mixed market basket consisting of injectable semaglutide, empagliflozin, and background therapy based on the placeholder net price and net prices used within the cost-effectiveness analysis. Switching of eligible adult T2DM patients to tirzepatide resulted in an average potential budgetary impact of approximately \$1,300 per patient per year when assuming our standard uptake of 20% per year. We suggest caution in interpreting the potential budget impact of tirzepatide due to the placeholder annual net price assumed.

Figure F2.1. Cumulative Annual Per Patient Treated with Tirzepatide at a Placeholder Price of \$4,643.50 per Year*



*Placeholder price was assumed. Interpret findings with caution.

Table F2.1 illustrates the average annual per-patient budget impact results in more detail, for tirzepatide’s placeholder price (\$4,643.50* per year), and the threshold prices to reach \$50,000, \$100,000, and \$150,000 per QALY (\$5,200, \$5,500, and \$5,700, per year, respectively) compared to injectable semaglutide added to background therapy.

Table F2.1. Average Annual Per-Patient Budget Impact Calculations Over a Five-year Time Horizon

	Average Annual Per-Patient Budget Impact for Each Calculated Price Point			
	Placeholder Price*	\$50,000/QALY	\$100,000/QALY	\$150,000/QALY
Tirzepatide and Background Therapy vs. Injectable Semaglutide, Empagliflozin, and Background Therapy	\$1,300	\$1,800	\$2,000	\$2,300

QALY: quality-adjusted life year

*This is an unvalidated placeholder price that is assumed to be equal to the discounted price for injectable semaglutide; injectable semaglutide price is sourced from Red Book, with corresponding discounts used to derive a net price being sourced from SSR Health, LLC ⁶²