

## **Brensocatib for Non-Cystic Fibrosis Bronchiectasis**

### **Revised Background and Scope**

**April 9, 2025**

### **Background**

Bronchiectasis is a chronic, progressive lung disease that affects breathing and coughing. Many patients have chronic daily cough, often bringing up mucus and saliva, and exacerbations that may involve worsening of these chronic symptoms along with shortness of breath, fatigue, and worsening lung function.<sup>1</sup> Bronchi, the small tubes that bring air to the lungs, often become enlarged and colonized with bacteria. Bronchiectasis is characterized by a “vicious cycle” of chronic infection, structural lung changes, inflammation, and deterioration in mucociliary clearance (i.e., the way that the body clears the lung of mucus). Given the complexity and synergy between these processes, some have proposed calling such an interaction a “vicious vortex.”<sup>4</sup> The coughing and mucus production make bronchiectasis similar to chronic obstructive pulmonary disease (COPD). However, unlike COPD, bronchiectasis is a disorder of the bronchi enlarging rather than collapsing.<sup>5</sup> In bronchiectasis, the enlargement of the bronchi can be seen on computed tomography (CT) scans of the lungs. The diagnosis of bronchiectasis as a clinical syndrome also requires the appearance of typical symptoms related to breathing and coughing. Bronchiectasis can result in significant negative effects on quality of life due to impaired social and physical functioning related to chronic cough and worsening lung function. Bronchiectasis is common among individuals with cystic fibrosis, a relatively rare condition. However, evidence has emerged that in some cases, the type of bronchiectasis that occurs with cystic fibrosis responds to different treatments than other types of bronchiectasis. In that context, bronchiectasis is often called non-cystic fibrosis bronchiectasis (NCFB).<sup>6</sup> This document focuses on the larger population of individuals with NCFB.

In the United States, NCFB is relatively common and increasingly recognized. An estimated 350,000 to 500,000 adults in the United States have NCFB, with 70,000 new cases emerging annually.<sup>7</sup> Older estimates previously suggested lower prevalence.<sup>8,9</sup> The increasing measured prevalence may reflect an actual increase in prevalence of NCFB and/or increased detection, particularly with higher use of high-resolution CT scans. The prevalence increases substantially with age, with large increases starting around age 45. Individuals with NCFB are often misdiagnosed with other

conditions, such as COPD. More proactive CT scan imaging can often clarify that NCFB is the underlying cause of symptoms.<sup>7</sup> The annual cost of NCFB care in the United States exceeds \$14 billion per year, about \$2 billion of which is for hospitalizations. Other important contributions to cost include laboratory testing, post-acute services, medical equipment, and outpatient care. Adjusted for population size, the cost of NCFB care in the United States is two to three times greater than in comparable countries.<sup>10</sup>

Among those with NCFB in the United States, about two-thirds have at least one exacerbation per year and about one-third have three or more exacerbations per year. Just under half have an exacerbation that requires either intravenous antibiotics and/or inpatient hospitalization.<sup>11</sup> Risk markers for prognosis in bronchiectasis include prior hospitalizations and exacerbations, severity of shortness of breath, lower forced expiratory volume in one second (amount of air the lung can force out in one second), colonization with bacteria (including the very resistant *Pseudomonas aeruginosa* and nontuberculous mycobacteria [NTM] such as *Mycobacterium avium*), the number of parts of the lung involved, age, and body-mass index.<sup>12</sup> Bronchiectasis can be caused by a prior lung infection, entry of food or liquid into the lung, inherited genetic disorders, structural disorders of the windpipe, problems with the immune system, and other conditions; in many cases, however, a specific cause cannot be found.<sup>1</sup>

Clinical guidelines for NCFB are based on generally low-quality evidence and interventions are directed at different components of the “vicious vortex.” For stable outpatients, regular airway clearance therapy at home after using humidification with saline nebulizers, hand-held oscillatory appliances and vests, chest physical and percussion therapy, and exercise are recommended. These therapies can take hours, require substantial effort and planning, and restrict daily activities, work, and travel. For stable outpatients with three or more exacerbations per year, long-term inhaled and oral antibiotics are recommended. These prolonged antibiotic courses can increase the risk of antimicrobial resistance, both for individuals with NCFB and larger populations. Pulmonary rehabilitation is recommended for individuals who are substantially limited by shortness of breath. In some severe cases, surgical resection of part of the lung or even lung transplantation is sometimes considered.<sup>13</sup> Unlike for cystic fibrosis, there are no treatments that are specifically approved for NCFB and there are not yet practice guidelines specific to the United States. As such, there is a substantial unmet need for patients with NCFB.

Brensocatib (Insmmed Incorporated) is a small molecule reversible inhibitor of dipeptidyl peptidase 1 (DPP1) that reduces signaling of neutrophils, which is thought to reduce the inflammation that is one of the drivers of the “vicious vortex.” The drug is delivered once daily via oral tablet. The manufacturer has announced the submission of a new drug application with the US Food and Drug Administration (FDA), with a decision expected August 12, 2025.<sup>14</sup>

## Stakeholder Input

This draft scoping document was developed through outreach and engagement of diverse stakeholders, including patients and their families, clinicians, researchers, and manufacturers of the agents of focus in this review. This document incorporates feedback gathered during preliminary calls with stakeholders and open input submissions from the public. ICER looks forward to continued engagement with stakeholders throughout its review and encourages comments to refine our understanding of the clinical effectiveness and value of preventive treatments.

Discussions with clinical experts emphasized the substantial burden that patients with NCFB face, often coughing with mucus production every day and spending 30-60 minutes or more daily with home respiratory therapy and nebulizers. There is hope that a new agent could reduce this amount of time in home self-care and potentially help individuals feel better later in the day. There is a lot of interest in brensocatib, since no FDA approved treatment exists and there have been a lot of therapeutic failures for NCFB (including the recognition that many cystic fibrosis treatments do not work well for NCFB).

Clinical experts also shared that they expect brensocatib to be used for outpatients with NCFB, potentially indefinitely. They believe that matching treatment to those most likely to benefit would be important, potentially aided with risk markers. How brensocatib affects utilization will be important. One clinical expert perceived different thresholds for hospital admission during exacerbations. For example, at highly specialized centers, the proportion of exacerbations that result in hospitalization may be lower than at other centers. Often outpatient visits occur every three to six months, with more frequent visits and interventions for those whose NCFB is not well controlled. The diagnosis of NCFB is clinical with supporting radiographic evidence, and there is no specific test. The apparent increase in prevalence is likely related in part to increased awareness and high-resolution imaging.

Our direct conversations with patients as well as academic reports based on patient perspectives suggest that patients and caregivers are concerned about symptoms including persistent cough, fatigue, shortness of breath, and mucus production. Misdiagnosis and delayed diagnosis are common, and associated with additional emotional burdens of anger, confusion, frustration, and anxiety.<sup>15</sup> Patients report that their quality of life is affected by social embarrassment, sleep disturbance, anxiety, and the need to modify daily and future activities.<sup>16</sup> Improving respiratory symptoms, reducing the frequency of exacerbations, and reducing progressive lung damage are key hopes for patients. Some validated quality of life tools have been developed for NCFB, and include a range of physical and emotional functioning, treatment burden, and other aspects of quality of life specific to NCFB.<sup>17</sup> Patients also report difficulty finding physicians and respiratory therapists with substantial experience in NCFB and frustration explaining their condition to others and connecting with supportive resources, given that NCFB is less well known than other lung disorders.<sup>18</sup> Given

fatigue and coughing, as well as the need for airway clearance devices , patients with NCFB report substantial limitations in choices of jobs, travel, and social activities.

## Report Aim

This project will evaluate the health and economic outcomes of brensocatib for NCFB. The ICER value framework includes both quantitative and qualitative comparisons across treatments to ensure that the full range of benefits and harms – including those not typically captured in the clinical evidence such as innovation, public health effects, reduction in disparities, and unmet medical needs – are considered in the judgments about the clinical and economic value of the interventions.

## Scope of Clinical Evidence Review

The proposed scope for this assessment is described on the following pages using the PICOTS (Population, Intervention, Comparators, Outcomes, Timing, and Settings) framework. Evidence will be abstracted from randomized controlled trials as well as high-quality systematic reviews; high-quality comparative cohort studies will be considered, particularly for long-term outcomes and uncommon adverse events. Our evidence review will include input from patients and patient advocacy organizations, data from regulatory documents, information submitted by manufacturers, and other grey literature when the evidence meets ICER standards (for more information, see ICER's [grey literature policy](#)).

All relevant evidence will be synthesized qualitatively or quantitatively. Wherever possible, we will seek out head-to-head studies of the interventions and comparators of interest. Data permitting, we will also consider combined use of direct and indirect evidence in network meta-analyses of selected outcomes. Full details regarding the literature search, screening strategy, data extraction, and evidence synthesis will be provided after the revised scope in a research protocol published on the Open Science Framework website (<https://osf.io/7awvd/>).

## Populations

The population of interest for this review is adolescents and adults with non-cystic fibrosis bronchiectasis (NCFB).

Data permitting, we will evaluate the evidence for treatment effect modification by subpopulations defined by:

- Sociodemographic factors (e.g., sex, age, race, ethnicity)
- Comorbidities (e.g., asthma, COPD)
- Pulmonary exacerbation rate in prior 12 months
- Chronic antibiotic use
- *Pseudomonas aeruginosa* culture status (positive, negative)
- Nontuberculous mycobacteria status (positive, negative)
- Bronchiectasis Severity Index Score

## Interventions

The intervention of interest of this review is:

- Brensocatib (Insmed Incorporated)

## Comparators

Data permitting, we intend to compare brensocatib as an add-on therapy to current usual care, which may include antibiotics, mucolytics, pulmonary rehabilitation, and airway clearance devices, versus usual care alone.

## Outcomes

The outcomes of interest are described in the list below.

- Patient-Important Outcomes
  - Pulmonary exacerbations
  - Exacerbation-related hospitalization or emergency room visit
  - Quality of life (e.g., quality of life-bronchiectasis questionnaire)
  - Lung function (e.g., FEV<sub>1</sub>)
  - Fatigue
  - Use of rescue medications, such as bronchodilators
  - All-cause mortality
- Other Outcome
  - Changes in biomarkers (e.g., neutrophil elastase)
- Adverse events (AEs) including but not limited to:
  - Serious AEs
  - Discontinuation due to AEs
  - Other AEs of interest
    - Hyperkeratosis

- Severe infection
- Pneumonia
- Gum disease

## **Timing**

Evidence on intervention effectiveness and harms will be derived from studies of any duration.

## **Settings**

All relevant settings will be considered, with a focus on outpatient settings in the United States.

## Benefits Beyond Health and Special Ethical Priorities

Our reviews seek to provide information on benefits beyond health and special ethical priorities offered by the intervention to the individual patient, caregivers, the delivery system, other patients, or the public that would not have been considered as part of the evidence on comparative clinical effectiveness. These general elements (i.e., not specific to a given disease) are listed in the table below.

**Table 1.1. Benefits Beyond Health and Special Ethical Priorities**

<b>Benefits Beyond Health and Special Ethical Priorities*</b>
There is substantial unmet need despite currently available treatments.
This condition is of substantial relevance for people from a racial/ethnic group that have not been equitably served by the healthcare system.
The treatment is likely to produce substantial improvement in caregivers' quality of life and/or ability to pursue their own education, work, and family life.
The treatment offers a substantial opportunity to improve access to effective treatment by means of its mechanism of action or method of delivery.

\*Benefits beyond health and special ethical priorities shape to some extent how the value of any effective treatments for a particular condition will be judged and are meant to reflect the broader effects of a specific treatment on patients, caregivers, and society. For additional information, please see the [ICER Value Assessment Framework](#).

ICER encourages stakeholders to provide input on these elements in their public comment submissions.

## Scope of Comparative Value Analyses

A detailed economic model analysis plan with proposed methodology, model structure, model parameters, model inputs, and model assumptions will be published on June 6, 2025. This scoping document provides early thoughts about the overall model structure.

As a complement to the evidence review, we will develop a *de-novo* economic model to assess the lifetime cost-effectiveness of brensocatib as an add-on therapy to usual care, which may include antibiotics, mucolytics, pulmonary rehabilitation, and airway clearance devices, compared with usual care alone. The model structure will be developed based in part on models identified through a targeted systematic literature review of economic analyses of treatments for bronchiectasis.<sup>19-22</sup>

Analyses will be conducted from the health care system perspective including direct medical care costs only. Patient and caregiver productivity, and other indirect costs will be considered in a modified societal perspective analysis. The modified societal perspective will be considered as a co-base case when (a) direct data on indirect costs are available, (b) the societal costs of care are large relative to direct health care costs, and (c) the impact of treatment on the indirect costs is substantial. This will most often occur in cases where the incremental cost-effectiveness ratio

changes by greater than 20%, greater than \$200,000 per QALY, and/or when the result crosses the threshold of \$100,000-\$150,000 per QALY gained. If direct data are lacking on patient and/or caregiver productivity, we will implement a method to capture the potential impacts of brensocatib on productivity (patient and caregiver).

The target population will consist of adolescents and adults with NCFB. The analytic cohort consists of patients who previously had at least two exacerbations in the previous year. The model will consist of seven Markov health states: 1) NCFB without exacerbation; 2) NCFB with exacerbation; 3) NCFB with chronic *Pseudomonas aeruginosa* infection without exacerbation; 4) NCFB with chronic *Pseudomonas aeruginosa* infection with exacerbation; 5) NCFB with chronic nontuberculous mycobacterial (NTM) infection without exacerbation; 6) NCFB with chronic NTM infection with exacerbation; and 7) Death. Provided that a significant proportion of exacerbation requires hospital admission and short-term extensive care, a stand-alone inpatient admission state may be added to the model as a transient state. A cohort of patients will be modeled as they transition between states during predetermined cycles (i.e., monthly) over a lifetime time horizon, modeling patients from treatment initiation until death over a lifetime time horizon, modeling patients from treatment initiation until death. Note that worsening NCFB is not a feature of the current model. If there is sufficient evidence supporting an effect of brensocatib on the progressive nature of NCFB, we will explore the potential impact of treatment with brensocatib on worsening NCFB in scenario analyses.

Key model inputs will include clinical probabilities, quality of life values, and health care costs. Probabilities, costs, and other inputs will differ to reflect varying effectiveness between the intervention and usual care. Treatment effectiveness will be estimated using data from clinical trials for both the treatment and comparator.<sup>23,24</sup> We will also consider expert opinion as the source of model inputs when insufficient evidence exists identifying those inputs from published literature.

Health outcomes and costs will be dependent on time spent in each health state, clinical events, adverse events (AEs), and direct medical costs. The health outcome of each intervention will be evaluated in terms of pulmonary exacerbations avoided, life-years gained, quality-adjusted life years (QALYs) gained, and equal value of life years gained ([evLYG](#)). Quality of life weights will be applied to each health state, including quality of life decrements for serious adverse events. The model will include direct medical costs, including but not limited to costs related to drug administration, drug monitoring, condition-related care, and serious adverse events. In addition, patient and caregiver productivity changes and other indirect costs will be included in a separate analysis, as available data allow. In separate analyses, we will explore the potential health care system budgetary impact of treatment over a five-year time horizon, utilizing published or otherwise publicly-available information on the potential population eligible for treatment and results from the economic model for treatment costs and cost offsets. This budgetary impact analysis will indicate the relation between treatment prices and level of use for a given potential



budget impact, and will allow assessment of any need for managing the cost of such interventions. More information on ICER's methods for estimating potential budget impact can be found [here](#).

## **Identification of Low-Value Services**

ICER includes in its reports information on wasteful or lower-value services in the same clinical area that could be reduced or eliminated to create additional resources in health care budgets for higher-value innovative services (for more information, see ICER's Value Assessment Framework). These services are ones that would not be directly affected by brensocatib (e.g., reduced need for emergency department visits and hospitalizations), as these services will be captured in the economic model. Rather, we are seeking services used in the current management of NCFB beyond the potential offsets that arise from a new intervention. ICER encourages all stakeholders to suggest services (including treatments and mechanisms of care) that could be reduced, eliminated, or made more efficient.

# References

---

1. Aliberti S, Goeminne PC, O'Donnell AE, et al. Criteria and definitions for the radiological and clinical diagnosis of bronchiectasis in adults for use in clinical trials: international consensus recommendations. *Lancet Respir Med*. Mar 2022;10(3):298-306. doi:10.1016/s2213-2600(21)00277-0
2. Cole PJ. Inflammation: a two-edged sword--the model of bronchiectasis. *Eur J Respir Dis Suppl*. 1986;147:6-15.
3. Polverino E, Goeminne PC, McDonnell MJ, et al. European Respiratory Society guidelines for the management of adult bronchiectasis. *European Respiratory Journal*. 2017;50(3):1700629. doi:10.1183/13993003.00629-2017
4. Flume PA, Chalmers JD, Olivier KN. Advances in bronchiectasis: endotyping, genetics, microbiome, and disease heterogeneity. *Lancet*. Sep 8 2018;392(10150):880-890. doi:10.1016/s0140-6736(18)31767-7
5. Hurst JR, Elborn JS, Soyza AD. COPD-bronchiectasis overlap syndrome. *European Respiratory Journal*. 2015;45(2):310-313. doi:10.1183/09031936.00170014
6. Dupont L. Lost in translation? Therapeutic contrasts in CF and non-CF bronchiectasis. *Pneumologie*. 2016/10/11 2016;70(10):A5. doi:10.1055/s-0036-1592229
7. Weycker D, Hansen GL, Seifer FD. Prevalence and incidence of noncystic fibrosis bronchiectasis among US adults in 2013. *Chron Respir Dis*. Nov 2017;14(4):377-384. doi:10.1177/1479972317709649
8. Weycker D, Edelsberg J, Oster G, Tino G. Prevalence and Economic Burden of Bronchiectasis. *Clinical Pulmonary Medicine*. 2005;12(4):205-209. doi:10.1097/01.cpm.0000171422.98696.ed
9. Seitz AE, Olivier KN, Adjemian J, Holland SM, Prevots DR. Trends in Bronchiectasis Among Medicare Beneficiaries in the United States, 2000 to 2007. *CHEST*. 2012;142(2):432-439. doi:10.1378/chest.11-2209
10. Roberts JM, Goyal V, Kularatna S, et al. The Economic Burden of Bronchiectasis: A Systematic Review. *CHEST*. 2023;164(6):1396-1421. doi:10.1016/j.chest.2023.06.040
11. Shoab S, Feliciano J, Dasenbrook EC, et al. Real-world disease burden, mortality, and healthcare resource utilization associated with bronchiectasis. *Chronic Respiratory Disease*. 2025;22:14799731241310897. doi:10.1177/14799731241310897
12. Chalmers JD, Goeminne P, Aliberti S, et al. The bronchiectasis severity index. An international derivation and validation study. *Am J Respir Crit Care Med*. Mar 1 2014;189(5):576-85. doi:10.1164/rccm.201309-1575OC
13. Hill AT, Sullivan AL, Chalmers JD, et al. British Thoracic Society Guideline for bronchiectasis in adults. *Thorax*. Jan 2019;74(Suppl 1):1-69. doi:10.1136/thoraxjnl-2018-212463
14. FDA Grants Priority Review to Insmed's Brensocatib for Treatment of Bronchiectasis with PDUFA Target Action Date Set for August 12, 2025. 2025. <https://investor.insmed.com/2025-02-06-FDA-Grants-Priority-Review-to-Insmeds-Brensocatib-for-Treatment-of-Bronchiectasis-with-PDUFA-Target-Action-Date-Set-for-August-12,-2025>
15. Delestre-Levai I, Aliberti S, Almagro M, et al. Patients' perspectives on bronchiectasis: findings from a social media listening study. *ERJ Open Research*. 2021;7(3):00096-2021. doi:10.1183/23120541.00096-2021
16. Dudgeon EK, Crichton M, Chalmers JD. "The missing ingredient": the patient perspective of health related quality of life in bronchiectasis: a qualitative study. *BMC Pulmonary Medicine*. 2018/05/22 2018;18(1):81. doi:10.1186/s12890-018-0631-7
17. Magge A, Ashraf S, Quittner AL, Metersky ML. Quality of life in patients with bronchiectasis: a 2-year longitudinal study. *Ann Transl Med*. Jul 2019;7(14):334. doi:10.21037/atm.2019.06.62

18. European Lung Foundation. Bronchiectasis Patient Perspectives (ERS Congress 2021). <https://europeanlung.org/en/people-and-partners/your-experiences/bronchiectasis-patient-perspectives/>
19. Bhattacharyya S.B. CF, Priedane E., Shirore R.M., Haworth C.S., Flume P.A., Sonathi V., Thomas S.K. A De-Novo ecoNoMic MoDel to Assess cliNicAl AND ecoNoMic coNsequeNces of BRoNchiectAsis. 2014:PA551.
20. Milne RJ, Hockey H, Rea H. Long-term air humidification therapy is cost-effective for patients with moderate or severe chronic obstructive pulmonary disease or bronchiectasis. *Value in health : the journal of the International Society for Pharmacoeconomics and Outcomes Research*. Jun 2014;17(4):320-7. doi:10.1016/j.jval.2014.01.007
21. van Wilder P, Odnoletkova I, Mouline M, de Vries E. Immunoglobulin Replacement Therapy is critical and cost-effective in increasing life expectancy and quality of life in patients suffering from Common Variable Immunodeficiency Disorders (CVID): A health-economic assessment. *PLoS One*. 2021;16(3):e0247941. doi:10.1371/journal.pone.0247941
22. Windegger TM, Nghiem S, Nguyen KH, Fung YL, Scuffham PA. Primary immunodeficiency disease: a cost-utility analysis comparing intravenous vs subcutaneous immunoglobulin replacement therapy in Australia. *Blood Transfus*. Mar 2020;18(2):96-105. doi:10.2450/2019.0083-19
23. Insmed Incorporated. ASPEN Topline Results. 2024.
24. Chalmers JD, Haworth CS, Metersky ML, et al. Phase 2 Trial of the DPP-1 Inhibitor Brensocatib in Bronchiectasis. *N Engl J Med*. Nov 26 2020;383(22):2127-2137. doi:10.1056/NEJMoa2021713